

ASYMPTOTIC THEORY OF CEPSTRAL RANDOM FIELDS

BY TUCKER S. MCELROY[†] AND SCOTT H. HOLAN^{*,‡}

U.S. Census Bureau[†] and University of Missouri[‡]

Random fields play a central role in the analysis of spatially correlated data and, as a result, have a significant impact on a broad array of scientific applications. This paper studies the cepstral random field model, providing recursive formulas that connect the spatial cepstral coefficients to an equivalent moving-average random field, which facilitates easy computation of the autocovariance matrix. We also provide a comprehensive treatment of the asymptotic theory for two-dimensional random field models: we establish asymptotic results for Bayesian, maximum likelihood, and quasi-maximum likelihood estimation of random field parameters and regression parameters. The theoretical results are presented generally and are of independent interest, pertaining to a wide class of random field models. The results for the cepstral model facilitate model-building: because the cepstral coefficients are unconstrained in practice, numerical optimization is greatly simplified, and we are always guaranteed a positive definite covariance matrix. We show that inference for individual coefficients is possible, and one can refine models in a disciplined manner. Our results are illustrated through simulation and the analysis of straw yield data in an agricultural field experiment.

1. Introduction. Spatial data feature heavily in many scientific disciplines including ecology, environmental science, epidemiology, geography, geology, small area estimation, and socio-demographics. Although spatial data can be broadly placed into three categories: *geostatistical data*, *lattice data*, and *spatial patterns* [11], our focus mainly resides in the development of cepstral random field models for spatial lattice data. That is, we consider random fields where the index set for the variables is \mathbb{Z}^2 – appropriate for image processing, for example.

Research on spatial random fields dates back over half a century, e.g., see Whittle [42]. Other references on spatial random fields include Besag [3, 4], Guyon [18], Rosenblatt [33], Besag and Green [5], and Rosenblatt

*Supported by NSF and the U.S. Census Bureau under NSF grant SES-1132031, through the NSF-Census Research Network (NCRN) program and by a University of Missouri Research Board grant.

MSC 2010 subject classifications: Primary 62F12, 62M30; Secondary 62F15, 62M10

Keywords and phrases: Bayesian estimation, Cepstrum, Exponential spectral representation, Lattice data, Spatial statistics, Spectral density.

[34], among others. Comprehensive overviews can be found in Cressie [11], Stein [38], Banerjee, Carlin and Gelfand [2], Cressie and Wikle [12], and the references therein. Recently, there has been a growing interest in modeling spatial random fields through the spectral domain. For example, see Fuentes [14], Fuentes, Guttorp and Sampson [15], Tonellato [41], Fuentes and Reich [16], Bandyopadhyay and Lahiri [1], and the references therein.

For a stationary Gaussian random field, it is natural to impose a Markov structure, as described in Rue and Held [35], in order to obtain an inverse covariance matrix (i.e., a precision matrix) that has a sparse structure, because this will ensure speedy computation of maximum likelihood estimates. Rue and Held [35] show how careful specification of conditional distributions generates a well-defined random field. However, this technique relies upon imposing *a priori* a sparse structure on the precision matrix, i.e., demanding that many conditional precisions be zero. In contrast, the cepstral random field does not generate a sparse covariance (or precision) matrix, and yet always yields a well-defined spatial random field; this occurs because the model is formulated in the frequency domain, by ensuring a positive spectral density. This frequency-domain approach provides a general way of specifying a non-isotropic random field, which is useful when we do not have prior notions about conditional variances or precisions.

The cepstral random field allows for unconstrained optimization of the objective function, i.e., each model coefficient can be any real number independently of the others; this appealing property is in marked contrast to other models and approaches, such as moving averages or Markov random fields (these require constraints on parameters to achieve identifiability and/or a well-defined process). In the development of this model, Solo [37] presents estimation approaches by both log periodogram regression and Whittle maximum likelihood, but does not derive the asymptotic properties of estimators. Based on information criterion, Mallows' C_p , and hypothesis testing, the author briefly describes methods for model selection. Some key advantages of the cepstral model are that it is well-defined (because it is defined through the spectral density), it is identifiable, and the cepstral parameter estimates are asymptotically uncorrelated with one another.

This paper provides a first comprehensive treatment of the theory for cepstral random field models. In particular, we establish recursive formulas for connecting cepstral random fields to moving average random fields, thus facilitating efficient computation of the spatial autocovariances, which are needed for likelihood evaluation and prediction. Critically, the resulting autocovariance matrix is guaranteed to be positive-definite; note that if we were to work with a moving average (MA) field instead, it would not be

identifiable without imposing further complicated parameter restrictions.

Additionally, we develop asymptotic results for Bayesian, maximum likelihood, and quasi-maximum likelihood estimation of field parameters and regression parameters under an expanding domain formulation. In particular, we establish asymptotic consistency in both the Bayesian and likelihood settings and provide Central Limit Theorems for the frequentist estimators we propose. We discuss the computational advantages of the cepstral model, and propose an exact Whittle likelihood that avoids the burdensome inversion of the autocovariance matrix. Although our primary focus is on cepstral models, the theoretical developments are presented for general random field models with regression effects. Our results are of independent interest and extend the existing results of Mardia and Marshall [24], providing a rigorous framework for conducting model building and inference under an expanding domain framework; this is applicable to lattice random field data that is sampled at regular fixed intervals, and for which in-filling is either impractical or of little interest.

As discussed in Sections 2 and 3, the proposed cepstral models are computationally advantageous over many current models (e.g., spatial autoregressive models), because no constraints need to be imposed on the parameters to ensure the resulting autocovariance matrix remains positive definite. In fact, given the recursive formulas of Section 2, one can model the two-dimensional cepstral coefficients (i.e., the Fourier coefficients of the two-dimensional log spectrum) and arrive at the autocovariances without the need for direct Fourier inversion.

Since the model's first inception [37], the cepstral random field literature has remained sparse, with relatively few examples to date. For example, Cressie [11, p. 448] makes brief mention of the model. In a different context, Noh and Solo [28] use cepstral random fields to test for space-time separability. Sandgren and Stoica [36] use two-dimensional cepstrum thresholding models to estimate the two-dimensional spectral density. However, this work does not treat the random field case. Related to our work, Kizilkaya and Kayran [22] derive an algorithm for computing cepstral coefficients from a known ARMA random field, whereas Kizilkaya [21] provides a recursive formula for obtaining a nonsymmetric half plane MA random field models for a given cepstral specification. In contrast, our recursive formulas provide unrestricted MA random fields as well as the necessary autocovariances for expressing the Gaussian likelihood.

This paper proceeds as follows. Section 2 describes the cepstral model and its computation. Specifically, this section lays out the recursive formulas that are needed to estimate the autocovariances given the cepstral coefficients.

Section 3 details the different model fitting methods, including Bayesian, maximum likelihood, quasi-maximum likelihood, and exact Whittle likelihood. Our theoretical results are provided in Section 4. Here, we establish consistency and asymptotic normality of the proposed estimators. Section 5 illustrates the models effectiveness through a simulation study, and Section 6 contains concluding discussion. Extensions to missing data, imputation, and signal extraction along with an application of our methodology to straw yield data from an agricultural experiment, as well as all proofs, are provided in a Supplementary Appendix.

2. The Cepstral Model and its Computation. We begin by introducing some basic concepts about spatial random fields, and then we specialize to the cepstral random field, with a focus on computation of auto-covariances. References on spatial random fields include Whittle [42], Besag [3], Rosenblatt [33, 34], Solo [37], Cressie [11], Kedem and Fokianos [20], and Rue and Held [35]. A random field $\mathbb{Y} = \{\mathbb{Y}_{s_1, s_2}\}$ is a process with indices on a lattice, which in this paper we take to be \mathbb{Z}^2 . Typically a random field has a mean function $\mu_{s_1, s_2} = \mathbb{E}\mathbb{Y}_{s_1, s_2}$, which may be modeled through regression variables (Cressie, 1993). The mean-corrected field $\mathbb{Y} - \{\mu_{s_1, s_2}\}$ will be denoted by \mathbb{W} .

Interest focuses upon weakly stationary random fields, which in practice is often adequate once mean effects are identified and accounted for. When all moments are defined, this is equivalent to the higher cumulants [8] being dependent only on lags between the mean-centered variables. The second cumulant function, or autocovariance function (acf), is defined via $\text{Cov}(\mathbb{Y}_{s_1, s_2}, \mathbb{Y}_{r_1, r_2}) = \mathbb{E}[\mathbb{W}_{s_1, s_2} \mathbb{W}_{r_1, r_2}] = \gamma_{s_1 - r_1, s_2 - r_2}$ for all $s_1, s_2, r_1, r_2 \in \mathbb{Z}$. It is convenient to summarize this second-order structure through the spectral density F defined on $[-\pi, \pi]^2$, which depends on two frequencies. Letting $Z_j = e^{-i\lambda_j}$ for $j = 1, 2$, the spectral density is related to the acf via the formula

$$(2.1) \quad F(\lambda_1, \lambda_2) = \sum_{h_1, h_2 \in \mathbb{Z}} \gamma_{h_1, h_2}(F) Z_1^{h_1} Z_2^{h_2}.$$

Here we write $\gamma(F)$ for the acf associated with the spectrum F , and it in turn is expressed in terms of F via Fourier inversion as

$$(2.2) \quad \gamma_{h_1, h_2}(F) = \frac{1}{4\pi^2} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} F(\lambda_1, \lambda_2) Z_1^{-h_1} Z_2^{-h_2} d\lambda_1 d\lambda_2.$$

As a general notation, let the normalized double integral over both frequencies be abbreviated by the expression $\langle \cdot \rangle$, so that $\gamma_{h_1, h_2}(F) = \langle F Z_1^{-h_1} Z_2^{-h_2} \rangle$

is compactly expressed. Now it follows elementarily from the commutativity of the field \mathbb{Y} variables that $\gamma_{h_1, h_2}(F) = \gamma_{-h_1, -h_2}(F)$, and hence the corresponding F in (2.1) must have mirror reflectional symmetry through both axes, i.e., $F(\lambda_1, \lambda_2) = F(-\lambda_1, -\lambda_2)$. Furthermore, the acf of a random field is always positive-definite [11] and the corresponding spectrum is non-negative [7].

2.1. The Cepstral Random Field Model. A spatial model for continuous-valued random variables should, at a minimum, capture second-order structure in the data, which is summarized through the acf. However, a putative acf may or may not have non-negative Discrete Fourier Transform (DFT) (2.1), whereas any valid acf of a stationary field *must* have non-negative spectrum F . One way to ensure our model has such a valid acf is to model F , utilizing some class of non-negative functions, and determine the corresponding covariances via (2.2). This is the philosophy behind the versatile exponential time series model of Bloomfield [6]. The idea there was to expand the log spectrum in the complex exponential basis functions, with a truncation of the expansion corresponding to a postulated model.

The same idea is readily adapted to the spatial context; Solo [37] seems to be the first formal presentation of this idea. Given that F is strictly positive and bounded, we can expand $\log F$ in each frequency concurrently, which yields

$$\log F(\lambda_1, \lambda_2) = \sum_{j_1, j_2 \in \mathbb{Z}} \Theta_{j_1, j_2} Z_1^{j_1} Z_2^{j_2}.$$

The coefficients $\{\Theta_{j_1, j_2} = \langle \log F Z_1^{-j_1} Z_2^{-j_2} \rangle\}$ are called the cepstral coefficients; see also the recent treatment of Kizilkaya and Kayran [22]. A pleasing feature of this representation is that F^{-1} has cepstral coefficients $\{-\Theta_{j_1, j_2}\}$. By truncating the summation, we obtain a parametric model that can approximate the second-order structure of *any* random field with bounded spectrum. So we obtain the cepstral model of order (p_1, p_2) given by

$$(2.3) \quad F(\lambda_1, \lambda_2) = \exp \left\{ \sum_{j_1=-p_1}^{p_1} \sum_{j_2=-p_2}^{p_2} \Theta_{j_1, j_2} Z_1^{j_1} Z_2^{j_2} \right\}.$$

Note that the cepstral coefficient $\Theta_{0,0}$ has no sinusoidal function multiplying it, and hence $\exp \Theta_{0,0}$ quantifies the scale of the data. In one dimension, this would be called the innovation variance; note that $\Theta_{0,0} = \langle \log F \rangle$. Because the complex exponentials form a complete orthonormal basis set, it is impossible for two distinct values of Θ to produce an identical function F ; hence the model is identifiable. Further special cases of the general cepstral

field model are considered in Solo [37]. Because F has mirror reflectional symmetry, the cepstral coefficients do as well, i.e., $\Theta_{j_1, j_2} = \Theta_{-j_1, -j_2}$.

In order to fit this model to Gaussian data, it is necessary to compute the acf from a given specification of cepstral coefficients. We next describe two approaches to this: one is approximate, and the other is exact. Both differ from the fitting techniques in Solo [37], who advocates an asymptotic likelihood (or Whittle) calculation.

2.2. Fast Calculation of Autocovariances. We here discuss a straightforward discretization of (2.2), together with (2.3), utilizing the Riemann approximation. So long as the spectrum is a bounded function, this method is arbitrarily accurate (since the practitioner controls the mesh size). In order to accomplish the computation, without loss of generality let $p_2 = p_1$, so that the cepstral coefficients are given by a $(2p_1 + 1) \times (2p_1 + 1)$ grid Θ (if $p_2 < p_1$, just fill in some entries of Θ with zeroes).

Now we refer to the entries of Θ via Θ_{j_1, j_2} with $-p_1 \leq j_1, j_2 \leq p_1$, which is a Cartesian mode of indexing; this differs from the style of indexing pertinent to matrices. We can map this grid to a matrix $[\Theta]$ (and back), with the following rule:

$$(2.4) \quad [\Theta]_{k_1, k_2} = \Theta_{k_2 - p_1 - 1, p_1 + 1 - k_1} \quad \Theta_{j_1, j_2} = [\Theta]_{p_1 + 1 - j_2, j_1 + p_1 + 1}$$

for $1 \leq k_1, k_2 \leq 2p_1 + 1$ and $-p_1 \leq j_1, j_2 \leq p_1$. We will consider a set of frequencies $\{\ell_1\pi/M, \ell_2\pi/M\}$ for $-M \leq \ell_1, \ell_2 \leq M$, which is an order M discretization of $[-\pi, \pi]^2$. Suppose that we wish to compute the grid of autocovariances given by $\Gamma = \{\gamma_{h_1, h_2}\}_{h_1, h_2 = -H}^H$ for some maximal lag H . To that end, we consider a complex-valued $2p_1 + 1 \times 2M + 1$ matrix E with entries $E_{k_1, k_2} = \exp\{i\pi(p_1 + 1 - k_1)(M - k_2 + 1)M^{-1}\}$ for $k_1 = 1, 2, \dots, 2p_1 + 1$ and $k_2 = 1, 2, \dots, 2M + 1$, and also define a $(2H + 1) \times (2M + 1)$ dimensional matrix G via $G_{j_1, j_2} = \exp\{i\pi(H + 1 - j_1)(M + 1 - j_2)M^{-1}\}$. Then with $[\Gamma]$ defined via $[\Gamma]_{k_1, k_2} = \gamma_{k_2 - H - 1, H + 1 - k_1}$, the following formula

$$(2.5) \quad [\Gamma] \cong (2M + 1)^{-2} G \exp\{\bar{E}'[\Theta]E\}\bar{G}'$$

provides a practical method of computation. In this formula, which is derived in the Supplement's Appendix B, we have written the exponential of a matrix, which here is *not* the "matrix exponential," but rather just consists of exponentiating each entry of the matrix. So (2.5) produces an arbitrarily fine approximation to the acf (taking M as large as desired). The algorithm takes a given Θ , produces $[\Theta]$ via (2.4), computes E and G (ahead of time, as they do not depend upon the parameters), and determines $[\Gamma]$ via (2.5).

2.3. *Exact Calculation of Autocovariances.* We now present an exact method for computing the acf from the cepstral matrix. Our approach is similar to that of Section 3 of Kizilkaya and Kayran [22], though with one important difference. They present an algorithm for computing cepstral coefficients from known coefficients of an ARMA random field. Instead, we take the cepstral coefficients as given, compute coefficients of certain corresponding MA random fields, and from there obtain the acf. In order to fit the Gaussian likelihood, we need to compute the acf from the cepstral matrix, not the reverse.

We introduce the device of a “causal” field and a “skew” field as follows. The causal field is a MA field that only involves coefficients with indices in the positive quadrant, whereas the skew field essentially is defined over the second quadrant. More precisely, we have

$$(2.6) \quad \gamma_{s_1, s_2}(\Psi) = \sum_{k_1, k_2 \geq 0} \psi_{s_1+k_1, s_2+k_2} \psi_{k_1, k_2}$$

$$\left| \sum_{j_1, j_2 \geq 0} \psi_{j_1, j_2} Z_1^{j_1} Z_2^{j_2} \right|^2 = \sum_{s_1, s_2 \in \mathbb{Z}} \gamma_{s_1, s_2}(\Psi) Z_1^{s_1} Z_2^{s_2}$$

for the causal field. The causal field may be written formally (in terms of backshift operators B_1, B_2) as $\Psi(B_1, B_2) = \sum_{j_1, j_2 \geq 0} \psi_{j_1, j_2} B_1^{j_1} B_2^{j_2}$. That is, the ψ_{j_1, j_2} coefficients define the moving average representation of the causal field, and $\{\gamma_{s_1, s_2}(\Psi)\}$ is its acf. It is important that we set $\psi_{0,0} = 1$. Similarly, let $\Phi(B_1, B_2) = \sum_{j_1, j_2 \geq 0} \phi_{j_1, j_2} B_1^{-j_1} B_2^{j_2}$ for the skew-field, which in the first index depends on the forward shift operator B_1^{-1} , but on the backshift operator B_2 in the second index. Thus,

$$(2.7) \quad \gamma_{s_1, s_2}(\Phi) = \sum_{k_1, k_2 \geq 0} \phi_{s_1+k_1, s_2+k_2} \phi_{k_1, k_2}$$

$$\left| \sum_{j_1, j_2 \geq 0} \phi_{j_1, j_2} Z_1^{-j_1} Z_2^{j_2} \right|^2 = \sum_{s_1, s_2 \in \mathbb{Z}} \gamma_{s_1, s_2}(\Phi) Z_1^{-s_1} Z_2^{s_2}.$$

We also have two time series, corresponding to the axes of the cepstral matrix, given by $\Xi(B_1) = \sum_{j_1 \geq 0} \xi_{j_1} B_1^{j_1}$ and $\Omega(B_2) = \sum_{j_2 \geq 0} \omega_{j_2} B_2^{j_2}$, which have acfs $\gamma_{h_1}(\Xi) = \sum_{k_1 \geq 0} \xi_{k_1+h_1} \xi_{k_1}$ and $\gamma_{h_2}(\Omega) = \sum_{k_2 \geq 0} \omega_{k_2+h_2} \omega_{k_2}$, respectively. Now each of these MA random fields has a natural cepstral representation, such that their acfs can be combined to produce the cepstral acf, as shown in the following result.

PROPOSITION 2.1. *The acf of the cepstral model is given by*

$$(2.8) \quad \gamma_{h_1, h_2}(F) = e^{\Theta_{0,0}} \sum_{j_1, j_2 \in \mathbb{Z}} \gamma_{j_1, j_2}(\Phi) \left[\sum_{k_1, k_2 \in \mathbb{Z}} \gamma_{h_1+j_1-k_1, h_2-j_2-k_2}(\Psi) \gamma_{k_1}(\Xi) \gamma_{k_2}(\Omega) \right],$$

where $\gamma(\Phi)$, $\gamma(\Psi)$, $\gamma(\Xi)$, and $\gamma(\Omega)$ can be calculated in terms of their coefficients, which are recursively given by

$$(2.9) \quad \psi_{j_1, j_2} = \frac{1}{j_1} \sum_{k_1=1}^{p_1} k_1 \left(\sum_{k_2=1}^{j_2} \psi_{j_1-k_1, j_2-k_2} \Theta_{k_1, k_2} \right)$$

$$(2.10) \quad \phi_{j_1, j_2} = \frac{1}{j_1} \sum_{k_1=1}^{p_1} k_1 \left(\sum_{k_2=1}^{j_2} \phi_{j_1-k_1, j_2-k_2} \Theta_{-k_1, k_2} \right)$$

$$(2.11) \quad \xi_{j_1} = \frac{1}{j_1} \sum_{k_1=1}^{p_1} k_1 \Theta_{k_1, 0} \xi_{j_1-k_1}$$

$$(2.12) \quad \omega_{j_2} = \frac{1}{j_2} \sum_{k_2=1}^{p_1} k_2 \Theta_{0, k_2} \omega_{j_2-k_2},$$

for $j_1 \geq 1$ and $j_2 \geq 1$.

Proposition 2.1 gives recursive formulas. In the causal case, one would compute $\psi_{1,1}, \psi_{2,1}, \dots, \psi_{p_1,1}, \psi_{1,2}, \psi_{2,2}, \dots$, etc. Alternative computational patterns could be utilized, noting that ψ_{j_1, j_2} only requires knowledge of ψ_{ℓ_1, ℓ_2} with $\ell_1 < j_1$ and $\ell_2 < j_2$. When $p_1 = \infty$, equation (2.8) gives the precise mapping of cepstral coefficients to various MA coefficients, and ultimately to the autocovariance function. If $p_1 < \infty$, it provides an algorithm for determining autocovariances for a given cepstral model. These formulas are already much more complicated than in the time series case [see 32], and for higher dimensional fields become intractable.

3. Model Fitting Methods. In this section we give additional details on various methods for fitting cepstral random field models, and present some tools for refining specified models. Once a model is specified, we can estimate the parameters via exact maximum likelihood, Bayesian posterior simulation, an approximate Whittle likelihood, or an exact Whittle likelihood. We focus on these four techniques due to their mixture of being flexible and possessing good statistical properties.

We first define Kullback-Leibler (KL) discrepancy, the exact Whittle likelihood, and the quasi-maximum likelihood estimate (QMLE), and then we

proceed to describe the distributional behavior of the maximum likelihood estimates (MLEs) and QMLEs, extending the results of Mardia and Marshall [24] to non-Gaussian fields, under an expanding domain asymptotic theory. These results, proved for fairly general linear random fields with regression effects, are then specialized to the case of the cepstral field, and model selection is afterwards described.

3.1. *Random Field Data.* Now we proceed to discuss spatial modeling (here we do not assume a cepstral random field structure), adapting the vector time series treatment in Taniguchi and Kakizawa [39]. Suppose that our data comes to us in gridded form, corresponding to a $N_1 \times N_2$ matrix \mathbb{Y}^N (with \mathbb{W}^N denoting the de-meaned version). We use the notation $N = \sqrt{N_1 \cdot N_2}$, so that N^2 is the sample size. Both \mathbb{Y}^N and \mathbb{W}^N can be vectorized into length N^2 vectors Y and W via the so-called lexicographical rule

$$Y_k = \mathbb{Y}_{s_1, s_2}^N, \quad W_k = \mathbb{W}_{s_1, s_2}^N, \quad k = N_2(s_1 - 1) + s_2.$$

Here, $Y = \text{vec}(\mathbb{Y}^{N'})$, where vec stands for the vector operation on a matrix, and $'$ is the transpose. Note that $s_1 - 1 = k \text{ div } N_2$ and $s_2 = k \text{ mod } N_2$. Also let $\mu = \mathbb{E}Y$, so that $\mu_k = \mathbb{E}Y_k = \mathbb{E}\mathbb{Y}_{s_1, s_2}^N = \mu_{s_1, s_2}$. In the simplest scenario the mean matrix $\{\mu_{s_1, s_2}\}$ is constant with respect to the indices s_1, s_2 . More generally, we might model the mean through regressor functions defined upon the grid, i.e., $\mu_{s_1, s_2} = \sum_{\ell=1}^L \beta_\ell X_\ell(s_1, s_2)$ for some specified lattice functions $\{X_\ell\}_{\ell=1}^L$. Then

$$\mu_k = \sum_{\ell=1}^L \beta_\ell X_\ell(k \text{ div } N_2 + 1, k \text{ mod } N_2) = \sum_{\ell=1}^L \beta_\ell \tilde{X}_\ell(k)$$

maps each X_ℓ from a lattice function to a function \tilde{X}_ℓ of the natural numbers. The parameters $\beta_1, \beta_2, \dots, \beta_L$ then enter the regression linearly, and we can express things compactly via $\mu = \tilde{X}\beta$, where \tilde{X} is the regression matrix with columns given by the various \tilde{X}_ℓ .

The spectral density of a mean zero random field \mathbb{W} has already been defined in (2.1), and the DFT of the field is now defined as

$$\tilde{\mathbb{W}}(\lambda_1, \lambda_2) = \sum_{t_1=1}^{N_1} \sum_{t_2=1}^{N_2} \mathbb{W}_{t_1, t_2}^N e^{-i\lambda_1 t_1} e^{-i\lambda_2 t_2} = \sum_{t_1=1}^{N_1} \sum_{t_2=1}^{N_2} W_{N_2(t_1-1)+t_2} Z_1^{t_1} Z_2^{t_2}$$

for $\lambda_1, \lambda_2 \in [-\pi, \pi]$. Note that we define this DFT over all pairs of frequencies, not just at the so-called Fourier frequencies. Also the DFT depends on β through the mean-centering; if we center the data \mathbb{Y}^N by any regression

parameter other than the true β , denoted $\tilde{\beta}$, some bias will be introduced. The periodogram will be defined at all frequencies, and is proportional to the squared magnitude of the DFT:

$$I_{\tilde{\beta}}(\lambda_1, \lambda_2) = N^{-2} |\widehat{\mathbb{W}}(\lambda_1, \lambda_2)|^2 = \sum_{|h_1| < N_1} \sum_{|h_2| < N_2} \gamma_{h_1, h_2}(I_{\tilde{\beta}}) Z_1^{h_1} Z_2^{h_2}.$$

Here $\gamma_{h_1, h_2}(I_{\beta})$ is defined as the sample acf of the series demeaned by $\mu = \tilde{X}\beta$ (see Appendix B for more detail); moreover it satisfies (2.2) with F replaced by I_{β} . We also will consider an unbiased acf estimate given by

$$\widehat{\gamma}_{h_1, h_2}(I_{\beta}) = \frac{N^2}{(N_1 - |h_1|)(N_2 - |h_2|)} \gamma_{h_1, h_2}(I_{\beta}).$$

We emphasize that the computation of this periodogram requires a choice of β , and so is written I_{β} . This can be used to assess the frequency domain information in the random field along any row or column; the periodogram can also be viewed as a crude estimate of the spectral density F [11].

In our context the treatment of the periodogram differs from the treatment provided in Fuentes [14]. In particular, we consider the periodogram defined at all frequencies, not just the Fourier frequencies. Additionally, the asymptotic properties developed in Fuentes [14] rely on shrinking domain asymptotics, whereas our asymptotic arguments rely on an expanding domain. Lastly, our periodogram is defined in terms of a mean centered random field and, thus, explicitly depends on the regression parameters β .

3.2. Model Fitting Criteria. Let the covariance matrix of \mathbb{W}^N be denoted $\Sigma(\tilde{F})$, which is defined via $\Sigma(\tilde{F}) = \mathbb{E}\mathbb{W}\mathbb{W}'$; the resulting block-Toeplitz structure of this matrix is analyzed in Section 4. The entries of this matrix can be determined from \tilde{F} via the algorithms of Section 2, along with careful bookkeeping. A model for the data involves a spectrum F_{θ} – let the associated block-Toeplitz covariance matrix be denoted $\Sigma(F_{\theta})$ – which is hoped to be a suitable approximation to $\Sigma(\tilde{F})$. Model fitting can be performed and assessed through the Kullback-Leibler (KL) discrepancy, just as with time series. Although KL is mentioned in Solo [37] and Cressie [11], we provide an in-depth treatment here – see Lemma 4.2, for example. If F and G are two (mean zero) random field spectral densities, their KL discrepancy is defined to be

$$\text{KL}(F, G) = \langle \log F + G/F \rangle.$$

This is a convenient mechanism, since KL is convex in F . As β parametrizes mean effects, we let θ be a parameter vector describing the second-order

structure. If the true data process has spectrum \tilde{F} , and we utilize a model with spectrum F_θ , then $\text{KL}(F_\theta, \tilde{F})$ can be used to assess proximity of the model to truth. The convexity of KL guarantees that when the model is correctly specified, the true parameter $\tilde{\theta}$ minimizes the discrepancy. When the model is misspecified, the minima $\tilde{\theta}$ are called pseudo-true values [cf. 39]. For the cepstral model, the parameter vector is $\theta = J \text{vec}\Theta$, where J is a selection matrix that eliminates redundancies in Θ due to symmetry. The full parameter vector is written ϕ , where $\phi' = [\theta', \beta']$.

It is natural to use KL to fit models as well. For this, consider $\text{KL}(F_\theta, I_\beta)$ – which is called the exact Whittle likelihood – and minimize with respect to θ , which produces by definition the estimate $\hat{\theta}_{QMLE}$. Then using (2.2) we obtain the practical expression

$$\text{KL}(F_\theta, I_\beta) = \langle \log F_\theta \rangle + \sum_{|h_1| < N_1} \sum_{|h_2| < N_2} \gamma_{h_1, h_2}(I_\beta) \cdot \gamma_{h_1, h_2}(F_\theta^{-1}).$$

This assumes that the correct regression parameters have been specified. In the case that F_θ is a cepstral spectrum (2.3), the above expression is even easier to compute: $\langle \log F_\theta \rangle = \Theta_{0,0}$ and $\gamma(F_\theta^{-1}) = \gamma(F_{-\theta})$, i.e., multiply each cepstral coefficient by -1 to obtain the acf of F_θ^{-1} from the acf of F_θ .

Unfortunately, $\gamma_{h_1, h_2}(I_{\tilde{\beta}})$ is biased as an estimate of $\gamma_{h_1, h_2}(\tilde{F})$, and this has a nontrivial impact for spatial data, though not for time series. Essentially, the presence of “corners” in the observed data set reduces the number of data points that are separated by a given lag (h_1, h_2) ; if either of $|h_1|$ or $|h_2|$ is large, we have a very biased estimate. Note, the impact of corners can be visualized by comparing the volume of a d -dimensional cube with that of an inscribed ball; the ratio is $\pi/(2d)$ for $d \geq 2$, which tends to zero as d increases. Thus, corners increasingly dominate the region as d increases, which interferes with one’s ability to measure correlation as a function of lag. This effect is more pronounced as the dimension increases. For this reason, we propose using $\hat{\gamma}_{h_1, h_2}(I_\beta)$ instead of $\gamma_{h_1, h_2}(I_\beta)$, because $\mathbb{E}\hat{\gamma}_{h_1, h_2}(I_{\tilde{\beta}}) = \gamma_{h_1, h_2}(\tilde{F})$. Let us call the modified $\text{KL}(F_\theta, I_\beta)$ by $\widehat{\text{KL}}(F_\theta)$:

$$\widehat{\text{KL}}(F_\theta) = \langle \log F_\theta \rangle + \sum_{|h_1| < N_1} \sum_{|h_2| < N_2} \hat{\gamma}_{h_1, h_2}(I_\beta) \cdot \gamma_{h_1, h_2}(F_\theta^{-1}).$$

Using this criterion instead will produce asymptotically normal cepstral parameter estimates, and therefore is to be preferred.

A drawback of utilizing $\hat{\gamma}_{h_1, h_2}(I_\beta)$ is that the corresponding spectral estimate – the DFT of the unbiased sample acf – need not be positive at all frequencies. Although this is irrelevant asymptotically, in finite samples it

can interfere with inference. The time domain representation of $\widehat{\text{KL}}(F_\theta)$ can still be computed, of course; but the second term in its formula might not be positive. Other types of autocovariance estimators could be utilized, being based on other kinds of spectral estimators (see Politis and Romano [31] for a discussion of the tradeoff between bias and non-negativity of the spectral estimate). These alternative estimators might be based on convolving the periodogram with a spectral window, or equivalently by using a taper (or lag window) with the sample acf. Tapers are known to modify the bias and variance properties of spectral estimators in time series – see Guyon [18], Dahlhaus and Künsch [13], and Politis and Romano [30].

If even faster computation of the objective function is desired, we may discretize $\widehat{\text{KL}}$ and utilize values of F directly, without having to compute the inverse DFT $\gamma(F_\theta^{-1})$. The result is the approximate Whittle likelihood, denoted $\widehat{\text{KL}}_N$, and is obtained by discretizing the integrals in $\text{KL}(F_\theta, I_\beta)$ with a mesh corresponding to Fourier frequencies, but replacing I_β with the DFT of the $\widehat{\gamma}_{h_1, h_2}(I_\beta)$ sequence, denoted by \widehat{I}_β . Then the discrepancy is

$$\widehat{\text{KL}}_N(F_\theta) = N^{-2} \sum_{j_1=-N_1}^{N_1} \sum_{j_2=-N_2}^{N_2} \left\{ \log F_\theta \left(\frac{\pi j_1}{N_1}, \frac{\pi j_2}{N_2} \right) + \frac{\widehat{I}_\beta \left(\frac{\pi j_1}{N_1}, \frac{\pi j_2}{N_2} \right)}{F_\theta \left(\frac{\pi j_1}{N_1}, \frac{\pi j_2}{N_2} \right)} \right\},$$

which can be minimized with respect to θ . The resulting estimate has asymptotic properties identical to the QMLE, and in practice one may use either $\widehat{\text{KL}}$ or $\widehat{\text{KL}}_N$ according to computational convenience. It will be convenient to present a notation for this double discrete sum, which is a Fourier approximation to $\langle \cdot \rangle$, denoted by $\langle \cdot \rangle_N$; then $\widehat{\text{KL}}_N(F_\theta) = \langle \log F_\theta + \widehat{I}_\beta / F_\theta \rangle_N$.

We can also extend the KL formula to handle regression effects:

$$(3.1) \quad \text{KL}(F_\theta, I_\beta) = \langle \log F_\theta \rangle + N^{-2} (Y - \widetilde{X}\beta)' \Sigma(F_\theta^{-1}) (Y - \widetilde{X}\beta).$$

This formula is proved in Appendix B. We propose using (3.1) to estimate regression parameters, but θ is to be determined by $\widehat{\text{KL}}$. The formula for the regression QMLE is then

$$(3.2) \quad \widehat{\beta}_{QMLE} = \left[\widetilde{X}' \Sigma(F_{\widehat{\theta}_{QMLE}}^{-1}) \widetilde{X} \right]^{-1} \widetilde{X}' \Sigma(F_{\widehat{\theta}_{QMLE}}^{-1}) Y,$$

where $\widehat{\theta}_{QMLE}$ minimizes $\widehat{\text{KL}}(F_\theta)$, which in turn depends upon $\widehat{\beta}_{QMLE}$ through \widehat{I}_β . These formulas do not apply when we use the approximate Whittle, although the same asymptotic properties will hold as for the exact Whittle.

On the other hand, we can also compute the exact Gaussian likelihood for the field. The log Gaussian likelihood is equal (up to constants) to

$$(3.3) \quad \mathcal{L}(\theta, \beta) = -\frac{1}{2} \log |\Sigma(F_\theta)| - \frac{1}{2} (Y - \tilde{X}\beta)' \Sigma^{-1}(F_\theta) (Y - \tilde{X}\beta).$$

Maximizing this function with respect to θ yields the MLE $\hat{\theta}_{MLE}$; also $\hat{\beta}_{MLE}$ is given by the Generalized Least Squares (GLS) estimate by standard arguments [see 24]:

$$(3.4) \quad \hat{\beta}_{MLE} = \left[\tilde{X}' \Sigma^{-1}(F_{\hat{\theta}_{MLE}}) \tilde{X} \right]^{-1} \tilde{X}' \Sigma^{-1}(F_{\hat{\theta}_{MLE}}) Y,$$

which expresses the regression parameter in terms of $\hat{\theta}_{MLE}$. For the computation of (3.3) we must calculate the acf corresponding to F_θ , which can be done using the algorithms of Section 2. Contrast (3.3) with (3.1); they are similar, the main difference being the replacement of the inverse of $\Sigma(F_\theta)$ by $\Sigma(F_\theta^{-1})$, which is equal to $\Sigma(F_{-\theta})$ for the cepstral model.

Most prior literature on random fields seems to utilize approximate Whittle estimation, or QMLE, since the objective function is quite simple to write down. The parameter MLEs do not have the bias problem of QMLEs, discussed above, but require more effort to compute due to matrix inversion. We can use the approximate algorithm given by equation (2.5), together with (3.3), to compute the MLEs. The QMLEs, based on unbiased $\hat{\gamma}_{h_1, h_2}(I_\beta)$ acf estimates, are faster to compute than MLEs and enjoy the same asymptotic normality and efficiency.

However, if one prefers a Bayesian estimation of θ (and β), it is necessary to compute $\exp \mathcal{L}(\theta, \beta)$, which is proportional to the data likelihood $p(Y|\theta, \beta)$. The posterior for θ is proportional to the likelihood times the prior, and one can use Markov chain Monte Carlo (MCMC) methods to approximate $p(\theta|Y)$ [17]. Recall that the mean of this distribution, which is the conditional expectation of θ given Y , is called the posterior mean, and will be denoted $\hat{\theta}_B$.

Note that equations (3.3) and (3.4) can each be used in an iterative estimation scheme. To determine the MLE, minimize (3.3) to obtain an estimate of θ for a given β computed via (3.4); then update $\hat{\beta}$ by plugging into (3.4), and iterate. For the QMLE, de-mean the data by computing $\widehat{Y} - \widehat{X}\beta$ and determining \widehat{I}_β , for a given β , and then minimize either \widehat{KL} or \widehat{KL}_N to obtain θ estimates (either exact or approximate); then update β by plugging into (3.2) and iterate. From now on, we refer to these estimates as the exact/approximate QMLEs (if using biased acf estimates $\gamma_{h_1, h_2}(I_\beta)$, only consistency holds, and not asymptotic normality).

3.3. *Distributional Properties of Parameter Estimates.* We now provide a description of the asymptotics for the various estimates; a rigorous treatment is given in Section 4, with formal statements of sufficient conditions and auxiliary results. Firstly, the Bayesian estimates $\hat{\theta}_B$ and $\hat{\beta}_B$ are consistent when the data is a Gaussian random field that satisfies suitable regularity conditions (Theorem 4.2). For the Frequentist case, recall that the number of observations equals $N_1 \cdot N_2$, so that a Central Limit Theorem result requires scaling by $\sqrt{N_1 \cdot N_2}$; we require that both dimensions expand, i.e., $\min\{N_1, N_2\} \rightarrow \infty$. Let the Hessian of the KL be denoted $H(\theta) = \nabla \nabla' KL(F_\theta, \tilde{F})$, which will be invertible at the unique pseudo-true value $\tilde{\theta}$ by assumption. Then the exact QMLE, approximate QMLE, and MLE for θ are all consistent, and are also asymptotically normal at rate N with mean $\tilde{\theta}$ and variance $H^{-1}(\tilde{\theta})V(\tilde{\theta})H^\dagger(\tilde{\theta})$, where \dagger denotes inverse transpose and $V(\theta) = 2 \langle \tilde{F}^2 \nabla F_\theta^{-1} \nabla' F_\theta^{-1} \rangle$. (This assumes that the fourth cumulants are zero; otherwise a more complicated expression for V results, involving the fourth-order spectral density.) The estimates of the regression parameters are asymptotically normal and independent of the θ estimates (when the third cumulants are zero), for all three types of estimates.

These theoretical results can be used to refine models. Typically, one uses these types of asymptotic results under the null hypothesis that the model is correctly specified, so that $\tilde{\theta}$ is the true parameter and $V = 2 \langle \nabla \log F_\theta \nabla' \log F_\theta \rangle$, which equals twice H . See McElroy and Holan [26] and McElroy and Findley [25] for more exposition on model misspecification in the frequency domain. Thus, the asymptotic variance is twice the inverse Hessian, or the inverse of $H/2$. Note that the Fisher information matrix is the Hessian of the asymptotic form of the Whittle likelihood, and hence is equal to one half of the Hessian of KL, i.e., $H/2$. Therefore when the model is correctly specified, parameter estimation is efficient.

Furthermore, the Fisher information matrix has a particularly elegant form in the case of a cepstral model. The gradient of the log spectrum is in this case just the various $Z_1^{j_1}$ or $Z_2^{j_2}$, so that as in the time series case the Hessian equals twice the identity matrix (because of mirror reflectional symmetry in Θ , there is a doubling that occurs), except for the case of the entry corresponding to $\Theta_{0,0}$ – in this case the derivative of the log spectrum with respect to $\Theta_{0,0}$ equals one. Thus the Fisher information matrix for all the parameters except $\Theta_{0,0}$ is equal to the identity matrix, and hence the asymptotic variance of any cepstral coefficient estimate is N^{-2} (or $2N^{-2}$ in the case of $\Theta_{0,0}$). The lack of cross-correlation in the parameter estimates asymptotically indicates there is no redundancy in the information they convey, which is a type of “maximal efficiency” in the cepstral model.

In terms of model-building with cepstral random fields, one procedure is the following: postulate a low order cepstral field model (e.g., order $p_1 = 1$) and jointly test for whether any coefficients (estimated via MLE or QMLE) are equal to zero. We might consider expanding the model – in the direction of one spatial axis or another as appropriate – if coefficients are significantly different from zero. Although this is not an *optimal* method of model selection, this type of forward addition strategy would stop once all additional coefficients are negligible. Alternatively, one could start with a somewhat larger cepstral model, and iteratively delete insignificant coefficients.

Gaussian Likelihood Ratio test statistics can be utilized for nested cepstral models, along the lines given in Taniguchi and Kakizawa [39] – which ultimately just depend on the asymptotic normality of the parameter estimates – in order to handle batches of parameters concurrently. Model selection and assessment can also be assisted by examination of spatial residuals, which are defined by applying the inverse square root of the estimated data covariance matrix $\Sigma(\hat{F}_{\hat{\theta}})$ to the vectorized centered data W – the result is a vectorized residual sequence, which should behave like white noise if the model has extracted all correlation structure. Note that examining whiteness of the vectorized residuals is equivalent to looking at all spatial correlations of the spatial residuals defined by undoing the vec operation. In the context of lattice data, one popular method for testing the null hypothesis of the absence of spatial autocorrelation is through the use of Moran’s I statistic [27, 10]. For a comprehensive discussion regarding Moran’s I statistic and its limitations see Cressie [11], Li, Calder and Cressie [23], Cressie and Wikle [12], and the references therein. In our case (Appendix A), we will evaluate goodness-of-fit by applying Moran’s I statistic to the spatial residuals obtained from the estimated model.

4. Theory of Inference. This section provides rigorous mathematical results regarding the inference problems delineated in Section 3. We do not assume a cepstral random field process, retaining greater generality, but assume a fair amount of regularity on the higher moments of the field through the Brillinger-type cumulant conditions [8]. We need not assume the field is Gaussian for Theorem 4.1, but we require a Gaussian assumption for Theorem 4.2. We first list technical assumptions, and then describe the mathematical results.

Previous rigorous work on asymptotics for parameter estimates of lattice random fields includes Guyon [18] and Mardia and Marshall [24]. Although Solo [37] advocates the approximate QMLE method in practice, asymptotic results are not proved in that paper. Our approach, like Mardia and Mar-

shall [24] and Pierce [29], handles regression effects together with parameter estimates, but we utilize broader data process assumptions formulated in terms of cumulants; Mardia and Marshall [24] assumes that the random field is Gaussian, whereas we do not. Pierce [29] treats the $d = 1$ time series case, and allows for non-Gaussian marginals, and shows that skewness can produce asymptotic correlation between regression and model parameter estimates; an analogous story for $d = 2$ is described in Theorem 4.1. Our contribution broadens the applicability of Mardia and Marshall [24] to non-Gaussian fields, and we moreover provide sufficient conditions under which our Lemma 4.1 yields the validity of condition (iii) of Theorem 2 of Mardia and Marshall [24]. This highlights our frequentist contribution; for Bayesian analysis, we are unaware of any published work on asymptotic concentration for random fields. Theorem 4.2 assumes a Gaussian field, which is natural given that the likelihood is Gaussian.

4.1. Regularity Assumptions. We first set out some notations and working assumptions: define a block-Toeplitz matrix $\Sigma(F)$ associated with spectral density F and an $N_1 \times N_2$ data matrix \mathbb{Y}^N to be $N^2 \times N^2$, with j_1, k_1 th block (for $1 \leq j_1, k_1 \leq N_1$) given by the $N_2 \times N_2$ -dimensional matrix $\Sigma(F_{j_1-k_1})$, which is defined as follows. If we integrate over the second variable of F we obtain a function of the first frequency:

$$F_{h_1}(\lambda_2) = \frac{1}{2\pi} \int_{-\pi}^{\pi} F(\lambda_1, \lambda_2) e^{ih_1\lambda_1} d\lambda_1, \quad \text{with } 0 \leq h_1 < N_1.$$

Then $\Sigma(F_{h_1})$ is the $N_2 \times N_2$ -dimensional matrix of inverse DFTs of F_{h_1} , with j_2, k_2 th entry given by

$$\gamma_{h_1, j_2-k_2}(F) = \frac{1}{4\pi^2} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} F(\lambda_1, \lambda_2) e^{ih_1\lambda_1} d\lambda_1 e^{i(j_2-k_2)\lambda_2} d\lambda_2 = \langle F Z_1^{-h_1} Z_2^{k_2-j_2} \rangle.$$

Based on how we have defined \mathbb{W}^N and $W = \text{vec}(\mathbb{W}^{N'})$, it follows that $\Sigma(\tilde{F}) = \mathbb{E}[WW']$, where \tilde{F} corresponds to the true data process. That is, lexicographical ordering of a stationary field produces this structure in the covariance matrix; there are N_1^2 blocks, each of which are $N_2 \times N_2$ -dimensional.

Also let \mathcal{F} denote the set of admissible spectra for two-dimensional random fields, defined as follows. For any spatial autocovariance function $\{\gamma_{h_1, h_2}\}$, consider the sums $S_{h_1, \cdot} = \sum_{h_2} |h_2| |\gamma_{h_1, h_2}|$, $S_{\cdot, h_2} = \sum_{h_1} |h_1| |\gamma_{h_1, h_2}|$, and $S_{\cdot, \cdot} = \sum_{h_1, h_2} |h_1| |h_2| |\gamma_{h_1, h_2}|$, and define the set

$$\mathcal{F} = \left\{ F : [-\pi, \pi]^2 \rightarrow \mathbb{R}^+, F(\lambda_1, \lambda_2) = \sum_{h_1, h_2} \gamma_{h_1, h_2}(F) Z_1^{h_1} Z_2^{h_2}, \right. \\ \left. S_{h_1, \cdot} < \infty \forall h_1, S_{\cdot, h_2} < \infty \forall h_2, S_{\cdot, \cdot} < \infty \right\}.$$

Note that this class excludes spectra with zeroes, which is a minor imposition in practice.

In this paper we take Brillinger's approach to asymptotic derivations, stipulating summability conditions on higher cumulants of the spatial field. Let us denote an integer-valued bivariate index by $t \in \mathbb{Z}^2$, which has integer coordinates (t_1, t_2) . Then a collection of spatial variables can be written $\{\mathbb{W}_{t^{(1)}}, \mathbb{W}_{t^{(2)}}, \dots\}$. The weak stationarity condition stipulates that joint moments of such variables only depend upon differences between indices: $t^{(1)} - t^{(2)} = (t_1^{(1)} - t_1^{(2)}, t_2^{(1)} - t_2^{(2)})$, etc. If we sum a function with respect to $t \in \mathbb{Z}^2$, the notation refers to a double sum over t_1 and t_2 . A similar notation is used for frequencies $\lambda \in [-\pi, \pi]^2$, in that $\lambda = (\lambda_1, \lambda_2)$.

Suppose that spatial data is sampled from a true spatial field with spectrum F , and that we have a collection of continuous weighting functions $G_j : [-\pi, \pi]^2 \mapsto \mathbb{R}^+$. The second cumulant function of the spatial field is the autocovariance function γ_h with $h \in \mathbb{Z}^2$, whereas the $(k + 1)$ th cumulant function is denoted

$$\gamma_{h^{(1)}, h^{(2)}, \dots, h^{(k)}} = \text{cum} [\mathbb{W}_t, \mathbb{W}_{t+h^{(1)}}, \mathbb{W}_{t+h^{(2)}}, \dots, \mathbb{W}_{t+h^{(k)}}].$$

We require absolute summability of these second and fourth cumulant functions. Then the fourth-order spectrum is well-defined via

$$\widetilde{FF}(\lambda^{(1)}, \lambda^{(2)}, \lambda^{(3)}) = \sum_{h^{(1)}, h^{(2)}, h^{(3)}} \gamma_{h^{(1)}, h^{(2)}, h^{(3)}} e^{\{-i\lambda^{(1)} \cdot h^{(1)} - i\lambda^{(2)} \cdot h^{(2)} - i\lambda^{(3)} \cdot h^{(3)}\}},$$

with \cdot denoting the dot product of bivariate vectors. More regularity can be imposed via the condition

$$(4.1) \quad \sum_{h^{(1)}, h^{(2)}, \dots, h^{(k)}} \left(1 + \overline{|h^{(1)}|} \overline{|h^{(2)}|} \dots \overline{|h^{(k)}|}\right) |\gamma_{h^{(1)}, h^{(2)}, \dots, h^{(k)}}| < \infty,$$

where \bar{t} denotes the product of the components of t . This will be referred to as Condition B_k , for any $k \geq 1$; note that B_2 implies the summability conditions of the set \mathcal{F} . Finally, recall that the periodogram is computed from a sample of size $N^2 = N_1 \cdot N_2$. When the regressors are correctly specified, we will write $\tilde{\beta}$ for the true parameter. Then $I_{\tilde{\beta}}$ denotes the periodogram of the data Y correctly adjusted for mean effects; equivalently, it is the periodogram of \mathbb{W}^N .

In addition to assuming that the regressors are correctly specified, with $\tilde{\beta}$ the true regression parameter and \tilde{X} the regression matrix, we require the following key assumptions.

Assumptions:**A1:** $\tilde{F} \in \mathcal{F}$.**A2:** The spectral density F_θ is twice continuously differentiable and uniformly bounded above and away from zero, and moreover all components of F_θ , ∇F_θ , $\nabla\nabla'F_\theta$ are in \mathcal{F} .**A3:** The process is weakly stationary of order k , and the Brillinger conditions B_k (4.1) hold for all $k \geq 1$.**A4:** The pseudo-true value $\tilde{\theta}$ exists uniquely in the interior of the parameter space.**A5:** $H(\theta) = \nabla\nabla'\text{KL}(F_\theta, \tilde{F})$ is invertible at $\tilde{\theta}$.

Conditions **A1**, **A3**, and **A5** cannot be verified from data, but some assumptions of this nature must be made to obtain asymptotic formulas. Condition **A2** will hold for cepstral models (and other random field models as well) by the following argument. The coefficients of the causal and skew fields will have exponential decay in either index argument, by extensions of the classical time series argument (see for example Hurvich [19]) applied to (2.9) and (2.10). (The time series argument can be directly applied to (2.11) and (2.12) as well.) Combining these results using (2.8), the acf of the cepstral field will also have exponential decay so that $F_\theta \in \mathcal{F}$. Of course, another way to verify this condition is to examine the boundedness of partial derivatives of the spectrum; at once we see that **A2** holds for the cepstral model, as it does for moving average random fields.

Although condition **A4** may be problematic for certain moving average models (which may have complicated constraints on coefficients), the cepstral model uses no constraints on θ , because the distinct entries of Θ can be any real number, independently of all other distinct entries. Euclidean space is open, so any pseudo-true value is necessarily contained in the interior. Also, existence of a pseudo-true value is guaranteed by convexity of the KL discrepancy.

For the result on Bayesian estimation, we will assume that the model is correctly specified; the model must also be identifiable, i.e., $F_{\theta_1} = F_{\theta_2}$ implies $\theta_1 = \theta_2$, which helps ensure asymptotic concentration of the likelihood. We assume the parameters belong to some compact subset of Euclidean space, and the true parameter vector lies in the interior. This assumption can often be accomplished by prior transformation (and is easily accomplished for the cepstral coefficients in the cepstral model). Also define the matrix 2-norm of a matrix A via the notation $\|A\|_2$.

4.2. *Technical Results.* We begin with an important Lemma that extends Lemma 4.1.2 of Taniguchi and Kakizawa [39] to the spatial context.

LEMMA 4.1. *Let $\Sigma(F_j)$ and $\Sigma(G_j)$ be block-Toeplitz matrices with $F_j, G_j^{-1} \in \mathcal{F}$ for $1 \leq j \leq m$. Assuming that $N_* = \min\{N_1, N_2\} \rightarrow \infty$, and $N = \sqrt{N_1 \cdot N_2}$,*

$$N^{-2} \text{tr}\{\Pi_{j=1}^m \Sigma(F_j) \Sigma^{-1}(G_j)\} = \langle \Pi_{j=1}^m F_j G_j^{-1} \rangle + O(N^{-2}).$$

Next, we discuss a Lemma that provides a Central Limit Theorem for weighted averages of the spatial periodogram, which is a natural extension of Lemma 3.1.1 of Taniguchi and Kakizawa [39]. Define the bias-correction quantities

$$(4.2) \quad B_1(\lambda) = \sum_{h_1, h_2} |h_1| \gamma_{h_1, h_2} e^{-i\lambda \cdot h} \quad B_2(\lambda) = \sum_{h_1, h_2} |h_2| \gamma_{h_1, h_2} e^{-i\lambda \cdot h},$$

and use $\langle\langle g(\lambda^{(1)}, \lambda^{(2)}) \rangle\rangle$ as a short hand for $(2\pi)^{-4} \int_{[-\pi, \pi]^4} g(\lambda^{(1)}, \lambda^{(2)}) d\lambda^{(1)} d\lambda^{(2)}$.

LEMMA 4.2. *Assume that $N_* = \min\{N_1, N_2\} \rightarrow \infty$ and let $N = \sqrt{N_1 \cdot N_2}$. Suppose Assumption **A3** holds, and that G_j for $1 \leq j \leq J$ are continuous functions. Let $G_j^*(\lambda) = G_j(-\lambda)$. Then:*

(i) *For the unbiased acf estimators, as $N_* \rightarrow \infty$, $\langle G_j \widehat{I}_{\tilde{\beta}} \rangle - \langle G_j \widehat{I}_{\tilde{\beta}} \rangle_N \xrightarrow{P} 0$ and $\langle G_j \widehat{I}_{\tilde{\beta}} \rangle \xrightarrow{P} \langle G_j \widetilde{F} \rangle$ for any $1 \leq j \leq J$. Also*

$$N \{ \langle G_j (\widehat{I}_{\tilde{\beta}} - \widetilde{F}) \rangle \}_{j=1}^J \xrightarrow{\mathcal{L}} \mathcal{N}(0, V),$$

where the covariance matrix V has jk th entry

$$\langle\langle G_j G_k \widetilde{F} \widetilde{F}(\lambda^{(1)}, -\lambda^{(2)}, \lambda^{(2)}) \rangle\rangle + \langle\langle (G_j G_k^* + G_j G_k) \widetilde{F}^2 \rangle\rangle.$$

(ii) *For the biased acf estimators, $\langle G_j I_{\tilde{\beta}} \rangle - \langle G_j I_{\tilde{\beta}} \rangle_N \xrightarrow{P} 0$ and $\langle G_j I_{\tilde{\beta}} \rangle \xrightarrow{P} \langle G_j \widetilde{F} \rangle$ for any $1 \leq j \leq J$. Also, for the same V given in case (i),*

$$N \{ \langle G_j (I_{\tilde{\beta}} - \widetilde{F} + N^{-1} B_1 + N^{-1} B_2) \rangle \}_{j=1}^J \xrightarrow{\mathcal{L}} \mathcal{N}(0, V),$$

where the bias correction terms are defined in (4.2).

The last assertion of Lemma 4.2 means that utilizing $I_{\tilde{\beta}}$ instead of $\widehat{I}_{\tilde{\beta}}$ will require a bias correction; cf. Guyon [18]. Both lemmas are important preliminary results for our main theorems, but also are of interest in their own right, extending known time series results to the spatial context. Although generalizations to dimensions higher than two seem feasible, the actual mechanics

become considerably more technical. We now state the limit theorems for our parameter estimates. For the QMLE estimates, we suppose that they are either exact or approximate Whittle estimates defined using the unbiased acf estimates.

THEOREM 4.1. *Assume that conditions **A1-A5** hold, and that the regressors are correctly specified with $(\tilde{X}'\tilde{X})^{-1} \rightarrow 0$ as $N_* = \min\{N_1, N_2\} \rightarrow \infty$. Then in the case of MLE or the QMLE, both $\hat{\beta}$ and $\hat{\theta}$ are jointly asymptotically normal with distributions given by*

$$N \left(\hat{\theta} - \tilde{\theta} \right) \xrightarrow{\mathcal{L}} \mathcal{N} \left(0, H^{-1}(\tilde{\theta})V(\tilde{\theta})H^\dagger(\tilde{\theta}) \right)$$

$$H(\theta) = \nabla \nabla' KL(F_\theta, \tilde{F})$$

$$V(\theta) = 2 \langle \tilde{F}^2 \nabla F_\theta^{-1} \nabla' F_\theta^{-1} \rangle + \langle \langle \nabla F_\theta^{-1} \nabla' F_\theta^{-1} \tilde{F} \tilde{F}' (\lambda^{(1)}, -\lambda^{(2)}, \lambda^{(2)}) \rangle \rangle,$$

where \dagger denotes an inverse transpose and $N = \sqrt{N_1 \cdot N_2}$. Also $N(\hat{\beta} - \tilde{\beta})$ is asymptotically normal with mean zero and covariance matrix

$$M_X^{-1}(\tilde{\theta}) \left[\tilde{X}' \Sigma(F_\theta^{-1}) \Sigma(\tilde{F}) \Sigma(F_\theta^{-1}) \tilde{X} \right] M_X^\dagger(\tilde{\theta}),$$

where $M_X(\theta) = \tilde{X}' \Sigma(F_\theta^{-1}) \tilde{X}$. Finally, $\hat{\beta}$ and $\hat{\theta}$ are asymptotically independent if the third cumulants of the process are zero.

REMARK 4.1. For a Gaussian process, third and fourth cumulants are zero, which implies that regression and model parameter estimates are asymptotically independent, and that V has a simpler form, being given just by $2 \langle \tilde{F}^2 \nabla F_\theta^{-1} \nabla' F_\theta^{-1} \rangle$.

REMARK 4.2. Application of the same techniques in the case of a one-dimensional random field, or time series, yields asymptotic normality of regression and time series parameters under Brillinger's conditions. To our knowledge, the only other results of this flavor for time series with regression effects is the work of Pierce [29], which focuses on ARIMA models but allows for skewed non-Gaussian distributions.

THEOREM 4.2. *Assume that the data process is Gaussian and **A2** and **A4** hold, and that the model is correctly specified and is identifiable. Also suppose that the regressors are correctly specified, with $N^{-2} \tilde{X}' \Sigma^{-1}(F_\theta) \tilde{X} \rightarrow M(\theta)$ for some $M(\theta)$ satisfying $0 < \sup_\theta \|M(\theta)\|_2 < \infty$. Then $\hat{\beta}_B \xrightarrow{P} \tilde{\beta}$ and $\hat{\theta}_B \xrightarrow{P} \tilde{\theta}$ as $N_* = \min\{N_1, N_2\} \rightarrow \infty$.*

It is worth comparing the conditions of the two theorems. In Theorem 4.2 the assumption of a correct model makes **A1** automatic, and the Gaussian assumption makes **A3** automatic. Furthermore, the assumption in Theorem 4.2 on the parameters – together with the assumption of a correct model – automatically entails **A4** as well. Theorem 4.1 also assumes **A5**, which is chiefly needed to establish asymptotic normality of the Frequentist estimates. The Bayesian result requires a slightly stronger assumption on the regression matrix in order to get asymptotic concentration of the likelihood. For example, if we seek to estimate a constant mean by taking \tilde{X} to be a column vector of all ones, then $M(\theta)$ exists and is just the scalar $F_{-\theta}(0, 0)$; this will be bounded away from zero and infinity in the cepstral model if all the cepstral coefficients are restricted to a range of values.

5. Simulation Study. To demonstrate the effectiveness of our approach, we conducted a small simulation study using maximum likelihood estimation as outlined in Sections 2 and 3. The model autocovariances were calculated according (2.5), with $M = 1000$ and $p_1 = p_2 = 2$. The exact parameter values for the simulation were calibrated to the straw yield data analysis presented in the Supplement’s Appendix A. Grid sizes of (15×15) , (20×20) , (20×25) , and (25×25) were considered, where (20×25) constitutes the size grid in our real-data example.

For this simulation, we generated 200 Gaussian datasets with parameters $\theta = J\Theta$ corresponding to quadrants I and II of the grid Θ^1 , and $\beta = (\beta_0, \beta_1, \beta_2)'$; see Table 1. In this case, β_1 and β_2 correspond to “row” and “column” effects, respectively, in the agricultural experiment considered. Here, the row and column effects are obtained by regressing the vectorized response on the corresponding row and column indices (since rows and columns are equally spaced). The \tilde{X} matrix used in this simulation consisted of a column of ones followed by columns associated with the row and column effects and was taken from the analysis presented in the Supplement’s Appendix A. Given θ and β , we simulate directly from the corresponding multivariate Gaussian distribution. However, in cases where the grid size is extremely large, another potential approach to simulation would be circular embedding [9, 43], though it would be necessary to properly account for any regression effects. The log Gaussian likelihood (up to constants) given by (3.3) was numerically maximized for each simulated dataset using the *optim* function in R [40].

As demonstrated in Table 1, through an assessment of mean square error

¹i.e., $\theta = [\Theta_{-2,2}, \Theta_{-2,1}, \Theta_{-1,2}, \Theta_{-1,1}, \Theta_{0,2}, \Theta_{0,1}, \Theta_{1,2}, \Theta_{1,1}, \Theta_{2,2}, \Theta_{2,1}, \Theta_{-2,0}, \Theta_{-1,0}, \Theta_{0,0}]'$
 $= (\theta_1, \theta_2, \dots, \theta_{13})'$.

(mse), the model parameters can be estimated with a high degree of precision. Additionally, Table 1 illustrates that our asymptotic theory agrees with the finite sample estimates for different grid sizes. Specifically, we provide the difference between the mean standard deviation (over all of the cepstral parameters – except $\Theta_{0,0}$) and the asymptotic standard deviation. This simulation shows that, as the grid size increases, the difference between the estimated standard error and the asymptotic standard error goes to zero on average. We also provide the mean, standard deviation, and mse for the individual parameters, including the mean parameters β ; this demonstrates the bias properties, as well as the fact that the mse goes to zero as the grid size increases. Lastly, the average p -value for the Shapiro-Wilks test of normality for each simulation grid size (over all of the cepstral parameters) was greater than 0.4, with only one parameter out of the thirteen cepstral parameters from each simulation not exhibiting normality. Hence, the estimated parameters converge to their asymptotic distribution and, as expected, their precision increases with sample size.

6. Conclusion. The general modeling approach and asymptotic theory we propose extends the spatial random field literature in several directions. By providing recursive formulas for calculating autocovariances, from a given cepstral random field model, we have facilitated usage of these models in both Bayesian and likelihood settings. This is extremely notable as many models suffer from a constrained parameter space, whereas the cepstral random field model imposes no constraints on the parameter values. More specifically, the autocovariance matrix obtained from our approach is guaranteed to be positive definite.

In addition, we establish results on consistency and asymptotic normality for an expanding domain. This provides a rigorous platform for conducting model selection and statistical inference. The asymptotic results are proven generally and can be viewed as an independent contribution to the random field literature, expanding on the results of Mardia and Marshall [24] and others, such as Guyon [18]. The simulation results support the theory, and the methods are illustrated through an application to straw yield data from an agricultural field experiment (Supplement’s Appendix A). In this setting, it is readily seen that our model is easily able to characterize the underlying spatial dependence structure.

Acknowledgements. We thank the A.E. and two anonymous referees for providing detailed comments that helped substantially improve this article. The authors would also like to thank Aaron Porter for his assistance with implementation of the Moran’s I statistic. This article is released to in-

form interested parties of research and to encourage discussion. The views expressed on statistical issues are those of the authors and not necessarily those of the U.S. Census Bureau. See [Supplement](#) for the supplementary material.

SUPPLEMENTARY MATERIAL

Supplement: Supplement to Asymptotic Theory of Cepstral Random Fields

(<http://www.e-publications.org/ims/support/download/imsart-ims.zip>). The supplement contains a description of further applications of the cepstral model, analysis of straw yield data, as well as all proofs.

References.

- [1] BANDYOPADHYAY, S. and LAHIRI, S. (2010). Asymptotic Properties of Discrete Fourier Transforms for Spatial Data. *Sankhya, Series A* **71-A** 221–259.
- [2] BANERJEE, S., CARLIN, B. P. and GELFAND, A. E. (2004). *Hierarchical Modeling and Analysis for Spatial Data* **101**. Chapman & Hall.
- [3] BESAG, J. (1972). On the Correlation Structure of Some Two-Dimensional Stationary Processes. *Biometrika* **59** 43–48.
- [4] BESAG, J. (1974). Spatial Interaction and the Statistical Analysis of Lattice Systems. *Journal of the Royal Statistical Society – Series B (Methodological)* 192–236.
- [5] BESAG, J. and GREEN, P. J. (1993). Spatial Statistics and Bayesian Computation. *Journal of the Royal Statistical Society – Series B (Methodological)* 25–37.
- [6] BLOOMFIELD, P. (1973). An Exponential Model for the Spectrum of a Scalar Time Series. *Biometrika* **60** 217–226.
- [7] BOCHNER, S. (1955). *Harmonic Analysis and the Theory of Probability*. Berkeley: University of California Press.
- [8] BRILLINGER, D. R. (2001). *Time Series: Data Analysis and Theory* **36**. Philadelphia: SIAM.
- [9] CHAN, G. and WOOD, A. T. (1999). Simulation of Stationary Gaussian Vector Fields. *Statistics and Computing* **9** 265–268.
- [10] CLIFF, A. D. and ORD, J. K. (1981). *Spatial Processes: Models & Applications* **44**. Pion London.
- [11] CRESSIE, N. (1993). *Statistics for Spatial Data*. New York: John Wiley and Sons.
- [12] CRESSIE, N. and WIKLE, C. K. (2011). *Statistics for Spatio-Temporal Data*. John Wiley and Sons.
- [13] DAHLHAUS, R. and KÜNSCH, H. (1987). Edge Effects and Efficient Parameter Estimation for Stationary Random Fields. *Biometrika* **74** 877–882.
- [14] FUENTES, M. (2002). Spectral Methods for Nonstationary Spatial Processes. *Biometrika* **89** 197–210.
- [15] FUENTES, M., GUTTORP, P. and SAMPSON, P. D. (2007). Using Transforms to Analyze Space-time Processes. *Statistical Methods for Spatio-Temporal Systems* 77–150.
- [16] FUENTES, M. and REICH, B. (2010). Spectral Domain. *Handbook of Spatial Statistics* 57–77.
- [17] GEWEKE, J. (2005). *Contemporary Bayesian Econometrics and Statistics* **537**. Wiley-Interscience.

- [18] GUYON, X. (1982). Parameter Estimation for a Stationary Process on a d -Dimensional Lattice. *Biometrika* **69** 95–105.
- [19] HURVICH, C. M. (2002). Multistep Forecasting of Long Memory Series Using Fractional Exponential Models. *International Journal of Forecasting* **18** 167–179.
- [20] KEDEM, B. and FOKIANOS, K. (2002). *Regression Models for Time Series Analysis* **323**. John Wiley and Sons.
- [21] KIZILKAYA, A. (2007). On the Parameter Estimation of 2-D Moving Average Random Fields. *Circuits and Systems II: Express Briefs, IEEE Transactions on* **54** 989–993.
- [22] KIZILKAYA, A. and KAYRAN, A. H. (2005). ARMA-Cepstrum Recursion Algorithm for the Estimation of the MA Parameters of 2-D ARMA Models. *Multidimensional Systems and Signal Processing* **16** 397–415.
- [23] LI, H., CALDER, C. A. and CRESSIE, N. (2007). Beyond Moran's I: testing for spatial dependence based on the spatial autoregressive model. *Geographical Analysis* **39** 357–375.
- [24] MARDIA, K. and MARSHALL, R. (1984). Maximum Likelihood Estimation of Models for Residual Covariance in Spatial Regression. *Biometrika* **71** 135–146.
- [25] MCELROY, T. S. and FINDLEY, D. F. (2010). Selection Between Models Through Multi-Step-Ahead Forecasting. *Journal of Statistical Planning and Inference* **140** 3655–3675.
- [26] MCELROY, T. S. and HOLAN, S. H. (2009). A Local Spectral Approach for Assessing Time Series Model Misspecification. *Journal of Multivariate Analysis* **100** 604–621.
- [27] MORAN, P. A. (1950). Notes on Continuous Stochastic Phenomena. *Biometrika* 17–23.
- [28] NOH, J. and SOLO, V. (2007). A True Spatio-Temporal Test Statistic for Activation Detection in fMRI by Parametric Cepstrum. In *Acoustics, Speech and Signal Processing, 2007. ICASSP 2007. IEEE International Conference on* **1** I–321. IEEE.
- [29] PIERCE, D. (1971). Least Squares Estimation in the Regression Model with Autoregressive-Moving Average Errors. *Biometrika* **58** 299–312.
- [30] POLITIS, D. N. and ROMANO, J. P. (1995). Bias-Corrected Nonparametric Spectral Estimation. *Journal of time series analysis* **16** 67–103.
- [31] POLITIS, D. N. and ROMANO, J. P. (1996). On Flat-top Kernel Spectral Density Estimators for Homogeneous Random Fields. *Journal of statistical planning and inference* **51** 41–53.
- [32] POURAHMADI, M. (1984). Taylor Expansion of $\exp\{\sum_{k=0}^{\infty} a_k z^k\}$ and Some Applications. *American Mathematical Monthly* **91** 303–307.
- [33] ROSENBLATT, M. (1985). *Stationary Sequences and Random Fields*. Birkhauser.
- [34] ROSENBLATT, M. (2000). *Gaussian and Non-Gaussian Linear Time Series and Random Fields*. Springer Verlag.
- [35] RUE, H. and HELD, L. (2010). Discrete Spatial Variation. In *Handbook of Spatial Statistics* (A. Gelfand, P. Diggle, M. Fuentes and P. Guttorp, eds.) Chapman and Hall, London.
- [36] SANDGREN, N. and STOICA, P. (2006). On Nonparametric Estimation of 2-D Smooth Spectra. *Signal Processing Letters, IEEE* **13** 632–635.
- [37] SOLO, V. (1986). Modeling of Two-Dimensional Random Fields by Parametric Cepstrum. *IEEE Transactions on Information Theory* **32** 743–750.
- [38] STEIN, M. L. (1999). *Interpolation of Spatial Data: Some Theory for Kriging*. Springer Verlag.
- [39] TANIGUCHI, M. and KAKIZAWA, Y. (2000). *Asymptotic Theory of Statistical Inference for Time Series*. New York: Springer.
- [40] R DEVELOPMENT CORE TEAM (2012). R: A Language and Environment for Sta-

tistical Computing R Foundation for Statistical Computing, Vienna, Austria ISBN 3-900051-07-0.

[41] TONELLATO, S. F. (2007). Random Field Priors for Spectral Density Functions. *Journal of Statistical Planning and Inference* **137** 3164–3176.
 [42] WHITTLE, P. (1954). On Stationary Processes in the Plane. *Biometrika* 434–449.
 [43] WOOD, A. T. and CHAN, G. (1994). Simulation of Stationary Gaussian Processes in $[0, 1]^d$. *Journal of Computational and Graphical Statistics* **3** 409–432.

TABLE 1

Simulation results for the simulation presented in Section 5 ($p_1 = p_2 = 2$). Note, there were 200 simulated datasets and $\bar{\sigma}_\theta$ denotes the mean standard deviation for parameters $\theta_1, \dots, \theta_{12}$ for a given simulation (over the 200 datasets). Recall that, for $j = 1, \dots, 12$, the asymptotic standard error for θ_j equals $1/N$. Therefore, $\bar{\sigma}_\theta - 1/N$ represents the average difference between the estimated and asymptotic standard error for $\theta_1, \dots, \theta_{12}$. The values in the table below are only reported to three decimal places and the elements of θ are described in Section 5. Note that $N = \sqrt{N_1 \cdot N_2}$ and that $\theta_{13} = \Theta_{0,0}$.

		$N_1 \times N_2$											
		15×15			20×20			20×25			25×25		
	True	Mean	SD	MSE	Mean	SD	MSE	Mean	SD	MSE	Mean	SD	MSE
θ_1	0.009	-0.019	0.084	0.008	0.000	0.056	0.003	-0.001	0.049	0.003	-0.002	0.041	0.002
θ_2	-0.028	-0.037	0.079	0.006	-0.029	0.057	0.003	-0.033	0.048	0.002	-0.039	0.044	0.002
θ_3	0.132	0.123	0.081	0.007	0.127	0.059	0.003	0.133	0.044	0.002	0.129	0.045	0.002
θ_4	0.067	0.054	0.080	0.007	0.057	0.059	0.004	0.058	0.047	0.002	0.063	0.049	0.002
θ_5	0.271	0.266	0.078	0.006	0.265	0.056	0.003	0.269	0.052	0.003	0.259	0.043	0.002
θ_6	0.383	0.367	0.074	0.006	0.370	0.057	0.003	0.377	0.047	0.002	0.379	0.038	0.001
θ_7	0.001	-0.006	0.079	0.006	-0.001	0.060	0.004	-0.005	0.050	0.003	-0.001	0.044	0.002
θ_8	-0.017	-0.028	0.077	0.006	-0.023	0.059	0.004	-0.022	0.049	0.002	-0.020	0.045	0.002
θ_9	-0.003	-0.023	0.082	0.007	-0.012	0.052	0.003	-0.009	0.047	0.002	-0.009	0.046	0.002
θ_{10}	-0.055	-0.090	0.079	0.008	-0.064	0.049	0.002	-0.053	0.053	0.003	-0.061	0.040	0.002
θ_{11}	-0.015	-0.035	0.085	0.008	-0.022	0.054	0.003	-0.021	0.048	0.002	-0.021	0.044	0.002
θ_{12}	0.144	0.129	0.072	0.005	0.134	0.053	0.003	0.138	0.047	0.002	0.137	0.042	0.002
θ_{13}	-0.871	-0.968	0.096	0.019	-0.922	0.080	0.009	-0.902	0.064	0.005	-0.902	0.060	0.004
β_0	7.646	7.632	0.275	0.076	7.651	0.200	0.040	7.627	0.167	0.028	7.645	0.173	0.030
β_1	-0.035	-0.035	0.020	0.000	-0.034	0.011	0.000	-0.034	0.011	0.000	-0.034	0.008	0.000
β_2	-0.059	-0.056	0.024	0.001	-0.060	0.013	0.000	-0.059	0.009	0.000	-0.060	0.009	0.000
$\bar{\sigma}_\theta - (1/N)$		0.0124			0.0059			0.0037			0.0033		

CENTER FOR STATISTICAL RESEARCH AND METHODOLOGY, U.S. CENSUS BUREAU, 4600 SILVER HILL ROAD, WASHINGTON, D.C. 20233-9100
 E-MAIL: tucker.s.mcelroy@census.gov

DEPARTMENT OF STATISTICS, UNIVERSITY OF MISSOURI, 146 MIDDLEBUSH HALL, COLUMBIA, MO, 65211-6100
 E-MAIL: holans@missouri.edu