

# Ising models on locally tree-like graphs

Amir Dembo\* and Andrea Montanari†

April 2008, Revised: June 26, 2009

## Abstract

We consider ferromagnetic Ising models on graphs that converge locally to trees. Examples include random regular graphs with bounded degree and uniformly random graphs with bounded average degree. We prove that the ‘cavity’ prediction for the limiting free energy per spin is correct for *any positive temperature and external field*. Further, local marginals can be approximated by iterating a set of mean field (cavity) equations. Both results are achieved by proving the local convergence of the Boltzmann distribution on the original graph to the Boltzmann distribution on the appropriate infinite random tree.

## 1 Introduction

A ferromagnetic *Ising model on the finite graph*  $G$  (with vertex set  $V$ , and edge set  $E$ ) is defined by the following Boltzmann distributions over  $\underline{x} = \{x_i : i \in V\}$ , with  $x_i \in \{+1, -1\}$

$$\mu(\underline{x}) = \frac{1}{Z(\beta, B)} \exp \left\{ \beta \sum_{(i,j) \in E} x_i x_j + B \sum_{i \in V} x_i \right\}. \quad (1.1)$$

These distributions are parametrized by the ‘magnetic field’  $B$  and ‘inverse temperature’  $\beta \geq 0$ , where the partition function  $Z(\beta, B)$  is fixed by the normalization condition  $\sum_{\underline{x}} \mu(\underline{x}) = 1$ . Throughout the paper, we will be interested in sequences of graphs<sup>1</sup>  $G_n = (V_n \equiv [n], E_n)$  of diverging size  $n$ .

Non-rigorous statistical mechanics techniques, such as the ‘replica’ and ‘cavity methods,’ allow to make a number of predictions on the model (1.1), when the graph  $G$  ‘lacks any finite-dimensional structure.’ The most basic quantity in this context is the asymptotic *free entropy density*

$$\phi(\beta, B) \equiv \lim_{n \rightarrow \infty} \frac{1}{n} \log Z_n(\beta, B) \quad (1.2)$$

(this quantity is also sometimes called in the literature also free energy or pressure). The limit free entropy density and the large deviation properties of Boltzmann distribution were characterized in great detail [EN79] in the case of a complete graph  $G_n = K_n$  (the inverse temperature must then be scaled by  $1/n$  to get a non-trivial limit). Statistical physics predictions exist however for a much wider class of graphs, including most notably sparse random graphs with bounded average degree, see for instance [JP98, DGM04, LV+02]. This is a direction of interest for at least two reasons:

---

\*Departments of Statistics and Mathematics, Stanford University

†Departments of Electrical Engineering and Statistics, Stanford University

Research partially supported by NSF grant #DMS-0806211.

**AMS (2000) Subject Classification:** Primary: 82B44 Secondary: 82B23, 60F10, 60K35, 05C80, 05C05

**Keywords:** Ising model, random sparse graphs, cavity method, Bethe measures, belief propagation, local weak convergence.

<sup>1</sup>We adopt the notation  $[i] = \{1, 2, \dots, i\}$  for the set of first  $i$  integers.

(i) Sparse graphical structures arise in a number of problems from combinatorics and theoretical computer science. Examples include random satisfiability, coloring of random graphs, graph partitioning [MM09]. In all of these cases, the uniform measure over solutions can be regarded as the Boltzmann distribution for a modified spin glass with multi-spin interactions. Such problems have been successfully attacked using non rigorous statistical mechanics techniques.

A mathematical foundation of this approach is still lacking, and would be extremely useful.

(ii) Sparse graphs allow to introduce a non-trivial notion of distance between vertices, namely the length of the shortest path connecting them. This geometrical structure allows for new characterizations of the measure (1.1) in terms of correlation decay. This type of characterization is in turn related to the theory of Gibbs measures on infinite trees [KM+07].

The asymptotic free entropy density (1.2) was determined rigorously only in a few cases for sparse graphs. In [GM07], this task was accomplished for random regular graphs. De Santis and Guerra [DG08] developed interpolation techniques for random graphs with independent edges (Erdős-Renyi type) but only determined the free entropy density at high temperature and at zero temperature (in both cases with vanishing magnetic field). The latter is in fact equivalent to counting the number of connected components of a random graph. Interestingly, the partition function  $Z_n(\beta, B)$  can be approximated in polynomial time for  $\beta \geq 0$ , using an appropriate Monte Carlo Markov Chain algorithm [JS90]. It is intriguing that no general approximation algorithms exists in the case  $\beta < 0$  (the ‘anti-ferromagnetic’ Ising model). Correspondingly, the statistical physics conjecture for the free entropy density [MM09] becomes significantly more intricate (presenting the so called ‘replica symmetry breaking’ phenomenon).

In this paper we generalize the previous results by rigorously verifying the validity of the Bethe free entropy prediction for the value of the limit in (1.2) for generic graph sequences that converge locally to trees. Indeed, we control the free entropy density by proving that the Boltzmann measure (1.1) converges locally to the Boltzmann measure of a model on a tree. The philosophy is related to the local weak convergence method of [AS03].

Finally, several of the proofs have an algorithmic interpretation, providing an efficient procedure for approximating the local marginals of the Boltzmann measure. The essence of this procedure consists in solving by iteration certain mean field (cavity) equations. Such an algorithm is known in artificial intelligence and computer science under the name of *belief propagation*. Despite its success and wide applicability, only weak performance guarantees have been proved so far. Typically, it is possible to prove its correctness in the high temperature regime, as a consequence of a uniform decay of correlations holding there (spatial mixing) [TJ02, BG07, SS09]. The behavior of iterative inference algorithms on Ising models was recently considered in [MK05, SWW07].

The emphasis of the present paper is on the low-temperature regime in which uniform decorrelation does not hold. We are able to prove that belief propagation converges exponentially fast on any graph, and that the resulting estimates are asymptotically exact for large locally tree-like graphs. The main idea is to introduce a magnetic field to break explicitly the  $+/-$  symmetry, and to carefully exploit the monotonicity properties of the model.

A key step consists in estimating the correlation between the root spin of an Ising model on a tree, and positive boundary conditions. Ising models on trees are interesting per se, and have been the object of significant mathematical work, see for instance [Lyo00, JS99, EK+00]. The question considered here appears however to be novel.

The next section provides the basic technical definitions (in particular concerning graphs and local convergence to trees), and the formal statement of our main results. Notations and certain key tools are described in Section 3 with Section 4 devoted to proofs of the relevant properties of Ising models on trees (which are of independent interest). The latter are used in Sections 5 and 6 to derive our main results concerning models on tree-like graphs. A companion paper [DGM09] deals with the related challenging

problem of spin glass models on sparse graphs.

## 2 Definitions and main results

The next subsections contain some basic definitions on graph sequences and the notion of local convergence to random trees. Sections 2.2 and 2.3 present our results on the free entropy density and the algorithmic implications of our analysis.

### 2.1 Locally tree-like graphs

Let  $P = \{P_k : k \geq 0\}$  a probability distribution over the non-negative integers, with finite, positive first moment, and denote by

$$\rho_k = \frac{kP_k}{\sum_{l=1}^{\infty} lP_l}, \quad (2.1)$$

its size-biased version. For any  $t \geq 0$ , we let  $\mathsf{T}(P, \rho, t)$  denote the random rooted tree generated as follows. First draw an integer  $k$  with distribution  $P_k$ , and connect the root to  $k$  offspring. Then recursively, for each node in the last generation, generate an integer  $k$  independently with distribution  $\rho_k$ , and connect the node to  $k - 1$  new nodes. This is repeated until the tree has  $t$  generations.

Sometimes it will be useful to consider the ensemble  $\mathsf{T}(\rho, t)$  whereby the root node has degree  $k - 1$  with probability  $\rho_k$ . We will drop the degree distribution arguments from  $\mathsf{T}(P, \rho, t)$  or  $\mathsf{T}(\rho, t)$  and write  $\mathsf{T}(t)$  whenever clear from the context. Notice that the infinite trees  $\mathsf{T}(P, \rho, \infty)$  and  $\mathsf{T}(\rho, \infty)$  are well defined.

The average branching factor of trees will be denoted by  $\bar{\rho}$ , and the average root degree by  $\bar{P}$ . In formulae

$$\bar{P} \equiv \sum_{k=0}^{\infty} kP_k, \quad \bar{\rho} \equiv \sum_{k=1}^{\infty} (k-1)\rho_k. \quad (2.2)$$

We denote by  $G_n = (V_n, E_n)$  a graph with vertex set  $V_n \equiv [n] = \{1, \dots, n\}$ . The distance  $d(i, j)$  between  $i, j \in V_n$  is the length of the shortest path from  $i$  to  $j$  in  $G_n$ . Given a vertex  $i \in V_n$ , we let  $\mathsf{B}_i(t)$  be the set of vertices whose distance from  $i$  is at most  $t$ . With a slight abuse of notation,  $\mathsf{B}_i(t)$  will also denote the subgraph induced by those vertices. For  $i \in V_n$ , we let  $\partial i$  denote the set of its neighbors  $\partial i \equiv \{j \in V_n : (i, j) \in E_n\}$ , and  $|\partial i|$  its size (i.e. the degree of  $i$ ).

This paper is concerned by sequence of graphs  $\{G_n\}_{n \in \mathbb{N}}$  of diverging size, that converge locally to trees. Consider two trees  $T_1$  and  $T_2$  with vertices labeled arbitrarily. We shall write  $T_1 \simeq T_2$  if the two trees become identical when vertices are relabeled from 1 to  $|T_1| = |T_2|$ , in a breadth first fashion, and following lexicographic order among siblings.

**Definition 2.1.** *Considering a sequence of graphs  $\{G_n\}_{n \in \mathbb{N}}$ , let  $\mathbb{P}_n$  denote the law induced on the ball  $\mathsf{B}_i(t)$  in  $G_n$  centered at a uniformly chosen random vertex  $i \in [n]$ . We say that  $\{G_n\}$  converges locally to the random tree  $\mathsf{T}(P, \rho, \infty)$  if, for any  $t$ , and any rooted tree  $T$  with  $t$  generations*

$$\lim_{n \rightarrow \infty} \mathbb{P}_n \{\mathsf{B}_i(t) \simeq T\} = \mathbb{P}\{\mathsf{T}(P, \rho, t) \simeq T\}. \quad (2.3)$$

**Definition 2.2.** *We say that a sequence of graphs  $\{G_n\}_{n \in \mathbb{N}}$  is uniformly sparse if*

$$\lim_{l \rightarrow \infty} \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i \in V_n} |\partial i| \mathbb{I}(|\partial i| \geq l) = 0. \quad (2.4)$$

## 2.2 Free entropy

According to the statistical physics derivation [LV+02], the model (1.1) has a line of first order phase transitions for  $B = 0$  and  $\beta > \beta_c$  (that is, where the continuous function  $B \mapsto \phi(\beta, B)$  exhibits a discontinuous derivative). The critical temperature depends on the graph only through the average branching factor and is determined by the condition

$$\bar{\rho} \tanh \beta_c = 1. \quad (2.5)$$

Notice that  $\beta_c \simeq 1/\bar{\rho}$  for large degrees.

The asymptotic free-entropy density is given in terms of the fixed point of a distributional recursion. One characterization of this fixed point is as follows.

**Lemma 2.3.** *Consider the sequence of random variables  $\{h^{(t)}\}$  defined by  $h^{(0)} = 0$  identically and, for  $t \geq 0$ ,*

$$h^{(t+1)} \stackrel{d}{=} B + \sum_{i=1}^{K-1} \xi(\beta, h_i^{(t)}), \quad (2.6)$$

where  $K$  is an integer valued random variable of distribution  $\rho$ ,

$$\xi(\beta, h) \equiv \operatorname{atanh}[\tanh(\beta) \tanh(h)], \quad (2.7)$$

and the  $h_i^{(t)}$ 's are i.i.d. copies of  $h^{(t)}$  that are independent of  $K$ . If  $B > 0$  and  $\rho$  has finite first moment, then the distributions of  $h^{(t)}$  are stochastically monotone and  $h^{(t)}$  converges in distribution to the unique fixed point  $h^*$  of the recursion (2.6) that is supported on  $[0, \infty)$ .

Our next result confirms the statistical physics prediction for the free entropy density.

**Theorem 2.4.** *Let  $\{G_n\}_{n \in \mathbb{N}}$  be a sequence of uniformly sparse graphs that converges locally to  $\mathbb{T}(P, \rho, \infty)$ . If  $\rho$  has finite first moment (that is if  $P$  has finite second moment), then for any  $B \in \mathbb{R}$  and  $\beta \geq 0$  the following limit exists*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log Z_n(\beta, B) = \phi(\beta, B). \quad (2.8)$$

Moreover, for  $B > 0$  the limit is given by

$$\begin{aligned} \phi(\beta, B) &\equiv \frac{\bar{P}}{2} \log \cosh(\beta) - \frac{\bar{P}}{2} \mathbb{E} \log[1 + \tanh(\beta) \tanh(h_1) \tanh(h_2)] \\ &+ \mathbb{E} \log \left\{ e^B \prod_{i=1}^L [1 + \tanh(\beta) \tanh(h_i)] + e^{-B} \prod_{i=1}^L [1 - \tanh(\beta) \tanh(h_i)] \right\}, \end{aligned} \quad (2.9)$$

where  $L$  has distribution  $P_l$  and is independent of the ‘cavity fields’  $h_i$  that are i.i.d. copies of the fixed point  $h^*$  of Lemma 2.3. Also,  $\phi(\beta, B) = \phi(\beta, -B)$  and  $\phi(\beta, 0)$  is the limit of  $\phi(\beta, B)$  as  $B \rightarrow 0$ .

The proof of Theorem 2.4 is based on two steps

- (a) Reduce the computation of  $\phi_n(\beta, B) = \frac{1}{n} \log Z_n(\beta, B)$  to computing expectations of local (in  $G_n$ ) quantities with respect to the Boltzmann measure (1.1). This is achieved by noticing that the derivative of  $\phi_n(\beta, B)$  with respect to  $\beta$  is a sum of such expectations.
- (b) Show that expectations of local quantities on  $G_n$  are well approximated by the same expectations with respect to an Ising model on the associated tree  $\mathbb{T}(P, \rho, t)$  (for  $t$  and  $n$  large.) This is proved by showing that, on such a tree, local expectations are insensitive to boundary conditions that dominate stochastically free boundaries. The theorem then follows by monotonicity arguments.

The key step is of course the last one. A stronger requirement would be that these expectation values are insensitive to any boundary condition, which would coincide with uniqueness of the Gibbs measure on  $\mathbb{T}(P, \rho, \infty)$ . Such a requirement would allow for an elementary proof, but holds only at ‘high’ temperature,  $\beta \leq \beta_c$ .

Indeed, insensitivity to positive boundary conditions is proved in Section 4 for the following collection of trees of conditionally independent (and of bounded average) offspring numbers.

**Definition 2.5.** *An infinite tree  $\mathbb{T}$  rooted at the vertex  $\phi$  is called conditionally independent if for each integer  $k \geq 0$ , conditional on the subtree  $\mathbb{T}(k)$  of the first  $k$  generations of  $\mathbb{T}$ , the number of offspring  $\Delta_j$  for  $j \in \partial\mathbb{T}(k)$  are independent of each other, where  $\partial\mathbb{T}(k)$  denotes the set of vertices at generation  $k$ . We further assume that the (conditional on  $\mathbb{T}(k)$ ) first moments of  $\Delta_j$  are uniformly bounded by a given non-random finite constant  $\Delta$ .*

Beyond the random tree  $\mathbb{T}(P, \rho, \infty)$ , these include deterministic trees with bounded degrees and certain multi-type branching processes (such as random bipartite trees and percolation clusters on deterministic trees of bounded degree). Consequently, Theorem 2.4 extends to any uniformly sparse graph sequence that converge locally to a random tree  $\mathbb{T}$  of the form of Definition 2.5 except that the formula  $\phi(\beta, B)$  is in general more involved than the one given in (2.9). For example, such an extension allows one to handle uniformly random bipartite graphs with different degree distributions  $P_k$  and  $Q_k$  for the two types of vertices.

While we refrain from formalizing and proving such generalizations, we note in passing that our derivation of the formula (2.9) implicitly uses the fact that  $\mathbb{T}(P, \rho, \infty)$  possesses the involution invariance of [AS03]. As pointed out in [AL07], every local limit of finite graphs must have the involution invariance property (which clearly not every conditionally independent tree has).

### 2.3 Algorithmic implications

The free entropy density is not the only quantity that can be characterized for Ising models on locally tree-like graphs. Indeed local marginals can be efficiently computed with good accuracy. The basic idea is to solve a set of mean field equations iteratively. These are known as Bethe-Peierls or cavity equations and the corresponding algorithm is referred to as ‘belief propagation’ (BP).

More precisely, associate to each directed edge in the graph  $i \rightarrow j$ , with  $(i, j) \in G$ , a distribution  $\nu_{i \rightarrow j}(x_i)$  over  $x_i \in \{+1, -1\}$ . In the computer science literature these distributions are referred to as ‘messages’. They are updated as follows

$$\nu_{i \rightarrow j}^{(t+1)}(x_i) = \frac{1}{z_{i \rightarrow j}^{(t)}} e^{Bx_i} \prod_{l \in \partial i \setminus j} \sum_{x_l} e^{\beta x_i x_l} \nu_{l \rightarrow i}^{(t)}(x_l). \quad (2.10)$$

The initial conditions  $\nu_{i \rightarrow j}^{(0)}(\cdot)$  may be taken to be uniform or chosen according to some heuristic. We will say that the initial condition is *positive* if  $\nu_{i \rightarrow j}^{(0)}(+1) \geq \nu_{i \rightarrow j}^{(0)}(-1)$  for each of these messages.

Our next result concerns the uniform exponential convergence of the BP iteration to the same fixed point of (2.10), irrespective of its positive initial condition. Here and below, we denote by  $\|p - q\|_{\text{TV}}$  the total variation distance between distributions  $p$  and  $q$ .

**Theorem 2.6.** *Assume  $\beta \geq 0$ ,  $B > 0$  and  $G$  is a graph of finite maximal degree  $\Delta$ . Then, there exists  $A = A(\beta, B, \Delta)$  finite,  $\lambda = \lambda(\beta, B, \Delta) > 0$  and a fixed point  $\{\nu_{i \rightarrow j}^*\}$  of the BP iteration (2.10) such that for any positive initial condition  $\{\nu_{l \rightarrow k}^{(0)}\}$  and all  $t \geq 0$ ,*

$$\sup_{(i,j) \in E} \|\nu_{i \rightarrow j}^{(t)} - \nu_{i \rightarrow j}^*\|_{\text{TV}} \leq A \exp(-\lambda t). \quad (2.11)$$

For  $i_* \in V$  let  $U \equiv \mathbb{B}_{i_*}(r)$  be the ball of radius  $r$  around  $i_*$  in  $G$ , denoting by  $E_U$  its edge set, by  $\partial U$  its border (i.e. the set of its vertices at distance  $r$  from  $i_*$ ), and for each  $i \in \partial U$  let  $j(i)$  denote any one fixed neighbor of  $i$  in  $U$ .

Our next result shows that the probability distribution

$$\nu_U(\underline{x}_U) = \frac{1}{z_U} \exp \left\{ \beta \sum_{(i,j) \in E_U} x_i x_j + B \sum_{i \in U \setminus \partial U} x_i \right\} \prod_{i \in \partial U} \nu_{i \rightarrow j(i)}^*(x_i), \quad (2.12)$$

with  $\{\nu_{i \rightarrow j}^*(\cdot)\}$  the fixed point of the BP iteration per Theorem 2.6, is a good approximation for the marginal  $\mu_U(\cdot)$  of variables  $\underline{x}_U \equiv \{x_i : i \in U\}$  under the Ising model (1.1).

**Theorem 2.7.** *Assume  $\beta \geq 0$ ,  $B > 0$  and  $G$  is a graph of finite maximal degree  $\Delta$ . Then, there exist finite  $c = c(\beta, B, \Delta)$  and  $\lambda = \lambda(\beta, B, \Delta) > 0$  such that for any  $i_* \in G$  and  $U = \mathbb{B}_{i_*}(r)$ , if  $\mathbb{B}_{i_*}(t)$  is a tree then*

$$\|\mu_U - \nu_U\|_{\text{TV}} \leq \exp \{c^{r+1} - \lambda(t - r)\}. \quad (2.13)$$

## 2.4 Examples

Many common random graph ensembles [JLR00] naturally fit our framework.

*Random regular graphs.* Let  $G_n$  be a uniformly random graph with degree  $k$ . As  $n \rightarrow \infty$ , the sequence  $\{G_n\}$  is obviously uniformly sparse, and converges locally almost surely to the rooted infinite tree of degree  $k$  at every vertex. Therefore, in this case Theorem 2.4 applies with  $P_k = 1$  and  $P_i = 0$  for  $i \neq k$ . The distributional recursion (2.6) then evolves with a deterministic sequence  $h^{(t)}$  recovering the result of [GM07].

*Erdős-Renyi graphs.* Let  $G_n$  be a uniformly random graph with  $m = n\gamma$  edges over  $n$  vertices. The sequence  $\{G_n\}$  converges locally almost surely to a Galton-Watson tree with Poisson offspring distribution of mean  $2\gamma$ . This corresponds to taking  $P_k = (2\gamma)^k e^{-2\gamma}/k!$ . The same happens to classical variants of this ensemble. For instance, one can add an edge independently for each pair  $(i, j)$  with probability  $2\gamma/n$ , or consider a multi-graph with Poisson( $2\gamma/n$ ) edges between each pair  $(i, j)$ .

The sequence  $\{G_n\}$  is with probability one uniformly sparse in each of these cases. Thus, Theorem 2.4 extends the results of [DG08] to arbitrary non-zero temperature and magnetic field.

*Arbitrary degree distribution.* Let  $P$  be a distribution with finite second moment and  $G_n$  a uniformly random graph with degree distribution  $P$  (more precisely, we set the number of vertices of degree  $k \geq 1$  to  $\lfloor nP_k \rfloor$ , adding one for  $k = 1$  if needed for an even sum of degrees). Then,  $\{G_n\}$  is uniformly sparse and with probability one it converges locally to  $\mathbb{T}(P, \rho, \infty)$ . The same happens if  $G_n$  is drawn according to the so-called configuration model (c.f. [Bo80]).

## 3 Preliminaries

We review here the notations and a couple of classical tools we use throughout this paper. To this end, when proving our results it is useful to allow for vertex-dependent magnetic fields  $B_i$ , that is, to replace the basic model (1.1) by

$$\mu(\underline{x}) = \frac{1}{Z(\beta, \underline{B})} \exp \left\{ \beta \sum_{(i,j) \in E} x_i x_j + \sum_{i \in V} B_i x_i \right\}. \quad (3.1)$$

Given  $U \subseteq V$ , we denote by  $(+)_U$  (respectively  $(-)_U$ ) the vector  $\{x_i = +1, i \in U\}$  (respectively,  $\{x_i = -1, i \in U\}$ ), dropping the subscript  $U$  whenever clear from the context. Further, we use  $\underline{x}_U \preceq \underline{x}'_U$

when two real valued vectors  $\underline{x}$  and  $\underline{x}'$  are such that  $x_i \leq x'_i$  for all  $i \in U$  and say that a distribution  $\rho_U(\cdot)$  over  $\mathbb{R}^U$  is dominated by a distribution  $\rho'_U(\cdot)$  over this set (denoted  $\rho_U \preceq \rho'_U$ ), if the two distributions can be coupled so that  $\underline{x}_U \preceq \underline{x}'_U$  for any pair  $(\underline{x}_U, \underline{x}'_U)$  drawn from this coupling. Finally, we use throughout the shorthand  $\langle \nu, f \rangle = \sum_x f(x)\nu(x)$  for a distribution  $\nu$  and function  $f$  on the same finite set, or  $\langle f \rangle$  when  $\nu$  is clear from the context.

The first classical result we need is Griffiths inequality (see [Lig85, Theorem IV.1.21]).

**Theorem 3.1.** *Consider two Ising models  $\mu(\cdot)$  and  $\mu'(\cdot)$  on graphs  $G = (V, E)$  and  $G' = (V, E')$ , inverse temperatures  $\beta$  and  $\beta'$ , and magnetic fields  $\{B_i\}$  and  $\{B'_i\}$ , respectively. If  $E \subseteq E'$ ,  $\beta \leq \beta'$  and  $0 \leq B_i \leq B'_i$  for all  $i \in V$ , then  $0 \leq \langle \mu, \prod_{i \in U} x_i \rangle \leq \langle \mu', \prod_{i \in U} x_i \rangle$  for any  $U \subseteq V$ .*

The second classical result we use is the GHS inequality (see [GHS70]) about the effect of the magnetic field  $\underline{B}$  on the local magnetizations at various vertices.

**Theorem 3.2** (Griffiths, Hurst, Sherman). *Let  $\beta \geq 0$  and for  $\underline{B} = \{B_i : i \in V\}$ , denote by  $m_j(\underline{B}) \equiv \mu(\{\underline{x} : x_j = +1\}) - \mu(\{\underline{x} : x_j = -1\})$  the local magnetization at vertex  $j$  in the Ising model (3.1). If  $B_i \geq 0$  for all  $i \in V$ , then for any three vertices  $j, k, l \in V$  (not necessarily distinct),*

$$\frac{\partial^2 m_j(\underline{B})}{\partial B_k \partial B_l} \leq 0. \quad (3.2)$$

Finally, we need the following elementary inequality

**Lemma 3.3.** *For any function  $f : \mathcal{X} \mapsto [0, f_{\max}]$  and distributions  $\nu, \nu'$  on the finite set  $\mathcal{X}$  such that  $\nu(f > 0) > 0$  and  $\nu'(f > 0) > 0$ ,*

$$\sum_x \left| \frac{\nu(x)f(x)}{\langle \nu, f \rangle} - \frac{\nu'(x)f(x)}{\langle \nu', f \rangle} \right| \leq \frac{3f_{\max}}{\max(\langle \nu, f \rangle, \langle \nu', f \rangle)} \|\nu - \nu'\|_{\text{TV}}. \quad (3.3)$$

*In particular, if  $0 < f_{\min} \leq f(x)$ , then the right hand side is bounded by  $(3f_{\max}/f_{\min})\|\nu - \nu'\|_{\text{TV}}$ .*

*Proof.* Assuming without loss of generality that  $\langle \nu', f \rangle \geq \langle \nu, f \rangle > 0$ , the left hand side of (3.3) can be bounded as

$$\begin{aligned} & \frac{1}{\langle \nu, f \rangle \langle \nu', f \rangle} \sum_x |\nu(x)f(x)\langle \nu', f \rangle - \nu'(x)f(x)\langle \nu, f \rangle| \leq \\ & \leq \frac{1}{\langle \nu', f \rangle} |\langle \nu, f \rangle - \langle \nu', f \rangle| + \frac{1}{\langle \nu', f \rangle} \sum_x |\nu(x)f(x) - \nu'(x)f(x)| \leq \\ & \leq \frac{f_{\max}}{\langle \nu', f \rangle} \|\nu - \nu'\|_{\text{TV}} + \frac{2f_{\max}}{\langle \nu', f \rangle} \|\nu - \nu'\|_{\text{TV}}. \end{aligned}$$

This implies the lemma. □

## 4 Ising models on trees

We prove in this section certain facts about Ising models on trees which are of independent interest and as a byproduct we deduce Lemma 2.3 and the theorems of Section 2.3. In doing so, recall that for each  $\ell \geq 1$

the Ising models on  $\mathbb{T}(\ell)$  with free and plus boundary conditions are

$$\mu^{\ell,0}(\underline{x}) \equiv \frac{1}{Z^{\ell,0}} \exp \left\{ \beta \sum_{(ij) \in \mathbb{T}(\ell)} x_i x_j + \sum_{i \in \mathbb{T}(\ell)} B_i x_i \right\}, \quad (4.1)$$

$$\mu^{\ell,+}(\underline{x}) \equiv \frac{1}{Z^{\ell,+}} \exp \left\{ \beta \sum_{(ij) \in \mathbb{T}(\ell)} x_i x_j + \sum_{i \in \mathbb{T}(\ell)} B_i x_i \right\} \mathbb{I}(\underline{x}_{\partial \mathbb{T}(\ell)} = (+)_{\partial \mathbb{T}(\ell)}). \quad (4.2)$$

Equivalently  $\mu^{\ell,0}$  is the Ising model (3.1) on  $\mathbb{T}(\ell)$  with magnetic fields  $\{B_i\}$  and  $\mu^{\ell,+}$  is the modified Ising model corresponding to the limit  $B_i \uparrow +\infty$  for all  $i \in \partial \mathbb{T}(\ell)$ . To simplify our notations we denote such limits hereafter simply by setting  $B_i = +\infty$  and use  $\mu^\ell$  for statements that apply to both free and plus boundary conditions.

We start with the following simple, but useful observation.

**Lemma 4.1.** *For a subtree  $U$  of a finite tree  $T$  let  $\partial_* U$  denote the subset of vertices of  $U$  connected by an edge to  $W \equiv T \setminus U$  and for each  $u \in \partial_* U$  let  $\langle x_u \rangle_W$  denote the root magnetization of the Ising model on the maximal subtree  $T_u$  of  $W \cup \{u\}$  rooted at  $u$ . The marginal on  $U$  of the Ising measure on  $T$ , denoted  $\mu_U^T$  is then an Ising measure on  $U$  with magnetic field  $B'_u = \text{atanh}(\langle x_u \rangle_W) \geq B_u$  for  $u \in \partial_* U$  and  $B'_u = B_u$  for  $u \notin \partial_* U$ .*

*Proof.* Since  $U$  is a subtree of the tree  $T$ , the subtrees  $T_u$  for  $u \in \partial_* U$  are disjoint. Therefore, with  $\hat{\mu}_u(\underline{x})$  denoting the Ising model distribution for  $T_u$  we have that

$$\mu_U^T(\underline{x}_U) = \frac{1}{\hat{Z}} f(\underline{x}_U) \prod_{u \in \partial_* U} \hat{\mu}_u(x_u), \quad (4.3)$$

for the Boltzmann weight

$$f(\underline{x}_U) = \exp \left\{ \beta \sum_{(uv) \in U} x_u x_v + \sum_{u \in U \setminus \partial_* U} B_u x_u \right\}.$$

Further,  $x_u \in \{+1, -1\}$  so for each  $u \in \partial_* U$  and some constants  $c_u$ ,

$$\hat{\mu}_u(x_u) = \frac{1}{2} (1 + x_u \langle x_u \rangle_W) = c_u \exp(\text{atanh}(\langle x_u \rangle_W) x_u).$$

Embedding the normalization constants  $c_u$  within  $\hat{Z}$  we thus conclude that  $\mu_U^T$  is an Ising measure on  $U$  with the stated magnetic field  $B'_u$ . Finally, comparing the root magnetization for  $T_u$  with that for  $\{u\}$  we have by Griffiths inequality that  $\langle x_u \rangle_W \geq \tanh(B_u)$ , as claimed.  $\square$

**Theorem 4.2.** *Suppose  $\mathbb{T}$  is a conditionally independent infinite tree of average offspring numbers bounded by  $\Delta$ , as in Definition 2.5. For  $0 < B_{\min} \leq B_{\max}$ ,  $\beta_{\max}$  and  $\Delta$  finite, there exist  $M = M(\beta_{\max}, B_{\min}, \Delta)$  and  $C = C(\beta_{\max}, B_{\max})$  finite such that if  $B_i \leq B_{\max}$  for all  $i \in \mathbb{T}(r-1)$  and  $B_i \geq B_{\min}$  for all  $i \in \mathbb{T}(\ell)$ ,  $\ell > r$ , then*

$$\mathbb{E} \|\mu_U^{\ell,+} - \mu_U^{\ell,0}\|_{\text{TV}} \leq \delta(\ell - r) \mathbb{E}\{C^{|\mathbb{T}(r)|}\}, \quad (4.4)$$

for  $\delta(t) = M/t$ , all  $U \subseteq \mathbb{T}(r)$  and  $\beta \leq \beta_{\max}$ .

*Proof.* Fixing  $\ell > r$  it suffices to consider  $U = \mathbb{T}(r)$  (for which the left side of (4.4) is maximal). For this  $U$  and  $T = \mathbb{T}(\ell)$  we have that  $\partial_* U = \partial \mathbb{T}(r)$  and  $U \setminus \partial_* U = \mathbb{T}(r-1)$ , where in this case the Boltzmann weight  $f(\cdot)$  in (4.3) is bounded above by  $f_{\max} = c^{|\mathbb{T}(r)|}$  and below by  $f_{\min} = 1/f_{\max}$  for  $c = \exp(\beta_{\max} + B_{\max})$ . Further, the plus and free boundary conditions then differ in (4.3) by having the corresponding boundary conditions at generation  $\ell - r$  of each subtree  $T_u$ , which we distinguish by using  $\hat{\mu}_u^{+/0}(x_u)$  instead of  $\hat{\mu}_u(x_u)$ . Since the total variation distance between two product measures is at most the sum of the distance between their marginals, upon applying Lemma 3.3 we deduce from (4.3) that

$$\|\mu_{\mathbb{T}(r)}^{\ell,+} - \mu_{\mathbb{T}(r)}^{\ell,0}\|_{\text{TV}} \leq \frac{3}{2} c^{2|\mathbb{T}(r)|} \sum_{i \in \partial \mathbb{T}(r)} |\hat{\mu}_i^+(x_i = 1) - \hat{\mu}_i^0(x_i = 1)|.$$

By our assumptions, conditional on  $U = \mathbb{T}(r)$ , the subtrees  $T_i$  of  $T = \mathbb{T}(\ell)$  denoted hereafter also by  $\mathbb{T}_i$  are for  $i \in \partial \mathbb{T}(r)$  independent of each other. Further,  $2\hat{\mu}_i^{+/0}(x_i = 1) - 1$  is precisely the magnetization of their root vertex under plus/free boundary conditions at generation  $\ell - r$ . Thus, taking  $C = ec^2$  (and using the inequality  $y \leq e^y$ ), it suffices to show that the magnetizations  $m^{\ell,+/0}(\underline{B}) = \langle \mu^{\ell,+/0}, x_\phi \rangle$  at the root of any such conditionally independent infinite tree  $\mathbb{T}$  satisfy  $\mathbb{E}\{m^{\ell,+}(\underline{B}) - m^{\ell,0}(\underline{B})\} \leq \frac{M}{\ell}$ , for some  $M = M(\beta_{\max}, B_{\min}, \Delta)$  finite, all  $\beta \leq \beta_{\max}$  and  $\ell \geq 1$ , where we have removed the absolute value since  $m^{\ell,+}(\underline{B}) \geq m^{\ell,0}(\underline{B})$  by Griffiths inequality. For greater convenience of the reader, this fact is proved in the next lemma.  $\square$

**Lemma 4.3.** *Suppose  $\mathbb{T}$  is a conditionally independent infinite tree of average offspring numbers bounded by  $\Delta$ . For  $0 < B_{\min} \leq B_{\max}$ ,  $\beta_{\max}$  and  $\Delta$  finite, there exist  $M = M(\beta_{\max}, B_{\min}, \Delta)$  such that*

$$\mathbb{E}\{m^{\ell,+}(\underline{B}) - m^{\ell,0}(\underline{B})\} \leq \frac{M}{\ell}, \quad (4.5)$$

where  $m^{\ell,+/0}(\underline{B}) = \langle \mu^{\ell,+/0}, x_\phi \rangle$  are the root magnetizations under + and free boundary condition on  $\mathbb{T}$ .

*Proof.* Note that (4.5) trivially holds for  $\beta = 0$  (in which case  $\mu^{\ell,+}(x_\phi) = \mu^{\ell,0}(x_\phi)$ ). Assuming hereafter that  $\beta > 0$  we proceed to prove (4.5) when each vertex of  $\mathbb{T}(\ell - 1)$  has a non-zero offspring number. To this end, for  $\underline{H} = \{H_i \in \mathbb{R} : i \in \partial \mathbb{T}(k)\}$  let

$$\mu^{k,\underline{H}}(\underline{x}) \equiv \frac{1}{Z^{k,0}} \exp \left\{ \beta \sum_{(ij) \in \mathbb{T}(k)} x_i x_j + \sum_{i \in \mathbb{T}(k)} B_i x_i + \sum_{i \in \partial \mathbb{T}(k)} H_i x_i \right\}$$

and denote by  $m^k(\underline{B}, \underline{H})$  the corresponding root magnetization. Writing  $H$  instead of  $\underline{H}$  for constant magnetic field on the leaf nodes, that is, when  $H_i = H$  for each  $i \in \partial \mathbb{T}(k)$ , we note that  $m^{k,+}(\underline{B}) = m^k(\underline{B}, \infty)$  and  $m^{k,0}(\underline{B}) = m^k(\underline{B}, 0)$ . Further, applying Lemma 4.1 for the subtree  $\mathbb{T}(k-1)$  of  $\mathbb{T}(k)$  we represent  $m^k(\underline{B}, \infty)$  as the root magnetization  $m^{k-1}(\underline{B}', 0)$  on  $\mathbb{T}(k-1)$  where  $B'_i = B_i + \beta \Delta_i$  for  $i \in \partial \mathbb{T}(k-1)$  and  $B'_i = B_i$  for all other  $i$ . Consequently,

$$m^k(\underline{B}, \infty) = m^{k-1}(\underline{B}, \{\beta \Delta_i\}). \quad (4.6)$$

Recall that if  $\frac{\partial^2 g}{\partial z_i^2} \leq 0$  for  $i = 1, \dots, s$ , then applying Jensen's inequality one variable at a time we have that  $\mathbb{E}g(Z_1, \dots, Z_s) \leq g(\mathbb{E}Z_1, \dots, \mathbb{E}Z_s)$  for any independent random variables  $Z_1, \dots, Z_s$ . By the GHS inequality, this is the case for  $\underline{H} \mapsto m^{k-1}(\underline{B}, \underline{H})$ , hence with  $\mathbb{E}_k$  denoting the conditional on  $\mathbb{T}(k)$  expectation over the independent offspring numbers  $\Delta_i$  for  $i \in \partial \mathbb{T}(k)$ , we deduce that

$$\mathbb{E}_{k-1} m^k(\underline{B}, \infty) \leq m^{k-1}(\underline{B}, \{\beta \mathbb{E}_{k-1} \Delta_i\}) \leq m^{k-1}(\underline{B}, \beta \Delta), \quad (4.7)$$

where the last inequality is a consequence of Griffiths inequality and our assumption that  $\mathbb{E}_t \Delta_i \leq \Delta$  for any  $i \in \partial\mathbb{T}(t)$  and all  $t \geq 0$ . Since each  $i \in \partial\mathbb{T}(k-1)$  has at least one offspring whose magnetic field is at least  $B_{\min}$ , it follows by Griffiths inequality that  $m^{k,0}(\underline{B})$  is bounded below by the magnetization at the root of the subtree  $T$  of  $\mathbb{T}(k)$  where  $\Delta_i = 1$  for all  $i \in \partial\mathbb{T}(k-1)$  and  $B_i = B_{\min}$  for all  $i \in \partial\mathbb{T}(k)$ . Applying Lemma 4.1 for  $T$  and  $U = \mathbb{T}(k-1)$ , the root magnetization for the Ising distribution on  $T$  turns out to be precisely  $m^{k-1}(\underline{B}, \xi)$  for  $\xi = \xi(\beta, B_{\min}) > 0$  of (2.7). Thus, one more application of Griffiths inequality yields that

$$m^k(\underline{B}, 0) \geq m^{k-1}(\underline{B}, \xi) \geq m^{k-1}(\underline{B}, 0). \quad (4.8)$$

Next note that  $\xi(\beta, B) \leq \beta \leq \beta\Delta$  and by GHS inequality  $H \mapsto m^{k-1}(\underline{B}, H)$  is concave. Hence,

$$m^{k-1}(\underline{B}, \beta\Delta) - m^{k-1}(\underline{B}, 0) \leq M[m^{k-1}(\underline{B}, \xi) - m^{k-1}(\underline{B}, 0)], \quad (4.9)$$

for the finite constant

$$M \equiv \sup_{0 < \beta \leq \beta_{\max}} \frac{\beta\Delta}{\xi(\beta, B_{\min})}$$

and all  $\beta \leq \beta_{\max}$ . Combining (4.7), (4.8) and (4.9) we obtain that

$$\begin{aligned} \mathbb{E}_{k-1}\{m^{k,+}(\underline{B}) - m^{k,0}(\underline{B})\} &\leq m^{k-1}(\underline{B}, \Delta\beta) - m^{k-1}(\underline{B}, 0) \\ &\leq M[m^{k-1}(\underline{B}, \xi) - m^{k-1}(\underline{B}, 0)] \leq M[m^k(\underline{B}, 0) - m^{k-1}(\underline{B}, 0)]. \end{aligned}$$

We have seen in (4.8) that  $k \mapsto m^{k,0}(\underline{B})$  is non-decreasing whereas from (4.6) and Griffiths inequality we have that  $k \mapsto m^{k,+}(\underline{B})$  is non-increasing. With magnetization bounded above by one, we thus get upon summing the preceding inequalities for  $k = 1, \dots, \ell$  that

$$\ell \mathbb{E}_{\ell-1}[m^{\ell,+}(\underline{B}) - m^{\ell,0}(\underline{B})] \leq \sum_{k=1}^{\ell} \mathbb{E}_{k-1}[m^{k,+}(\underline{B}) - m^{k,0}(\underline{B})] \leq M,$$

from which we deduce (4.5).

Considering now the general case where the infinite tree  $\mathbb{T}$  has vertices (other than the root) of degree one, let  $\mathbb{T}^*(\ell)$  denote the ‘backbone’ of  $\mathbb{T}(\ell)$ , that is, the subtree induced by vertices along self-avoiding paths between  $\emptyset$  and  $\partial\mathbb{T}(\ell)$ . Taking  $U = \mathbb{T}^*(\ell)$  as the subtree of  $T = \mathbb{T}(\ell)$  in Lemma 4.1, note that for each  $u \in \partial_* U$  the subtree  $T_u$  contains no vertex from  $\partial\mathbb{T}(\ell)$ . Consequently, the marginal measures  $\mu_U^{\ell,+/0}$  are Ising measures on  $U$  with the same magnetic fields  $B'_i \geq B_i \geq B_{\min}$  outside  $\partial\mathbb{T}(\ell)$ . Thus, with  $m_*^{\ell,+/0}(\underline{B})$  denoting the corresponding magnetizations at the root for  $\mathbb{T}^*(\ell)$ , we deduce that  $m^{\ell,+/0}(\underline{B}) = m_*^{\ell,+/0}(\underline{B}')$  where  $B'_i \geq B_i \geq B_{\min}$  for all  $i$ . By definition every vertex of  $\mathbb{T}^*(\ell-1)$  has a non-zero offspring number and with  $B'_i \geq B_{\min}$ , the required bound

$$\mathbb{E}\{m^{\ell,+}(\underline{B}) - m^{\ell,0}(\underline{B})\} = \mathbb{E}\{m_*^{\ell,+}(\underline{B}') - m_*^{\ell,0}(\underline{B}')\} \leq \frac{M}{\ell}$$

follows by the preceding argument, since  $\mathbb{T}^*(\ell)$  is a conditionally independent tree whose offspring numbers  $\Delta_i^* \geq 1$  do not exceed those of  $\mathbb{T}(\ell)$ . Indeed, for  $k = 0, 1, \dots, \ell-1$ , given  $\mathbb{T}^*(k)$  the offspring numbers at  $i \in \partial\mathbb{T}^*(k)$  are independent of each other (with probability of  $\{\Delta_i^* = s\}$  proportional to the sum over  $t \geq 0$  of the product of the probability of  $\{\Delta_i = s+t\}$  and that of precisely  $s$  out of the  $s+t$  offspring of  $i$  in  $\mathbb{T}(\ell)$  having a line of descendants that survives additional  $\ell-k-1$  generations, for  $s \geq 1$ ).  $\square$

Simon’s inequality, see [Sim80, Theorem 2.1] allows one to bound the (centered) two point correlation functions in ferromagnetic Ising models with zero magnetic field. We provide next its generalization to arbitrary magnetic field, in the case of Ising models on trees.

**Lemma 4.4.** *If edge  $(i, j)$  is on the unique path from  $\emptyset$  to  $k \in \mathbb{T}(\ell)$ , with  $j$  a descendant of  $i \in \partial\mathbb{T}(t)$ ,  $t \geq 0$ , then*

$$\langle x_\emptyset; x_k \rangle_\emptyset^{(\ell)} \leq \cosh^2(2\beta + B_i) \langle x_\emptyset; x_i \rangle_\emptyset^{(t)} \langle x_j; x_k \rangle_j^{(\ell)}, \quad (4.10)$$

where  $\langle \cdot \rangle_i^{(r)}$  denotes the expectation with respect to the Ising distribution  $\hat{\mu}_i(\cdot)$  on the subtree  $\mathbb{T}_i$  of  $i$  and all its descendants in  $\mathbb{T}(r)$  and  $\langle x; y \rangle \equiv \langle xy \rangle - \langle x \rangle \langle y \rangle$  denotes the centered two point correlation function.

*Proof.* It is not hard to check that if  $x, y, z$  are  $\{+1, -1\}$ -valued random variables with  $x$  and  $z$  conditionally independent given  $y$ , then

$$\langle x; z \rangle = \frac{\langle x; y \rangle \langle y; z \rangle}{1 - \langle y \rangle^2}. \quad (4.11)$$

In particular, under  $\mu^{\ell, 0}$  the random variables  $x_\emptyset$  and  $x_k$  are conditionally independent given  $y = x_i$  with

$$\left| \log \left( \frac{\mu^{\ell, 0}(x_i = +1)}{\mu^{\ell, 0}(x_i = -1)} \right) \right| \leq 2(|\partial i| \beta + B_i).$$

Hence, if  $j$  is the unique descendant of  $i$  then  $|\langle x_i \rangle_\emptyset^{(\ell)}| \leq \tanh(2\beta + B_i)$  and we get from (4.11) that

$$\langle x_\emptyset; x_k \rangle_\emptyset^{(\ell)} \leq c \langle x_\emptyset; x_i \rangle_\emptyset^{(\ell)} \langle x_i; x_k \rangle_\emptyset^{(\ell)}$$

for  $c = \cosh^2(2\beta + B_i)$ . Next note that  $\langle x; y \rangle \leq 1 - \langle y \rangle^2$  for any two  $\{+1, -1\}$ -valued random variables, and since  $x_i$  and  $x_k$  are conditionally independent given  $y = x_j$  it follows from (4.11) that  $\langle x_i; x_k \rangle_\emptyset^{(\ell)} \leq \langle x_j; x_k \rangle_\emptyset^{(\ell)}$ . Further, if  $\langle \cdot \rangle$  is the expectation with respect to an Ising measure for some (finite) graph  $G$  then for any  $u, v \in G$

$$\frac{\partial \langle x_v \rangle}{\partial B_u} = \langle x_v x_u \rangle - \langle x_v \rangle \langle x_u \rangle = \langle x_v; x_u \rangle. \quad (4.12)$$

From Lemma 4.1 we know that computing the marginal of the Ising distribution for  $T = \mathbb{T}(\ell)$  on a smaller subtree  $U = \mathbb{T}_j$  of interest has the effect of increasing its magnetic field. Thus, combining the identity (4.12) with GHS inequality, we see that reducing this field (i.e. restricting to  $U$  the original Ising distribution), increases the centered two point correlation function. That is,  $\langle x_j; x_k \rangle_\emptyset^{(\ell)} \leq \langle x_j; x_k \rangle_j^{(\ell)}$ . Similarly, considering Lemma 4.1 for  $U = \mathbb{T}(t)$  we also have that  $\langle x_\emptyset; x_i \rangle_\emptyset^{(\ell)} \leq \langle x_\emptyset; x_i \rangle_\emptyset^{(t)}$  which completes our thesis in case  $j$  is the unique descendant of  $i$ .

Turning to the general case, we compare the thesis of the lemma for  $\mathbb{T}(\ell)$  and the subtree  $U = \mathbb{T}'(\ell)$  obtained upon deleting the subtrees rooted at descendants of  $i$  (and the corresponding edges to  $i$ ) except for  $\mathbb{T}_j$ . While  $\langle x_\emptyset; x_i \rangle_\emptyset^{(t)}$  and  $\langle x_j; x_k \rangle_j^{(\ell)}$  are unchanged by this modification of the underlying tree (as the relevant subgraphs are not modified), we have from Lemma 4.1 that  $\mu_U^{\ell, 0}(\cdot)$  is an Ising measure on  $U$  identical to the original but for an increase in the magnetic field at  $i$ . In view of (4.12) and the GHS inequality, we thus deduce that the value of  $\langle x_\emptyset; x_k \rangle_\emptyset^{(\ell)}$  is smaller for the Ising model on  $\mathbb{T}(\ell)$  than for the one on  $\mathbb{T}'(\ell)$  and since in  $\mathbb{T}'(\ell)$  the vertex  $j$  is the unique descendant of  $i$ , we are done.  $\square$

Equipped with the preceding lemma we next establish the exponential decay of correlations and of the effect of boundary conditions in Theorem 4.2.

**Corollary 4.5.** *There exist  $A$  finite and  $\lambda$  positive, depending only on  $\beta_{\max}$ ,  $B_{\min}$ ,  $B_{\max}$ , and  $\Delta$  such that*

$$\mathbb{E} \left\{ \sum_{i \in \partial\mathbb{T}(r)} \langle x_\emptyset; x_i \rangle_\emptyset^{(\ell)} \right\} \leq A e^{-\lambda r} \quad (4.13)$$

for any  $r \leq \ell$  and if  $B_i \leq B_{\max}$  for all  $i \in \mathbb{T}(\ell - 1)$  then Theorem 4.2 holds for  $\delta(t) = A \exp(-\lambda t)$ .

**Remark.** Taking  $B_i \uparrow +\infty$  for  $i \in \partial\mathbb{T}(\ell)$ , note that (4.13) applies when  $\langle \cdot \rangle^{(\ell)}$  is with respect to  $\mu^{\ell,+}(\cdot)$ .

*Proof.* Starting with the proof of (4.13) take  $\ell = r$  for which the left side is maximal (as we have seen while proving Lemma 4.4). Then, denoting by  $\langle \cdot \rangle_{H_r}$  the expectation under the Ising measure on  $\mathbb{T}(r)$  with a magnetic field  $H_r$  added to  $\underline{B}$  at all vertices  $i \in \partial\mathbb{T}(r)$ , it follows from (4.12) that

$$\sum_{i \in \partial\mathbb{T}(r)} \langle x_\emptyset; x_i \rangle_\emptyset^{(r)} = \sum_{i \in \partial\mathbb{T}(r)} \frac{\partial \langle x_\emptyset \rangle}{\partial B_i} = \left. \frac{\partial \langle x_\emptyset \rangle_{H_r}}{\partial H_r} \right|_{H_r=0}.$$

By GHS inequality the latter derivative is non-increasing in  $H_r$ , whence

$$\sum_{i \in \partial\mathbb{T}(r)} \langle x_\emptyset; x_i \rangle_\emptyset^{(r)} \leq \frac{2}{B_{\min}} [\langle x_\emptyset \rangle_{H_r=0} - \langle x_\emptyset \rangle_{H_r=-B_{\min}/2}].$$

Let  $B'_i = B_i - B_{\min}/2$  if  $i \in \partial\mathbb{T}(r)$  and  $B'_i = B_i$  otherwise, so  $\langle x_\emptyset \rangle_{H_r=-B_{\min}/2} = m^{r,0}(\underline{B}')$ . Further, from Griffiths inequality also  $\langle x_\emptyset \rangle_{H_r=0} \leq \langle x_\emptyset \rangle_{H_r=\infty} = m^{r,+}(\underline{B}')$  and it follows that

$$\Gamma_r \equiv \mathbb{E} \left\{ \sum_{i \in \partial\mathbb{T}(r)} \langle x_\emptyset; x_i \rangle_\emptyset^{(r)} \right\} \leq \frac{2}{B_{\min}} \mathbb{E} \{ m^{r,+}(\underline{B}') - m^{r,0}(\underline{B}') \}. \quad (4.14)$$

In particular, setting  $c = \cosh^2(2\beta_{\max} + B_{\max})$ , in view of Lemma 4.3 we find that  $\Gamma_{d-1} \leq 1/(ec\Delta)$  for  $d = 1 + \lceil 2ec\Delta M(\beta_{\max}, B_{\min}/2, \Delta)/B_{\min} \rceil$ . Further, since  $\mathbb{T}$  is conditionally independent, the same proof shows that if  $t + d = r' \leq r$  and  $\mathbb{T}_j$  is the subtree of  $\mathbb{T}(r)$  of depth  $d - 1$  rooted at  $j \in \partial\mathbb{T}(t + 1)$  then

$$\mathbb{E}_{t+1} \left\{ \sum_{k \in \partial\mathbb{T}_j} \langle x_j; x_k \rangle_j^{(r')} \right\} \leq \frac{1}{ec\Delta}.$$

Considering the inequality (4.10) of Lemma 4.4 for  $t = r - d \equiv r_1$  and all  $k \in \partial\mathbb{T}(r)$  we find that

$$\begin{aligned} \Gamma_r &\leq c \mathbb{E} \left\{ \sum_{\substack{i \in \partial\mathbb{T}(t) \\ j \in \partial\mathbb{T}(t+1) \cap \partial i}} \langle x_\emptyset; x_i \rangle_\emptyset^{(t)} \mathbb{E}_{t+1} \left[ \sum_{k \in \partial\mathbb{T}_j} \langle x_j; x_k \rangle_j^{(r')} \right] \right\} \\ &\leq \frac{1}{e\Delta} \mathbb{E} \left\{ \sum_{i \in \partial\mathbb{T}(t)} \Delta_i \langle x_\emptyset; x_i \rangle_\emptyset^{(t)} \right\} \leq e^{-1} \Gamma_{r_1}. \end{aligned}$$

Iterating the preceding bound at  $r_s = r - sd$ , for  $s = 1, \dots, \lfloor r/d \rfloor$  and noting that by (4.14) we have the bound  $\Gamma_{r'} \leq 2/B_{\min}$  at the last step, we get the uniform in  $\beta \leq \beta_{\max}$  exponential decay of (4.13).

Next, recall that the rate  $\delta(t)$  in Theorem 4.2 is merely the rate in the bound (4.5). For  $k \equiv |\partial\mathbb{T}(\ell)|$  we choose uniformly and independently of everything else a one to one mapping  $i : \{1, \dots, k\} \mapsto \partial\mathbb{T}(\ell)$ , and let  $\underline{B}^{(s)}$  for  $s \geq 1$  denote the magnetic field configuration obtained when taking  $B_{i(j)} \uparrow +\infty$  for all  $j \leq s$  (with  $\underline{B}^{(0)} = \underline{B}$ ). Since

$$m^{\ell,+}(\underline{B}) - m^{\ell,0}(\underline{B}) = \sum_{s=0}^{k-1} [m^{\ell,0}(\underline{B}^{(s+1)}) - m^{\ell,0}(\underline{B}^{(s)})],$$

we get the rate  $\delta(t) = A \exp(-\lambda t)$  from (4.13) as soon as we show that for  $i = i(s+1)$  and  $s = 0, \dots, k-1$ ,

$$m^{\ell,0}(\underline{B}^{(s+1)}) - m^{\ell,0}(\underline{B}^{(s)}) \leq \langle x_\emptyset; x_i \rangle_\emptyset^{(\ell)}. \quad (4.15)$$

To this end, let  $\langle \cdot \rangle_s$  denote the expectation under  $\mu^{\ell,0}$  with magnetic field  $\underline{B}^{(s)}$  so  $m^{\ell,0}(\underline{B}^{(s)}) = \langle x_\emptyset \rangle_s$ . Further, fixing  $i = i(s+1)$

$$m^{\ell,0}(\underline{B}^{(s+1)}) = \frac{\langle x_\emptyset \mathbb{I}(x_i = 1) \rangle_s}{\langle \mathbb{I}(x_i = 1) \rangle_s} = \frac{\langle x_\emptyset x_i \rangle_s + \langle x_\emptyset \rangle_s}{1 + \langle x_i \rangle_s}$$

(since  $\mathbb{I}(x_i = 1) = (1 + x_i)/2$ ). Since  $\langle x_i \rangle_s \geq 0$  by Griffiths inequality, it follows that

$$m^{\ell,0}(\underline{B}^{(s+1)}) - m^{\ell,0}(\underline{B}^{(s)}) \leq \langle x_\emptyset x_i \rangle_s - \langle x_\emptyset \rangle_s \langle x_i \rangle_s = \frac{\partial m_\emptyset(\underline{B}^{(s)})}{\partial B_i},$$

which by GHS inequality is maximal at  $s = 0$ , yielding (4.15) and completing the proof.  $\square$

As promised, Lemma 2.3 follows from the preceding results.

**Proof of Lemma 2.3.** Consider the Galton-Watson tree  $\mathbb{T}(\rho, \infty)$  of Section 2.1 and the corresponding Ising models  $\mu^{t,+/\circ}(\underline{x})$  of constant magnetic field  $B_i = B > 0$  on the subtrees  $\mathbb{T}(\rho, t)$ . It is easy to check that the random variables  $h^{(t)} = \text{atanh}(m^{t,0}(B))$  satisfy the distributional recursion (2.6) starting at  $h^{(0)} = 0$ . By Griffiths inequality  $m^{t,0}(B)$ , hence  $h^{(t)}$ , is non-decreasing in  $t$ , and so converges almost surely as  $t \rightarrow \infty$  to a limiting random variable  $h^*$ . Further, the bounds  $0 = h^{(0)} \leq h^{(t)} \leq B + \Delta_\emptyset$  hold for all  $t$  and hence also for  $h^*$ . We thus deduce that the distributions  $Q_t$  of  $h^{(t)}$  as determined by (2.6) are stochastically monotone (in  $t$ ) and converge weakly to some law  $Q^*$  of  $h^*$  that is supported on  $[0, \infty)$ .

Next, recall that for any fixed  $k$  and  $F(\cdot)$  continuous and bounded on  $\mathbb{R}^k$ , the functional  $\Psi_F(Q) = \int F(h_1, \dots, h_k) dQ(h_1) \cdots dQ(h_k)$  is continuous with respect to weak convergence of probability measures on  $[0, \infty)$  (for example, see [DZ98, Lemma 7.3.12]). Fixing  $g : \mathbb{R} \mapsto [-C, C]$  continuous, clearly

$$g_j(h_1, \dots, h_j) = g\left(B + \sum_{i=1}^{j-1} \xi(\beta, h_i)\right)$$

are continuous and bounded. Further, it follows from (2.6) that for all  $t$

$$\left| \int g dQ_{t+1} - \sum_{j=1}^k \mathbb{P}(K = j) \Psi_{g_j}(Q_t) \right| \leq C \mathbb{P}(K > k).$$

Taking  $t \rightarrow \infty$  followed by  $k \rightarrow \infty$ , we deduce by the preceding arguments (and the uniform boundedness  $|\Psi_{g_j}(Q^*)| \leq C$  for all  $j$ ), that

$$\int g dQ^* = \sum_{j=1}^{\infty} \mathbb{P}(K = j) \Psi_{g_j}(Q^*).$$

As this applies for every bounded continuous function  $g(\cdot)$ , we conclude that  $h^*$  and its law  $Q^*$  are a fixed point of the distributional recursion (2.6).

Next note that the random variables  $h_+^{(t)} = \text{atanh}[m^{t,+}(B)]$  form a non-increasing sequence that satisfies the same distributional recursion, but with the initial condition  $h_+^{(0)} = +\infty$ . Consequently, by the same arguments we have used before, the laws  $Q_{t,+}$  of  $h_+^{(t)}$  converge weakly to some fixed point  $Q_+^*$  of (2.6) that is also supported on  $[0, \infty)$ . Further,  $Q_t \preceq Q^{**} \preceq Q_{t,+}$  for  $t = 0$  and any (other) possible law  $Q^{**}$  of a fixed point  $h^{**}$  of (2.6) that is supported on  $[0, \infty)$ . Coupling so as to have the same value of  $K$ , evidently the recursion (2.6) preserves this stochastic order, which thus applies for all  $t$ . In the limit  $t \rightarrow \infty$  we thus deduce that  $Q^* \preceq Q^{**} \preceq Q_+^*$ . Since  $\rho$  has finite first moment, by (4.5) of Theorem 4.2,  $\mathbb{E}|\tanh(h_+^{(t)}) - \tanh(h^{(t)})| \rightarrow 0$  as  $t \rightarrow \infty$ . Thus, the expectation of the monotone increasing continuous

and bounded function  $\tanh(h)$  is the same under both  $Q^*$  and  $Q_+^*$ . Necessarily this is also the expectation of  $\tanh(h)$  under  $Q^{**}$  and the uniqueness of the non-negative fixed point of (2.6) follows.  $\square$

We next control the dependence on  $\beta$  of the distribution of the fixed point  $h^*$  from Lemma 2.3.

**Lemma 4.6.** *Let  $\|X - Y\|_{\text{MK}}$  denote the Monge-Kantorovich-Wasserstein distance between given laws of random variables  $X$  and  $Y$  (that is, the infimum of  $\mathbb{E}|X - Y|$  over all couplings of  $X$  and  $Y$ ). For any  $B > 0$  and  $\beta_{\max}$  finite there exists a constant  $C = C(\beta_{\max}, B)$  such that if  $h_{\beta_1}^*, h_{\beta_2}^*$  are the fixed points of the recursion (2.6) for  $0 \leq \beta_1, \beta_2 \leq \beta_{\max}$ , then*

$$\|\tanh(h_{\beta_2}^*) - \tanh(h_{\beta_1}^*)\|_{\text{MK}} \leq C |\beta_2 - \beta_1|. \quad (4.16)$$

*Proof.* Fixing a random tree  $\mathbb{T} = \mathbb{T}(\rho, \infty)$  of degree distribution  $\rho$ , recall that while proving Lemma 2.3 we provided a coupling of the random variables  $\tanh(h_\beta^*)$  and the Ising root magnetizations  $m^{t,+/\circ}(\beta, B)$  at  $\beta$  such that

$$m^{t,0}(\beta, B) \leq \tanh(h_\beta^*) \leq m^{t,+}(\beta, B)$$

for each  $\beta$  and all  $t$ . By Griffiths inequality the magnetizations at the root are non-decreasing in  $\beta$  so from the bound (4.5) we get that for  $M = M(\beta_{\max}, B, \bar{\rho})$  and any  $\beta_1 \leq \beta_2 \leq \beta_{\max}$ ,

$$\begin{aligned} \mathbb{E}|\tanh(h_{\beta_2}^*) - \tanh(h_{\beta_1}^*)| &\leq \mathbb{E}m^{t,0}(\beta_2, B) - \mathbb{E}m^{t,0}(\beta_1, B) + \frac{M}{t} \\ &\leq (\beta_2 - \beta_1) \sup_{\beta \leq \beta_{\max}} \mathbb{E}\left\{\frac{\partial m^{t,0}}{\partial \beta}\right\} + \frac{M}{t}, \end{aligned}$$

where the expectations are over the random tree  $\mathbb{T}(\rho, \infty)$ . Considering  $t \rightarrow \infty$  it thus suffices to show that  $\mathbb{E}[\partial m^{\ell,0}/\partial \beta]$  is bounded, uniformly in  $\ell$  and  $\beta \leq \beta_{\max}$ . To this end, a straightforward calculation yields

$$\frac{\partial m^{\ell,0}}{\partial \beta}(\beta, B) = \sum_{(i,j) \in \mathbb{T}(\ell)} (\langle x_\emptyset x_i x_j \rangle - \langle x_\emptyset \rangle \langle x_i x_j \rangle),$$

with  $\langle \cdot \rangle$  denoting the expectation with respect to the Ising measure  $\mu^{\ell,0}$ . If  $i$  is on the path in  $\mathbb{T}(\ell)$  between the root and  $j$ , then under the measure  $\mu^{\ell,0}$  the variables  $x_\emptyset$  and  $x_j$  are conditionally independent given  $x_i$ . Further, as  $x_i \in \{-1, 1\}$  it is easy to check that in this case

$$\langle x_\emptyset x_i x_j \rangle - \langle x_\emptyset \rangle \langle x_i x_j \rangle = \gamma \langle x_\emptyset; x_i \rangle,$$

where  $\gamma$  is the arithmetic mean of the conditional expected value of  $x_j$  for  $x_i = -1$  and the conditional expected value of  $x_j$  for  $x_i = 1$ . Thus,  $|\gamma| \leq 1$  and recalling (4.12) that  $\langle x_\emptyset; x_i \rangle$  is non-negative by Griffiths inequality, we deduce that

$$\frac{\partial m^{\ell,0}}{\partial \beta}(\beta, B) \leq \sum_{i \in \mathbb{T}(\ell-1)} \Delta_i \langle x_\emptyset; x_i \rangle = \sum_{r=0}^{\ell-1} V_{r,\ell},$$

where  $\Delta_i$  denotes the offspring number at  $i \in \mathbb{T}$  and by (4.12)

$$V_{r,\ell} \equiv \sum_{i \in \partial \mathbb{T}(r)} \Delta_i \langle x_\emptyset; x_i \rangle = \sum_{i \in \partial \mathbb{T}(r)} \Delta_i \partial_{B_i} m^\ell(\underline{B}, 0) \Big|_{\underline{B}=B}$$

(with  $m^k(\underline{B}, \underline{H})$  the root magnetization for the measure  $\mu^{\underline{B}, \underline{H}}$  of (4.6)). In view of Lemma 4.1 we have that  $m^k(\underline{B}, 0) = m^{k-1}(\underline{B}, \underline{H})$  for some non-negative vector  $\underline{H}$ . By GHS inequality we deduce that for any  $i \in \mathbb{T}(k-1)$

$$\partial_{B_i} m^k(\underline{B}, 0) = \partial_{B_i} m^{k-1}(\underline{B}, \underline{H}) \leq \partial_{B_i} m^{k-1}(\underline{B}, 0).$$

Consequently,  $V_{r,\ell}$  is non-increasing in  $\ell$  and

$$\mathbb{E}\left[\frac{\partial m^{\ell,0}}{\partial \beta}\right] \leq \sum_{r=0}^{\ell-1} \mathbb{E}V_{r,\ell} \leq \sum_{r=0}^{\ell-1} \mathbb{E}V_{r,r} \leq \sum_{r=0}^{\infty} \mathbb{E}V_{r,r}.$$

Further,  $m^r(\underline{B}, 0)$  is independent of the offspring numbers at  $\partial\mathbb{T}(r)$  whose expectation with respect to the random tree  $\mathbb{T}(\rho, \infty)$  is  $\bar{\rho}$ . Thus, applying (4.13) of Corollary 4.5 for  $\ell = r$ ,  $\mathbb{T} = \mathbb{T}(\rho, \infty)$  and constant magnetic field, we find that for some  $A$  finite,  $\lambda > 0$ , any  $r \geq 0$  and all  $\beta \leq \beta_{\max}$

$$\mathbb{E}V_{r,r} = \bar{\rho} \mathbb{E}\left[\sum_{i \in \partial\mathbb{T}(r)} \partial_{B_i} m^r(\underline{B}, 0) |_{\underline{B}=B}\right] = \bar{\rho} \mathbb{E}\left[\sum_{i \in \partial\mathbb{T}(r)} \langle x_{\emptyset}; x_i \rangle\right] \leq \bar{\rho} A e^{-\lambda r}.$$

Summing over  $r$  gives us the required uniform boundedness of  $\mathbb{E}[\partial m^{\ell,0}/\partial \beta]$  in  $\ell$  and  $\beta \leq \beta_{\max}$ .  $\square$

## 5 Algorithms

The theorems stated in Section 2.3 are in fact consequences of Corollary 4.5.

**Proof of Theorem 2.6.** The proof is based on the well known representation of the iteration (2.10) in terms of ‘computation tree’ [TJ02]. Namely,  $\nu_{i \rightarrow j}^{(t)}(\cdot)$  coincides with the marginal at the root of the Ising model (1.1) on a properly constructed, deterministic tree  $\mathbb{T}_{i \rightarrow j}^c(t)$  of  $t$  generations. While we refer to the literature for the precise definition of  $\mathbb{T}_{i \rightarrow j}^c(t)$ , here are some immediate properties:

- (a) One can construct an infinite tree  $\mathbb{T}_{i \rightarrow j}^c(\infty)$  such that, for any  $t$ ,  $\mathbb{T}_{i \rightarrow j}^c(t)$  is the subtree formed by the first  $t$  generations of  $\mathbb{T}_{i \rightarrow j}^c(\infty)$ .
- (b) The maximal degree of  $\mathbb{T}_{i \rightarrow j}^c(\infty)$  is bounded by the maximal degree of  $G$  (and equal to the latter when  $G$  is connected).
- (c) A positive initialization corresponds to adding  $H_{l \rightarrow k} = \text{atanh}(\nu_{l \rightarrow k}^{(0)}(+1) - \nu_{l \rightarrow k}^{(0)}(-1))$  non-negative to the field  $B$  on the  $t$ -th generation vertices of  $\mathbb{T}_{i \rightarrow j}^c(t)$ .

Denote by  $\nu_{i \rightarrow j}^{+, (t)}(\cdot)$ ,  $\nu_{i \rightarrow j}^{0, (t)}(\cdot)$  the messages obtained under initializations  $\nu_{k \rightarrow l}^{+, (0)}(+1) = 1$  and  $\nu_{k \rightarrow l}^{0, (0)}(+1) = \nu_{k \rightarrow l}^{0, (0)}(-1) = 1/2$ , respectively. By Griffiths inequality,  $\nu_{i \rightarrow j}^{+, (t)}(+1)$  is non-increasing in  $t$ ,  $\nu_{i \rightarrow j}^{0, (t)}(+1)$  is non-decreasing in  $t$  and any positive initialization results with  $\nu_{i \rightarrow j}^{(t)}(\cdot)$  such that

$$\nu_{i \rightarrow j}^{+, (t)}(+1) \geq \nu_{i \rightarrow j}^{(t)}(+1) \geq \nu_{i \rightarrow j}^{0, (t)}(+1).$$

By Corollary 4.5 we have that  $\nu_{i \rightarrow j}^{+, (t)}(+1) - \nu_{i \rightarrow j}^{0, (t)}(+1) \leq A e^{-\lambda t}$  for all  $t \geq 0$ . Since  $A < \infty$  and  $\lambda > 0$  depend only on  $\beta$ ,  $B$  and the maximal degree of  $G$ , this immediately yields our thesis.  $\square$

**Proof of Theorem 2.7.** We use an additional property of the computation tree:

- (d) If  $\mathbb{B}_i(k)$  is a tree then  $\mathbb{T}_{i \rightarrow j}^c(k)$  is a tree rooted at  $i \rightarrow j$  whose vertices are the directed edges on the maximal subtree of  $\mathbb{B}_i(k)$  rooted at  $i$  that does not include  $j$ .

Without loss of generality we may and shall assume that  $t > r$ . For  $U = \mathbb{B}_{i_*}(r)$  consider the local marginal approximations  $\nu_U^+(\cdot)$ ,  $\nu_U^0(\cdot)$  defined as in (2.12) except that the fixed point messages  $\nu_{i \rightarrow j(i)}^*(\cdot)$  at  $i \in \partial\mathbb{B}_{i_*}(r)$  are replaced by those obtained after  $(t - r)$  iterations starting at  $\nu_{k \rightarrow l}^{+, (0)}(+1) = 1$  and

$\nu_{k \rightarrow l}^{0,(0)}(+1) = \nu_{k \rightarrow l}^{0,(0)}(-1) = 1/2$ , respectively. Since  $\mathbf{B}_{i_*}(t)$  is a tree, here  $j(i)$  is necessarily the neighbor of  $i$  on the path from  $i_*$  to  $i \in \partial \mathbf{B}_{i_*}(r)$  and from the preceding property (d) we see that  $\mathbf{T}_{i \rightarrow j(i)}^c(t-r)$  corresponds to the subtree of  $i$  and its lines of descendant in  $\mathbf{B}_{i_*}(t)$ . By property (c) we thus have that  $\nu_U^+(\cdot)$  and  $\nu_U^0(\cdot)$  are the marginals on  $U$  of the Ising model  $\nu^+$  on  $G$  with  $B_i = \infty$  at all  $i \notin \mathbf{B}_{i_*}(t)$  and the Ising model  $\nu^0$  on the vertices of  $G$  and the edges within the tree  $\mathbf{B}_{i_*}(t)$ . Such reasoning also shows that the probability measure  $\nu_U$  of (2.12) is the marginal on  $U$  of the Ising model  $\nu$  on vertices of  $G$  and edges of  $\mathbf{B}_{i_*}(t)$  with an additional non-negative magnetic field  $H_{l \rightarrow k} = \text{atanh}(\nu_{l \rightarrow k}^*(+1) - \nu_{l \rightarrow k}^*(-1))$  at  $\partial \mathbf{B}_{i_*}(t)$ . Consequently, with  $x_F \equiv \prod_{i \in F} x_i$  we have by Griffiths inequality that for any  $F \subseteq U$

$$\langle \nu^0, x_F \rangle \leq \langle \nu, x_F \rangle \leq \langle \nu^+, x_F \rangle, \quad \langle \nu^0, x_F \rangle \leq \langle \mu, x_F \rangle \leq \langle \nu^+, x_F \rangle,$$

and we deduce that for any  $F \subseteq U$ ,

$$|\langle \mu, x_F \rangle - \langle \nu, x_F \rangle| \leq \langle \nu^+, x_F \rangle - \langle \nu^0, x_F \rangle \leq 2 \|\nu_U^+ - \nu_U^0\|_{\text{TV}}.$$

Recall that since  $x_i \in \{-1, 1\}$ , for any possible value  $\underline{y} = \{y_i, i \in U\}$  of  $\underline{x}_U$ ,

$$\mathbb{I}(\underline{x}_U = \underline{y}) = 2^{-|U|} \prod_{i \in U} (1 + y_i x_i) = 2^{-|U|} \sum_{F \subseteq U} y_F x_F,$$

and with  $|y_F| \leq 1$  it follows that

$$|\mu_U(\underline{y}) - \nu_U(\underline{y})| = 2^{-|U|} \left| \sum_{F \subseteq U} y_F (\langle \mu_U, x_F \rangle - \langle \nu_U, x_F \rangle) \right| \leq \max_{F \subseteq U} |\langle \mu_U, x_F \rangle - \langle \nu_U, x_F \rangle| \leq 2 \|\nu_U^+ - \nu_U^0\|_{\text{TV}}.$$

This applies for any of the  $2^{|U|}$  possible values of  $\underline{x}_U$ , so

$$\|\mu_U(\cdot) - \nu_U(\cdot)\|_{\text{TV}} \leq 2^{|U|} \|\nu_U^+(\cdot) - \nu_U^0(\cdot)\|_{\text{TV}}.$$

Applying Corollary 4.5 for the deterministic tree  $\mathbf{B}_{i_*}(t)$  rooted at  $i_*$ , we get the bound (4.4) on the right side of the preceding inequality with  $\delta(k) = A \exp(-\lambda k)$ , some finite  $A$  and  $\lambda > 0$  that depend only on  $\beta$ ,  $B$  and  $\Delta$ . Thus, noting that  $|U| = |\mathbf{B}_{i_*}(r)| \leq \Delta^{r+1} + 1$  we establish our thesis upon choosing  $c = c(A, C, \Delta)$  large enough.  $\square$

## 6 From trees to graphs

We start with the following technical lemma.

**Lemma 6.1.** *Consider a convex set  $\mathcal{K} \subseteq \mathbb{R}$  and symmetric twice differentiable functions  $F_\ell : \mathcal{K}^\ell \rightarrow \mathbb{R}$  with  $F_0$  constant, such that for some finite constant  $c$ ,*

$$\sup_{\ell} \sup_{\mathcal{K}^\ell} \left| \frac{\partial^2 F_\ell}{\partial x_1 \partial x_2} \right| \leq 2c.$$

*Suppose i.i.d.  $X, X_i \in \mathcal{K}$  are such that  $\ell^{-1} \mathbb{E} |\partial_{x_1} F_\ell(x, X_2, \dots, X_\ell)|$  is bounded uniformly in  $\ell$  and  $x \in \mathcal{K}$  and the independent, square-integrable, non-negative integer valued random variable  $L$  satisfies*

$$\mathbb{E}[L \partial_{x_1} F_L(x, X_2, \dots, X_L)] = 0, \quad \forall x \in \mathcal{K}. \quad (6.1)$$

*Then, for any i.i.d.  $Y, Y_i \in \mathcal{K}$  also independent of  $L$ ,*

$$|\mathbb{E}[F_L(Y_1, \dots, Y_L) - F_L(X_1, \dots, X_L)]| \leq c \mathbb{E}[L(L-1)] \|X - Y\|_{\text{MK}}^2. \quad (6.2)$$

*Proof.* Our thesis trivially holds if either  $\|X - Y\|_{\text{MK}} = 0$  or  $\|X - Y\|_{\text{MK}} = \infty$ , so without loss of generality, fixing  $\gamma > 1$  we assume hereafter that  $(X_i, Y_i)$  are i.i.d. pairs, independent on  $L$  and coupled in such a way that  $\mathbb{E}|X_i - Y_i| \leq \gamma \|X - Y\|_{\text{MK}}$  is finite. It is easy to check that almost surely,

$$F_\ell(Y_1, \dots, Y_\ell) - F_\ell(X_1, \dots, X_\ell) = \sum_{i=1}^{\ell} \Delta_i F_\ell + \sum_{i \neq j}^{\ell} f_{ij}^{(\ell)}(Y_i - X_i)(Y_j - X_j), \quad (6.3)$$

where  $\Delta_i F_\ell = (Y_i - X_i) \int_0^1 \partial_{x_i} F_\ell(X_1, \dots, tY_i + (1-t)X_i, \dots, X_\ell) dt$  and each of the terms

$$f_{ij}^{(\ell)} = \int_0^1 \int_0^t \frac{\partial^2 F_\ell}{\partial x_i \partial x_j}(sY_1 + (1-s)X_1, \dots, tY_i + (1-t)X_i, \dots, sY_\ell + (1-s)X_\ell) ds dt,$$

is bounded by  $c$ . For i.i.d.  $(X_i, Y_i)$ , by the symmetry of the functions  $F_\ell$  with respect to their arguments, the assumed boundedness of  $\ell^{-1} \mathbb{E}|\partial_{x_1} F_\ell(x, X_2, \dots, X_\ell)|$  implies integrability of  $\Delta_i F_\ell$  with  $\mathbb{E}\Delta_i F_\ell$  independent of  $i$  and  $\ell^{-1} \mathbb{E}|\Delta_i F_\ell|$  uniformly bounded. This in turn implies the integrability of  $\sum_{i=1}^L \Delta_i F_L$  for any  $L$  square integrable and independent of  $(X_i, Y_i)$ , so by Fubini's theorem and our assumption (6.1),

$$\mathbb{E}\left[\sum_{i=1}^L \Delta_i F_L\right] = \mathbb{E}[L\Delta_1 F_L] = \mathbb{E}\left[(Y_1 - X_1) \int_0^1 \mathbb{E}[L\partial_{x_1} F_L(tY_1 + (1-t)X_1, X_2, \dots, X_L) | X_1, Y_1] dt\right] = 0.$$

Thus, considering the expectation of (6.3), by the uniform boundedness of  $f_{ij}^{(\ell)}$  and the independence of  $L$  on the i.i.d. pairs  $(X_i, Y_i)$ , we deduce that

$$\left| \mathbb{E}[F_L(Y_1, \dots, Y_L) - F_L(X_1, \dots, X_L)] \right| \leq c \mathbb{E} \sum_{i \neq j}^L |Y_i - X_i| |Y_j - X_j| \leq \gamma^2 c \mathbb{E}[L(L-1)] \|X - Y\|_{\text{MK}}^2.$$

Finally, taking  $\gamma \downarrow 1$  yields the bound (6.2). □

**Remark 6.2.** *It is not hard to adapt the proof of the lemma so as to replace  $F_1 : \mathcal{K} \mapsto \mathbb{R}$  by  $0.5F_1(x, y)$  for a twice differentiable symmetric function  $F_1 : \mathcal{K}^2 \mapsto \mathbb{R}$ . Taking  $P_\ell = \mathbb{P}(L = \ell)$  the contribution of  $L = 1$  to the left side of (6.1) is then  $P_1 \mathbb{E}[\partial_{x_1} F_1(x, X_2)]$  and the bound (6.2) is modified to*

$$\left| \frac{P_1}{2} \mathbb{E}[F_1(Y_1, Y_2) - F_1(X_1, X_2)] + \sum_{\ell \geq 2} P_\ell \mathbb{E}[F_\ell(Y_1, \dots, Y_\ell) - F_\ell(X_1, \dots, X_\ell)] \right| \leq c \mathbb{E}[L^2] \|X - Y\|_{\text{MK}}^2. \quad (6.4)$$

Consider the functional  $h \mapsto \varphi_h$  that, given a random variable  $h$ , evaluates the right hand side of Eq. (2.9). It is not hard to check that  $\varphi_h$  is well-defined and finite for every random variable  $h$ . The following corollary of Lemma 6.1 plays an important role in the proof of Theorem 2.4.

**Corollary 6.3.** *There exist non-decreasing finite  $c(|\beta|)$  such that if  $\bar{\rho} < \infty$  and  $h^*$  is a fixed point of the distributional identity (2.6) for some  $\beta, B \in \mathbb{R}$  then*

$$|\varphi_h(\beta, B) - \varphi_{h^*}(\beta, B)| \leq c(|\beta|) \bar{P} \bar{\rho} \|\tanh(h) - \tanh(h^*)\|_{\text{MK}}^2. \quad (6.5)$$

*Proof.* Setting  $u = \tanh(\beta)$  so  $|u| < 1$ , we verify the conditions of Lemma 6.1 when  $X_i$  are i.i.d. copies of  $X = \tanh(h^*)$  and  $Y_i$  i.i.d. copies of  $Y = \tanh(h)$ , all of whom take values in  $\mathcal{K} = [-1, 1]$  and are

independent of the random variable  $L$ . We apply the lemma in this setting for the symmetric, twice differentiable functions

$$F_\ell(x_1, \dots, x_\ell) = -\frac{1}{(\ell-1)} \sum_{1 \leq i < j \leq \ell} \log(1 + ux_i x_j) + \log \left\{ e^B \prod_{i=1}^{\ell} (1 + ux_i) + e^{-B} \prod_{i=1}^{\ell} (1 - ux_i) \right\},$$

for  $\ell \geq 2$ , and as in Remark 6.2,

$$F_1(x_1, x_2) = -\log(1 + ux_1 x_2) + \log \{ e^B (1 + ux_1) + e^{-B} (1 - ux_1) \} + \log \{ e^B (1 + ux_2) + e^{-B} (1 - ux_2) \}.$$

Indeed, setting  $\psi(x, y) = uy/(1 + uxy)$  and for each  $\ell \geq 1$

$$g_\ell(x_2, \dots, x_\ell) = \tanh \left( B + \sum_{j=2}^{\ell} \operatorname{atanh}(ux_j) \right), \quad (6.6)$$

(so  $g_1 = \tanh(B)$ ), it is not hard to verify that  $\partial_{x_1} F_1(x_1, x_2) = \psi(x_1, g_1) - \psi(x_1, x_2)$  while for  $\ell \geq 2$

$$\partial_{x_1} F_\ell(x_1, \dots, x_\ell) = \psi(x_1, g_\ell(x_2, \dots, x_\ell)) - \frac{1}{\ell-1} \sum_{j=2}^{\ell} \psi(x_1, x_j). \quad (6.7)$$

In particular,  $g_\ell(\cdot)$  are differentiable functions from  $\mathcal{K}^{\ell-1}$  to  $\mathcal{K}$ , such that  $\partial_{x_2} g_\ell$  are uniformly bounded (by  $a = |u|/(1 - u^2)$ ) and  $\partial_y \psi(x, y)$  is uniformly bounded on  $\mathcal{K}^2$  (by  $b = |u|/(1 - |u|^2)$ ). Consequently,  $\partial_{x_1} F_\ell$  and  $\partial^2 F_\ell / \partial x_1 \partial x_2$  are also uniformly bounded (by  $2/(1 - |u|)$  and  $b(a+1) = 2c(|\beta|)$ , respectively). Further,  $h^*$  is a fixed point of (2.6), hence  $X_1 \stackrel{d}{=} g_K(X_2, \dots, X_K)$ . With  $X_i$  identically distributed and  $\bar{P}\rho_k = kP_k$  we thus find as required in (6.1) that

$$\begin{aligned} P_1 \mathbb{E}[\partial_{x_1} F_1(x, X_2)] + \sum_{k \geq 2} k P_k \mathbb{E}[\partial_{x_1} F_k(x, X_2, \dots, X_k)] \\ = \bar{P} \left\{ \sum_{k=1}^{\infty} \rho_k \mathbb{E}[\psi(x, g_k(X_2, \dots, X_k))] - \mathbb{E}\psi(x, X_1) \right\} = 0. \end{aligned} \quad (6.8)$$

Noting that  $\mathbb{E}[L^2] = \bar{P}\bar{\rho}$  our thesis is merely the bound (6.4) upon confirming that

$$\begin{aligned} \varphi_h &= F_0 + \frac{P_1}{2} \mathbb{E}F_1(Y_1, Y_2) + \sum_{\ell \geq 2} P_\ell \mathbb{E}F_\ell(Y_1, \dots, Y_\ell) \\ \varphi_{h^*} &= F_0 + \frac{P_1}{2} \mathbb{E}F_1(X_1, X_2) + \sum_{\ell \geq 2} P_\ell \mathbb{E}F_\ell(X_1, \dots, X_\ell), \end{aligned}$$

for some constant  $F_0$  and that both series are absolutely summable.  $\square$

Let  $\bar{\mathbb{T}}(\rho, \infty)$  denote the infinite random tree  $\rho$  obtained by ‘gluing’ two independent trees from the ensemble  $\mathbb{T}(\rho, \infty)$  through an extra edge  $e$  between their roots and considering  $e$  as the root of  $\bar{\mathbb{T}}(\rho, \infty)$  denote by  $\bar{\mathbb{T}}(\rho, t)$  the subtree formed by its first  $t$  generations (i.e. consisting of  $e$  and the corresponding two independent copies from  $\mathbb{T}(\rho, t)$ ). An alternative way to sample from  $\bar{\mathbb{T}}(\rho, \infty)$  is to have independent offspring number  $k-1$  with probability  $\rho_k$  at each end of the root edge  $e$  and thereafter independently sample from this offspring distribution at each revealed new node of the tree. Equipped with these notations we have the following consequence of the local convergence of the graph sequence  $\{G_n\}$ .

**Lemma 6.4.** *Suppose a uniformly sparse graph sequence  $\{G_n\}$  converges locally to the random tree  $\mathbb{T}(P, \rho, \infty)$ . Fixing a non-negative integer  $t$ , for each  $(i, j) \in E_n$  denote the subgraph of  $G_n$  induced by vertices at distance at most  $t$  from  $(i, j)$  by  $\mathbf{B}_{ij}(t)$ . Let  $F(\cdot)$  be a fixed, bounded function on the collection of all possible subgraphs that may occur as  $\mathbf{B}_{ij}(t)$ , such that  $F(T_1) = F(T_2)$  whenever  $T_1 \simeq T_2$ . Then,*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{(i,j) \in E_n} F(\mathbf{B}_{ij}(t)) = \frac{\bar{P}}{2} \mathbb{E}\{F(\bar{\mathbb{T}}(\rho, t))\}. \quad (6.9)$$

*Proof.* Denoting by  $\mathbb{E}_{(ij)}(\cdot)$  the expectation with respect to a uniformly chosen edge  $(i, j)$  in  $E_n$ , the left side of (6.9) is merely  $(|E_n|/n)\mathbb{E}_{(ij)}\{F(\mathbf{B}_{ij}(t))\}$ . A uniformly chosen edge can be sampled by first selecting a vertex  $i$  with probability proportional to its degree  $|\partial i|$  and then picking one of its neighbors  $j = j(i)$  uniformly. Thus, denoting by  $\mathbb{E}_n(\cdot)$  the expectation with respect to a uniformly chosen random vertex  $i \in [n]$ , we have that

$$\mathbb{E}_{(ij)}\{F(\mathbf{B}_{ij}(t))\} = \frac{\mathbb{E}_n\{|\partial i| F(\mathbf{B}_{ij(i)}(t))\}}{\mathbb{E}_n\{|\partial i|\}}.$$

Marking uniformly at random one offspring of  $\emptyset$  in  $\mathbb{T}(P, \rho, t+1)$  (as corresponding to  $j(i)$ ), let  $\mathbb{T}_*(t+1)$  denote the subtree induced by vertices whose distance from either  $\emptyset$  or its marked offspring is at most  $t$ . Since  $\mathbf{B}_{ij(i)}(t) \subseteq \mathbf{B}_i(t+1)$  and with probability  $q_{t,k} \rightarrow 1$  as  $k \rightarrow \infty$  the random tree  $\mathbb{T}(P, \rho, t+1)$  belongs to the finite collection of trees with  $t+1$  generations and maximal degree at most  $k$ , it follows by dominated convergence and the local convergence of  $\{G_n\}$  that for any fixed  $l$ ,

$$\lim_{n \rightarrow \infty} \mathbb{E}_n [|\partial i| \mathbb{I}(|\partial i| \leq l) F(\mathbf{B}_{ij(i)}(t))] = \mathbb{E}_\rho \{\Delta_\emptyset \mathbb{I}(\Delta_\emptyset \leq l) F(\mathbb{T}_*(t+1))\},$$

where  $\mathbb{E}_\rho(\cdot)$  and  $\Delta_\emptyset$  denote expectations and the degree of the root, respectively, in  $\mathbb{T}(P, \rho, \infty)$ . Similarly,

$$\lim_{n \rightarrow \infty} \mathbb{E}_n \{|\partial i| \mathbb{I}(|\partial i| \leq l)\} = \mathbb{E}_\rho \Delta_\emptyset \mathbb{I}(\Delta_\emptyset \leq l).$$

Further, by the uniform sparsity of  $\{G_n\}$ ,

$$\limsup_{n \rightarrow \infty} \left| \mathbb{E}_n [|\partial i| \mathbb{I}(|\partial i| > l) F(\mathbf{B}_{ij(i)}(t))] \right| \leq \|F\|_\infty \limsup_{n \rightarrow \infty} \mathbb{E}_n [|\partial i| \mathbb{I}(|\partial i| > l)],$$

goes to zero as  $l \rightarrow \infty$ . Since  $P$  has a finite first moment,  $\Delta_\emptyset$  is integrable, so by the preceding, upon taking  $l \rightarrow \infty$  we deduce by dominated convergence that

$$\lim_{n \rightarrow \infty} \mathbb{E}_{(ij)}\{F(\mathbf{B}_{ij}(t))\} = \frac{\mathbb{E}_\rho\{\Delta_\emptyset F(\mathbb{T}_*(t+1))\}}{\mathbb{E}_\rho\{\Delta_\emptyset\}}.$$

To complete the proof note that the right side of the last expression is precisely  $\mathbb{E}\{F(\bar{\mathbb{T}}(\rho, t))\}$  and we have also shown that  $2|E_n|/n = \mathbb{E}_n\{|\partial i|\} \rightarrow \mathbb{E}_\rho \Delta_\emptyset = \bar{P}$ .  $\square$

**Proof of Theorem 2.4.** Since  $\phi_n(\beta, B) \equiv \frac{1}{n} \log Z_n(\beta, B)$  is invariant under  $B \rightarrow -B$  and is uniformly (in  $n$ ) Lipschitz continuous in  $B$  with Lipschitz constant one, it suffices to fix  $B > 0$  and show that  $\phi_n(\beta, B)$  converges as  $n \rightarrow \infty$  to the predicted  $\varphi_{h^*}(\beta, B)$  of (2.9), whereby  $h^* = h_\beta^*$  is the unique fixed point of the recursion (2.6) that is supported on  $[0, \infty)$  (see Lemma 2.3).

This is obviously true for  $\beta = 0$  since  $\phi_n(0, B) = \log(2 \cosh B) = \varphi_h(0, B)$ . Next, denoting by  $\langle \cdot \rangle_n$  the expectation with respect to the Ising measure on  $G_n$  (at parameters  $\beta$  and  $B$ ), it is easy to see that

$$\partial_\beta \phi_n(\beta, B) = \frac{1}{n} \sum_{(i,j) \in E_n} \langle x_i x_j \rangle_n. \quad (6.10)$$

Clearly  $|\partial_\beta \phi_n(\beta, B)| \leq |E_n|/n$  is bounded by the uniform sparsity of  $\{G_n\}$  so it is enough to show that the expression in (6.10) converges to the partial derivative of  $\varphi_{h_\beta^*}(\beta, B)$  with respect to  $\beta$ . Turning to compute the latter derivative, by Lemma 4.6 and Corollary 6.3 we can ignore the dependence of  $h_\beta^*$  on  $\beta$ . That is, we simply compute the partial derivative in  $\beta$  of the expression (2.9) while considering (the law of)  $h_i$  to be fixed. Indeed, with notations  $u = \tanh(\beta)$  and  $X_i = \tanh(h_i)$  as in the derivation of Corollary 6.3, a direct computation leads by the exchangeability of  $X_i$  to

$$\partial_\beta \varphi(\beta, B) = \frac{\bar{P}}{2}u - \frac{\bar{P}}{2}(1 - u^2)\mathbb{E}[\psi(X_1, X_2)] + (1 - u^2)\mathbb{E}[L\psi(X_1, g_L(X_2, \dots, X_L))],$$

for  $\psi(x, y) = xy/(1 + uxy)$  and  $g_\ell(x_2, \dots, x_\ell)$  of (6.6). Further, the fixed point property (6.8) applies for any bounded measurable  $\psi(\cdot)$ , so we deduce that

$$\mathbb{E}[L\psi(X_1, g_L(X_2, \dots, X_L))] = \bar{P}\mathbb{E}[\psi(X_1, g_K(X_2, \dots, X_K))] = \bar{P}\mathbb{E}[\psi(X_1, X_2)].$$

Consequently, it is not hard to verify that

$$\partial_\beta \varphi(\beta, B) = \frac{\bar{P}}{2}\mathbb{E}\left\{\frac{u + X_1X_2}{1 + uX_1X_2}\right\} = \frac{\bar{P}}{2}\mathbb{E}\left[\langle x_i x_j \rangle_{\bar{\Gamma}}\right], \quad (6.11)$$

where  $\langle \cdot \rangle_{\bar{\Gamma}}$  denotes the expectation with respect to the Ising model

$$\mu_{\bar{\Gamma}}(x_i, x_j) = \frac{1}{z_{ij}} \exp\{\beta x_i x_j + H_i x_i + H_j x_j\},$$

on one edge  $(ij)$  and random magnetic fields  $H_i$  and  $H_j$  that are independent copies of  $h_\beta^*$ .

In comparison, fixing a positive integer  $t$ , by Griffiths inequality the correlation  $\langle x_i x_j \rangle_n$  lies between the correlations  $F_0(\mathbf{B}_{ij}(t)) \equiv \langle x_i x_j \rangle_{\mathbf{B}_{ij}(t)}^0$  and  $F_+(\mathbf{B}_{ij}(t)) \equiv \langle x_i x_j \rangle_{\mathbf{B}_{ij}(t)}^+$  for the Ising model on the subgraph  $\mathbf{B}_{ij}(t)$  with free and plus, respectively, boundary conditions at  $\partial\mathbf{B}_{ij}(t)$ . Thus, in view of (6.10)

$$\frac{1}{n} \sum_{(i,j) \in E_n} F_0(\mathbf{B}_{ij}(t)) \leq \partial_\beta \phi_n(\beta, B) \leq \frac{1}{n} \sum_{(i,j) \in E_n} F_+(\mathbf{B}_{ij}(t)),$$

and taking  $n \rightarrow \infty$  we get by Lemma 6.4 that

$$\frac{\bar{P}}{2}\mathbb{E}[F_0(\bar{\Gamma}(\rho, t))] \leq \liminf_{n \rightarrow \infty} \partial_\beta \phi_n(\beta, B) \leq \limsup_{n \rightarrow \infty} \partial_\beta \phi_n(\beta, B) \leq \frac{\bar{P}}{2}\mathbb{E}[F_+(\bar{\Gamma}(\rho, t))].$$

To compute  $F_{0/+}(\bar{\Gamma}(\rho, t))$  we first sum over the values of  $x_k$  for  $k \in \bar{\Gamma}(\rho, t) \setminus \{i, j\}$ . This has the effect of reducing  $F_{0/+}(\bar{\Gamma}(\rho, t))$  to a form of  $\langle x_i x_j \rangle_{\bar{\Gamma}}$ . Further, as shown in the proof of Lemma 2.3, we get  $F_{0/+}(\bar{\Gamma}(\rho, t))$  by setting for  $H_i$  and  $H_j$  two independent copies of the variables  $h^{(t)}$  and  $h_+^{(t)}$ , respectively, which converge in law to  $h_\beta^*$  when  $t \rightarrow \infty$ . We also saw there that the functional  $\Psi_U(\nu) = \mathbb{E}[\langle x_i x_j \rangle_{\bar{\Gamma}}]$  (for continuous and bounded  $U(H_i, H_j) = (u + \tanh(H_i) \tanh(H_j))/(1 + u \tanh(H_i) \tanh(H_j))$ ), is continuous with respect to the weak convergence of the law  $\nu$  of  $H_i$ . Consequently, by (6.11)

$$\lim_{t \rightarrow \infty} \frac{\bar{P}}{2}\mathbb{E}[F_{0/+}(\bar{\Gamma}(\rho, t))] = \partial_\beta \varphi(\beta, B),$$

which completes the proof of the theorem.  $\square$

## References

- [AL07] D. Aldous and R. Lyons, *Processes on unimodular random networks*, *Elect. J. Probab.*, **12** (2007), 1454–1508.
- [AS03] D. Aldous and J. M. Steele, *The objective method: probabilistic combinatorial optimization and local weak convergence*, in “Probability on discrete structures”, H. Kesten Ed., Springer Verlag, New York, 2003.
- [BG07] A. Bandyopadhyay, D. Gamarnik, *Counting without sampling. New algorithms for enumeration problems using statistical physics*, Proceedings of the 17th ACM-SIAM Symposium on Discrete Algorithm, Miami, Flo., 2007.
- [Bo80] B. Bollobás, *A probabilistic proof of an asymptotic formula for the number of labeled regular graphs*, *Eur. J. Combinatorics*, **1** (1980), 311–316.
- [DGM09] A. Dembo, A. Gerschenfeld and A. Montanari, *Spin glasses on locally tree-like graphs*, in preparation, 2009.
- [DZ98] A. Dembo and O. Zeitouni, “Large deviations techniques and applications: Second edition”, Springer Verlag, New York, 1998.
- [DG08] L. De Sanctis, F. Guerra, *Mean field dilute ferromagnet I. High temperature and zero temperature behavior*, *J. Stat. Phys.* **132** (2008), 759–785.
- [DGM04] S. N. Dorogovtsev, A. V. Goltsev and J. F. F. Mendes, *Ising model on networks with an arbitrary distribution of connections* *Phys. Rev. E* **66** (2002), 016104.
- [EN79] R. S. Ellis and C. M. Newman, *The statistics of Curie-Weiss models*, *J. Stat. Phys.* **19** (1978), 149–161.
- [EK+00] W. Evans, C. Kenyon, Y. Peres and L. J. Schulman *Broadcasting on trees and the Ising model*, *Ann. Appl. Prob.* **10** (2000), 410–433.
- [GM07] A. Gerschenfeld and A. Montanari, *Reconstruction for models on random graphs*, 48nd Annual Symposium on Foundations of Computer Science, Providence, RI, 2007.
- [GHS70] R. B. Griffiths, C. A. Hurst and S. Sherman, *Concavity of magnetization of an Ising ferromagnet in a positive external field*, *J. Math Phys.* **11** (1970), 790–795.
- [JLR00] S. Janson, T. Luczak and A. Ruciński, “Random graphs,” John Wiley and Sons, 2000.
- [JS90] M. Jerrum, A. Sinclair, *Polynomial-time approximation algorithms for the Ising model*, Automata, Languages and Programming, LNCS-443 (1990), 462–475.
- [JP98] D. A. Johnston and P. Plechác, *Equivalence of ferromagnetic spin models on trees and random graphs*, *J. Phys. A*, **31** (1998), 475–482.
- [JS99] J. Jonasson and J. E. Steif, *Amenability and phase transition in the Ising model*, *J. Theor. Prob.*, **12** (1999), 549–559.
- [KM+07] F. Krzakala, A. Montanari, F. Ricci-Tersenghi, G. Semerjian and L. Zdeborová, *Gibbs states and the set of solutions of random constraint satisfaction problems*, *Proc. Natl. Acad. Sci. (USA)*, **104** (2007), 10318–10323.

- [LV+02] M. Leone, A. Vázquez, A. Vespignani and R. Zecchina, *Ferromagnetic ordering in graphs with arbitrary degree distribution*, Eur. Phys. J. B **28** (2002), 191–197.
- [Lig85] T. M. Liggett “Interacting particle systems”, Springer Verlag, New York, 1985.
- [Lyo00] R. Lyons, *Phase transitions on nonamenable graphs*, J. Math. Phys. **41** (2000), 1099.
- [MM09] M. Mézard and A. Montanari, “Information, physics, and computation”, Oxford University Press, Oxford, 2009.
- [MK05] J. M. Mooij and H. J. Kappen, *On the properties of the Bethe approximation and loopy belief propagation on binary networks*, J. Stat. Mech. (2005), P11012.
- [SS09] J. Salez and D. Shah, *Belief propagation: an asymptotically optimal algorithm for the random assignment problem*, arXiv:0902.0585 (2009)
- [SWW07] E. Sudderth, M. Wainwright, and A. Willsky *Loop series and Bethe variational bounds in attractive graphical models*, to appear at Neural Information Processing Systems, 2007.
- [Sim80] B. Simon, *Correlation inequalities and the decay of correlations in ferromagnets*, Commun. Math Phys. **77** (1980), 111–126.
- [TJ02] S. Tatikonda and M. I. Jordan, *Loopy belief propagation and Gibbs measures*, Uncertainty in Artificial Intelligence, 2002.