

## ON OPTIMAL ARBITRAGE\*

BY DANIEL FERNHOLZ AND IOANNIS KARATZAS

*University of Texas at Austin and INTECH*

In a Markovian model for a financial market, we characterize the best arbitrage with respect to the market portfolio that can be achieved using non-anticipative investment strategies, in terms of the smallest positive solution to a parabolic partial differential inequality; this is determined entirely on the basis of the covariance structure of the model. The solution is intimately related to properties of strict local martingales, and is used to generate the investment strategy which realizes the best possible arbitrage. Some extensions to non-Markovian situations are also presented.

**1. Introduction.** In a Markovian model for an equity market with mean rates of return  $b_i(\mathfrak{X}(t))$  and covariance rates  $a_{ij}(\mathfrak{X}(t))$ ,  $1 \leq i, j \leq n$  for its asset capitalizations  $\mathfrak{X}(t) = (X_1(t), \dots, X_n(t))' \in (0, \infty)^n$  at time  $t$ , what is the highest return on investment (as in (6.3) below) that can be achieved relative to the market on a given time-horizon  $[0, T]$ , using non-anticipative investment strategies? What are the weights assigned to the different assets by such an investment strategy that accomplishes this?

Answers: under suitable conditions,  $1/U(T, \mathfrak{X}(0))$  and

$$X_i(t) D_i \log U(T-t, \mathfrak{X}(t)) + \frac{X_i(t)}{X_1(t) + \dots + X_n(t)}, \quad i = 1, \dots, n, \quad t \in [0, T]$$

respectively. Here  $U : [0, \infty) \times (0, \infty)^n \rightarrow (0, 1]$  is the smallest non-negative solution of the linear parabolic partial differential inequality

$$(1.1) \quad \frac{\partial U}{\partial \tau}(\tau, \mathbf{x}) \geq \widehat{\mathcal{L}}U(\tau, \mathbf{x}), \quad (\tau, \mathbf{x}) \in (0, \infty) \times (0, \infty)^n$$

---

\*The second author is on leave from the Department of Mathematics at Columbia University, where his research was supported by NSF Grant DMS-06-01774. We wish to thank G. Žitković, N. Sesum, E.R. Fernholz, A. Banner, V. Papathanakos, T. Ichiba and most notably J. Ruf for several helpful discussions. We are grateful to M. Sirbu, M. Soner, F. Delbaen, W. Schachermayer, J. Hugonnier, J. Teichmann and C. Kardaras for their comments. We are also deeply indebted to the referees and Associate Editor for their meticulous readings, and for suggestions that improved this paper greatly.

*AMS 2000 subject classifications:* Primary 60H10, 91B28; secondary 60G44, 35B50.

*Keywords and phrases:* portfolios, arbitrage, parabolic operators, maximum principle, strict local martingales, exit measures for supermartingales, diffusions, Fichera drift.

subject to the initial condition  $U(0, \cdot) \equiv 1$ , for the linear operator

$$(1.2) \quad \widehat{\mathcal{L}}f := \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n x_i x_j a_{ij}(\mathbf{x}) D_{ij}^2 f + \sum_{i=1}^n x_i \left( \sum_{j=1}^n \frac{x_j a_{ij}(\mathbf{x})}{x_1 + \cdots + x_n} \right) D_i f$$

with  $D_i = \partial / \partial x_i$ ,  $D = (\partial / \partial x_1, \dots, \partial / \partial x_n)'$  and  $D_{ij}^2 = \partial^2 / \partial x_i \partial x_j$ . Furthermore,  $U(T, \mathfrak{X}(0))$  is the probability that the  $([0, \infty)^n \setminus \{\mathbf{0}\})$ -valued diffusion process  $\mathfrak{Y}(\cdot) = (Y_1(\cdot), \dots, Y_n(\cdot))'$  with infinitesimal generator  $\widehat{\mathcal{L}}$  as above and  $\mathfrak{Y}(0) = \mathfrak{X}(0) \in (0, \infty)^n$ , does not hit the boundary of the orthant  $[0, \infty)^n$  by time  $t = T$ . We note that the answers involve only the covariance structure of the market, not the actual rates of return; the only rôle these latter play is to ensure that the diffusion  $\mathfrak{X}(\cdot)$  lives in  $(0, \infty)^n$ .

Arbitrage relative to the market exists on  $[0, T]$ , iff  $U(T, \mathfrak{X}(0)) < 1$ ; this is deeply related to the importance of *strict local martingales* in the present context, and amounts to failure of uniqueness for the Cauchy problem

$$\frac{\partial U}{\partial \tau}(\tau, \mathbf{x}) = \widehat{\mathcal{L}}U(\tau, \mathbf{x}), \quad (\tau, \mathbf{x}) \in (0, \infty) \times (0, \infty)^n \quad \text{and} \quad U(0, \cdot) \equiv 1.$$

Sufficient conditions for such failure of uniqueness are provided.

Consider an “auxiliary market”, with capitalizations  $\mathfrak{Y}(\cdot) = (Y_1(\cdot), \dots, Y_n(\cdot))'$  as above. The probabilistic significance of the change of drift inherent in the definition of the operator  $\widehat{\mathcal{L}}$ , from  $b_i(\mathbf{x})$  for  $\mathfrak{X}(\cdot)$  to  $\sum_{j=1}^n (x_j a_{ij}(\mathbf{x})) / (x_1 + \cdots + x_n)$  for  $\mathfrak{Y}(\cdot)$ , is that it corresponds to a change of probability measure which makes the weights  $\nu_i(\cdot) := Y_i(\cdot) / (Y_1(\cdot) + \cdots + Y_n(\cdot))$ ,  $i = 1, \dots, n$  of the auxiliary market portfolio martingales. Its financial significance is that it bestows to the auxiliary market portfolio  $\nu(\cdot) = (\nu_1(\cdot), \dots, \nu_n(\cdot))'$  the so-called *numéraire property*: any strategy’s relative performance in the market with capitalizations  $\mathfrak{Y}(\cdot)$  is a supermartingale, so this market cannot be outperformed. This change need not come from a Girsanov-type (absolutely continuous) transformation; rather it corresponds to, and represents, the *exit measure* of Föllmer (1972) for an appropriate supermartingale.

Sections 2 and 3 set up the model, whereas section 4 introduces the notion and offers examples of relative arbitrage; section 5 makes the connection with strict local martingales. Section 6 formulates the problem and section 7 offers some preliminary results, actually in some modest generality (including non-Markovian cases). Section 8 sets up the Markovian model; the results are presented in earnest in sections 9-11, section 12 discusses a couple of examples in detail, and a few open questions are raised in section 13.

*Related Literature:* The questions raised in this study are related to the work of Delbaen & Schachermayer (1995.b). They bear an even closer connection with issues raised in the Finance literature under the general rubric

of “bubbles” (see Definition 5 and Theorem 1 in Ruf (2009) for the precise connection). The literature on this topic is large, so let us mention the papers by Loewenstein & Willard (2000), Pal & Protter (2007) and, most significantly, Heston et al. (2007), as the closest in spirit to our approach here. We note the recent preprint by Hugonnier (2007), which demonstrates that arbitrage opportunities can arise in equilibrium models; this preprint, and Heston et al. (2007), can be consulted for an up-to-date survey of the literature on this subject and for some explicit computations of trading strategies that lead to arbitrage. The need to consider state-price-density processes that are only local (as opposed to true) martingales has also been noticed in the context of “stochastic volatility” models (e.g. Sin (1998), Wong & Heyde (2006)) and of pricing with long maturities (e.g. Hulley & Platen (2008)).

**2. The Model.** We consider a model consisting of a money-market  $dB(t) = B(t)r(t)dt$ ,  $B(0) = 1$  and of  $n$  stocks with capitalizations

$$(2.1) \quad dX_i(t) = X_i(t) \left( \beta_i(t)dt + \sum_{k=1}^K \sigma_{ik}(t) dW_k(t) \right), \quad X_i(0) = x_i > 0$$

for  $i = 1, \dots, n$ . These are defined on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  and are driven by the Brownian motion  $W(\cdot) = (W_1(\cdot), \dots, W_K(\cdot))'$  whose  $K \geq n$  independent components are the model’s “factors”.

We shall assume throughout that the interest rate process of the money-market is  $r(\cdot) \equiv 0$ , identically equal to zero; and that the vector-valued process  $\mathfrak{X}(\cdot) = (X_1(\cdot), \dots, X_n(\cdot))'$  of capitalizations, the vector-valued process  $\beta(\cdot) = (\beta_1(\cdot), \dots, \beta_n(\cdot))'$  of *mean rates of return* for the various stocks, and the  $(n \times K)$ -matrix-valued process  $\sigma(\cdot) = (\sigma_{ik}(\cdot))_{1 \leq i \leq n, 1 \leq k \leq K}$  of *volatilities*, are all progressively measurable with respect to a right-continuous filtration  $\mathbb{F} = \{\mathcal{F}(t)\}_{0 \leq t < \infty}$  which represents the “flow of information” in the market with  $\mathcal{F}(0) = \{\emptyset, \Omega\}$ , mod.  $\mathbb{P}$ . Let  $\alpha(\cdot) := \sigma(\cdot)\sigma'(\cdot)$  be the *covariance process* of the stocks in the market, and impose for  $\mathbb{P}$ -a.e.  $\omega \in \Omega$  the condition

$$(2.2) \quad \sum_{i=1}^n \int_0^T \left( |\beta_i(t, \omega)| + \alpha_{ii}(t, \omega) \right) dt < \infty, \quad \forall T \in (0, \infty).$$

Under this condition the processes  $X_1(\cdot), \dots, X_n(\cdot)$  can be expressed as  $X_i(\cdot) = x_i \exp \left\{ \int_0^\cdot (\beta_i(t) - \frac{1}{2} \alpha_{ii}(t)) dt + \sum_{k=1}^K \int_0^\cdot \sigma_{ik}(t) dW_k(t) \right\} > 0$ .

In this setting, the Brownian motion  $W(\cdot)$  need not be adapted to the “observations” filtration  $\mathbb{F}$ . It *is* adapted, though, to the  $\mathbb{P}$ -augmentation  $\mathbb{G} = \{\mathcal{G}(t)\}_{0 \leq t < \infty}$  of the filtration  $\mathbb{F}$ , provided that  $K = n$  and that the matrix-valued process  $\sigma(\cdot)$  is invertible – as in Assumption B below.

**3. Strategies and Portfolios.** Consider now a *small investor* who decides, at each time  $t$ , which proportion  $\pi_i(t)$  of current wealth  $V(t)$  to invest in the  $i^{\text{th}}$  stock,  $i = 1, \dots, n$ ; the proportion  $1 - \sum_{i=1}^n \pi_i(t) =: \pi_0(t)$  gets invested in the money market. Thus, the wealth  $V(\cdot) \equiv V^{v,\pi}(\cdot)$  for an initial capital  $v \in (0, \infty)$  and an investment strategy  $\pi(\cdot) = (\pi_1(\cdot), \dots, \pi_n(\cdot))'$  satisfies the initial condition  $V(0) = v$  and

$$(3.1) \quad \frac{dV(t)}{V(t)} = \sum_{i=1}^n \pi_i(t) \frac{dX_i(t)}{X_i(t)} + \pi_0(t) \frac{dB(t)}{B(t)} = \pi'(t) [\beta(t)dt + \sigma(t)dW(t)].$$

We shall call *investment strategy* a  $\mathbb{G}$ -progressively measurable process  $\pi : [0, \infty) \times \Omega \rightarrow \mathbb{R}^n$  which satisfies for  $\mathbb{P}$ -a.e.  $\omega \in \Omega$  the analogue  $\int_0^T (|\pi'(t, \omega) \beta(t, \omega)| + \pi'(t, \omega) \alpha(t, \omega) \pi(t, \omega)) dt < \infty$ ,  $\forall T \in (0, \infty)$  of (2.2). The collection of investment strategies will be denoted by  $\mathcal{H}$ .

A strategy  $\pi(\cdot) \in \mathcal{H}$  with  $\sum_{i=1}^n \pi_i(t, \omega) = 1$  for all  $(t, \omega) \in [0, \infty) \times \Omega$  will be called *portfolio*. A portfolio never invests in the money market, and never borrows from it. We shall say that a process  $\pi(\cdot)$  is *bounded*, if for it there exists a real constant  $C_\pi > 0$  such that  $\|\pi(t, \omega)\| \leq C_\pi$  holds for all  $(t, \omega) \in [0, \infty) \times \Omega$ . We shall call *long-only portfolio* one that satisfies  $\pi_1(t, \omega) \geq 0, \dots, \pi_n(t, \omega) \geq 0$ ,  $\forall (t, \omega) \in [0, \infty) \times \Omega$ , i.e., never sells any stock short. Clearly, a long-only portfolio is also bounded.

Corresponding to an investment strategy  $\pi(\cdot)$  and initial capital  $v > 0$ , the associated wealth process, i.e., the solution of the equation (3.1), is

$$V^{v,\pi}(\cdot) = v \exp \left\{ \int_0^\cdot \pi'(t) \left( \beta(t) - \frac{\alpha(t)}{2} \pi(t) \right) dt + \int_0^\cdot \pi'(t) \sigma(t) dW(t) \right\} > 0.$$

The strategy  $\varrho(\cdot) \equiv 0$  invests only in the money market at all times; it results in  $V^{v,\varrho}(\cdot) \equiv v$ , i.e., in hoarding the initial wealth under the mattress.

**3.1. The Market Portfolio.** An important long-only portfolio is the market portfolio; this invests in all stocks in proportion to their relative weights

$$(3.2) \quad \mu_i(t) := \frac{X_i(t)}{X(t)}, \quad i = 1, \dots, n, \quad \text{where } X(t) := X_1(t) + \dots + X_n(t).$$

Clearly  $V^{v,\mu}(\cdot) = vX(\cdot)/X(0)$ , and the resulting vector process  $\mu(\cdot) = (\mu_1(\cdot), \dots, \mu_n(\cdot))'$  of market weights takes values in the positive simplex  $\Delta_+^n := \{(m_1, \dots, m_n)' \in (0, 1)^n \mid \sum_{i=1}^n m_i = 1\}$  of  $\mathbb{R}^n$ . An application of Itô's rule gives, after some computation, the dynamics of this process as

$$(3.3) \quad d\mu_i(t) = \mu_i(t) \left[ \gamma_i^\mu(t) dt + \sum_{k=1}^K \tau_{ik}^\mu(t) dW_k(t) \right], \quad i = 1, \dots, n.$$

Here  $\tau^\mu(t)$  is the matrix with entries  $\tau_{ik}^\mu(t) := \sigma_{ik}(t) - \sum_{j=1}^n \mu_j(t) \sigma_{jk}(t)$ , the vector  $\gamma^\mu(t) := (\gamma_1^\mu(t), \dots, \gamma_n^\mu(t))'$  has components

$$(3.4) \quad \gamma_i^\mu(t) := (\mathbf{e}_i - \mu(t))'(\beta(t) - \alpha(t)\mu(t)),$$

and  $\mathbf{e}_i$  the  $i^{\text{th}}$  unit vector in  $\mathbb{R}^n$ .

**4. Relative Arbitrage.** The following notion was introduced in Fernholz (2002): given a real number  $T > 0$  and any two investment strategies  $\pi(\cdot)$  and  $\rho(\cdot)$ , we call  $\pi(\cdot)$  an *arbitrage relative to*  $\rho(\cdot)$  over  $[0, T]$ , if

$$(4.1) \quad \mathbb{P}(V^{1,\pi}(T) \geq V^{1,\rho}(T)) = 1 \quad \text{and} \quad \mathbb{P}(V^{1,\pi}(T) > V^{1,\rho}(T)) > 0.$$

We call such relative arbitrage *strong*, if  $\mathbb{P}(V^{1,\pi}(T) > V^{1,\rho}(T)) = 1$ .

Arbitrage (resp., strong arbitrage) relative to  $\varrho(\cdot) \equiv 0$  that invests only in the money market, is called just that, without the qualifier ‘‘relative’’.

4.1. *Examples of Arbitrage Relative to the Market.* Here are some examples taken from the survey Fernholz & Karatzas (2009), especially Sections 7 and 8, Remark 11.4, Examples 11.1 and 11.2. Suppose first that

$$(4.2) \quad \sum_{i=1}^n \mu_i(t) \alpha_{ii}(t) - \sum_{i=1}^n \sum_{j=1}^n \mu_i(t) \alpha_{ij}(t) \mu_j(t) \geq h, \quad \forall 0 \leq t < \infty$$

holds almost surely for some constant  $h > 0$ . Then the long-only portfolio  $\pi_i(t) = \mu_i(t)(c - \log \mu_i(t))/J(t)$ ,  $i = 1, \dots, n$ ,  $J(t) := \sum_{j=1}^n \mu_j(t)(c - \log \mu_j(t))$  is, for sufficiently large  $c > 0$ , a strong arbitrage relative to the market portfolio  $\mu(\cdot)$  over any time-horizon  $[0, T]$  with  $T > (2 \log n)/h$ .

Another condition guaranteeing the existence of strong arbitrage relative to the market, is that there exist a real constant  $h > 0$  with

$$(4.3) \quad \sqrt{\mu_1(t) \cdots \mu_n(t)} \left[ \sum_{i=1}^n \alpha_{ii}(t) - \frac{1}{n} \sum_{i=1}^n \sum_{j=1}^n \alpha_{ij}(t) \right] \geq h, \quad \forall 0 \leq t < \infty$$

a.s. Then for  $c > 0$  sufficiently large, the long-only portfolio  $\pi_i(t) = \lambda(t)(1/n) + (1 - \lambda(t))\mu_i(t)$ ,  $1 \leq i \leq n$ ,  $1/\lambda(t) := 1 + ((\mu_1(t) \cdots \mu_n(t))^{1/n}/c)$ , is strong arbitrage relative to the market over any  $[0, T]$  with  $T > (2n^{1-(1/n)})/h$ .

**Remark 1:** Suppose that all the eigenvalues of the covariance matrix-valued process  $\alpha(\cdot)$  are bounded away from both zero and infinity, uniformly on  $[0, \infty) \times \Omega$ ; and that (4.2) holds. Then, for any given constant  $p \in (0, 1)$ , the long-only portfolio  $\mu_i^{(p)}(t) = (\mu_i(t))^p (\sum_{j=1}^n (\mu_j(t))^p)^{-1}$ ,  $i = 1, \dots, n$  leads again to strong arbitrage relative to the market portfolio over sufficiently long time-horizons. It is also of great interest that appropriate modifications of the portfolio  $\mu^{(p)}(\cdot)$  yield such arbitrage over any time-horizon  $[0, T]$ .

**5. Market Price of Risk and Strict Local Martingales.** We shall assume from now on that there exists a *market price of risk*  $\vartheta : [0, \infty) \times \Omega \rightarrow \mathbb{R}^K$ , an  $\mathbb{F}$ -progressively measurable process that satisfies

$$(5.1) \quad \begin{aligned} & \sigma(t, \omega)\vartheta(t, \omega) = \beta(t, \omega), \quad \forall (t, \omega) \in [0, \infty) \times \Omega; \quad \text{and} \\ & \mathbb{P} \left( \int_0^T \|\vartheta(t, \omega)\|^2 dt < \infty, \quad \forall T \in (0, \infty) \right) = 1. \end{aligned}$$

The existence of a market-price-of-risk process  $\vartheta(\cdot)$  allows us to introduce an associated exponential local martingale

$$(5.2) \quad Z(t) := \exp \left\{ - \int_0^t \vartheta'(s) dW(s) - \frac{1}{2} \int_0^t \|\vartheta(s)\|^2 ds \right\}, \quad 0 \leq t < \infty.$$

This process is also a supermartingale; it is a martingale, if and only if  $\mathbb{E}(Z(T)) = 1$  holds for all  $T \in (0, \infty)$ . For the purposes of this work it is important to allow such exponential processes to be strict local martingales; that is, not to exclude the possibility  $\mathbb{E}(Z(T)) < 1$  for some  $T \in (0, \infty)$ .

From (5.2) and (3.1), now written in the form

$$(5.3) \quad dV^{v, \pi}(t) = V^{v, \pi}(t) \pi'(t) \sigma(t) d\widehat{W}(t), \quad \widehat{W}(t) := W(t) + \int_0^t \vartheta(s) ds$$

on the strength of (5.1), the product rule of Itô's calculus shows that

$$(5.4) \quad Z(\cdot) V^{v, \pi}(\cdot) = v + \int_0^\cdot Z(t) V^{v, \pi}(t) (\sigma'(t) \pi(t) - \vartheta(t))' dW(t)$$

is a positive local martingale and a supermartingale, for every  $\pi(\cdot) \in \mathcal{H}$ .

If  $\alpha(\cdot)$  is invertible, we can take  $\vartheta(\cdot) = \sigma'(\cdot) \alpha^{-1}(\cdot) \beta(\cdot)$  as market price of risk in (5.1). If  $\beta(\cdot) = \alpha(\cdot) \mu(\cdot)$  holds we can select  $\vartheta(\cdot) = \sigma'(\cdot) \mu(\cdot)$  and get  $Z(\cdot) \equiv v/V^{v, \mu}(\cdot) \equiv X(0)/X(\cdot)$  from (5.4); there is then no arbitrage relative to the market because  $V^{v, \pi}(\cdot)/V^{v, \mu}(\cdot)$  is a *supermartingale for all*  $\pi(\cdot) \in \mathcal{H}$ , thus  $\mathbb{E}[V^{1, \pi}(T)/V^{1, \mu}(T)] \leq 1$ , a conclusion at odds with (4.1).

**5.1. Strict Local Martingales.** Suppose the covariance process  $\alpha(\cdot)$  is bounded, and (4.1) holds for two bounded portfolios  $\pi(\cdot)$  and  $\rho(\cdot)$ . Then, for *any* market-price-of-risk process  $\vartheta(\cdot)$  as in (5.1), the positive local martingales  $Z(\cdot)$  and  $Z(\cdot) V^{v, \rho}(\cdot)$  of (5.2), (5.4) are strict:  $\mathbb{E}[Z(T) V^{v, \rho}(T)] < v$ ,  $\mathbb{E}(Z(T)) < 1$  (Fernholz & Karatzas (2009), section 6).

In particular, if the matrix  $\alpha(\cdot)$  is bounded and (4.1) holds for some bounded portfolio  $\pi(\cdot)$  and for the market portfolio  $\rho(\cdot) \equiv \mu(\cdot)$  (these assumptions are satisfied, e.g., under the conditions in Remark 1), then

$$(5.5) \quad \begin{aligned} & \mathbb{E}(Z(T)) < 1, \quad \mathbb{E}[Z(T) X(T)] < X(0), \\ & \mathbb{E}[Z(T) X_i(T)] < X_i(0), \quad i = 1, \dots, n. \end{aligned}$$

**6. Optimal Arbitrage Relative to the Market.** The possibility of strong arbitrage relative to the market, defined and exemplified in section 4, raises an obvious question: *What is the best possible arbitrage of this kind?*

One way to cast this question is as follows: On a given time-horizon  $[0, T]$ , what is the *smallest* relative amount

$$(6.1) \quad u(T) := \inf \{w > 0 \mid \exists \pi(\cdot) \in \mathcal{H} \text{ s.t. } V^{wX(0), \pi}(T) \geq X(T), \text{ a.s.}\}$$

of initial capital, starting with which one can match or exceed at time  $t = T$  the market capitalization  $X(T)$ ? Clearly,  $0 < u(T) \leq 1$ ; and for  $0 < w < u(T)$ , no strategy starting with initial capital  $wX(0)$  can outperform the market almost surely, over the horizon  $[0, T]$ . That is, for every  $\pi(\cdot) \in \mathcal{H}$  and  $0 < w < u(T)$ , we have  $\mathbb{P}[V^{wX(0), \pi}(T) \geq X(T)] < 1$ .

We shall impose from now on the following structural assumption on the filtration  $\mathbb{F} = \{\mathcal{F}(t)\}_{0 \leq t < \infty}$ , the “flow of information” in the market.

**Assumption A:** *Every local martingale of the filtration  $\mathbb{F}$  can be represented as a stochastic integral, with respect to the driving Brownian motion  $W(\cdot)$  in (2.1), of some  $\mathbb{G}$ -progressively measurable integrand.*

**Assumption B:** *We have  $K = n$  and  $\sigma(t)$  is invertible,  $\forall t \in [0, T]$ .*

Under these two assumptions, general results about hedging in so-called complete markets (e.g., Karatzas & Shreve (1998); section 10 in Fernholz & Karatzas (2009); or Ruf (2009)) based on martingale representation results, show that the quantity of (6.1) is given as

$$(6.2) \quad u(T) = \mathbb{E}[Z(T)X(T)]/X(0); \quad \text{that } V^{u(T)X(0), \hat{\pi}}(T) = X(T)$$

holds a.s. for some  $\hat{\pi}(\cdot) \in \mathcal{H}$ ; and that  $1/u(T)$  gives the highest return

$$(6.3) \quad \sup \{q \geq 1 \mid \exists \pi(\cdot) \in \mathcal{H} \text{ s.t. } V^{1, \pi}(T) \geq qV^{1, \mu}(T), \text{ a.s.}\}$$

on investment, that one can achieve relative to the market over  $[0, T]$ . Arbitrage relative to the market is possible on  $[0, T]$ , if and only if  $u(T) < 1$ .

The result in (6.2) provides no information about the strategy  $\hat{\pi}(\cdot)$  that implements this ‘best possible’ arbitrage, apart from ascertaining its existence. In section 8 we shall specialize the model of (2.1) to a Markovian context and describe  $\hat{\pi}(\cdot)$  in terms of partial differential equations (section 11). We shall also characterize the quantity  $u(T)$  in terms of the smallest solution to a parabolic partial differential inequality, and as the probability of non-absorption by time  $T$  for a suitable diffusion (Theorems 1, 2).

Assumption A holds when  $\mathbb{F}$  is (the augmentation of)  $\mathbb{F}^W$ , the filtration generated by the Brownian motion  $W(\cdot)$ ; as well as when Assumption B holds, the  $\beta_i(\cdot)$ ,  $\sigma_{i\nu}(\cdot)$  are all progressively measurable with respect to  $\mathbb{F}^{\mathfrak{X}} = \{\mathcal{F}^{\mathfrak{X}}(t)\}_{0 \leq t < \infty}$ ,  $\mathcal{F}^{\mathfrak{X}}(t) := \sigma(\mathfrak{X}(s), 0 \leq s \leq t)$ , and  $\mathbb{F} \equiv \mathbb{F}_+^{\mathfrak{X}}$  (Jacod (1977)).

6.1. *Generalized Likelihood Ratios.* The positive local martingale  $Z(\cdot)X(\cdot)$ , whose expectation appears in (6.2), can be expressed as

$$(6.4) \quad Z(t)X(t) = X(0) \cdot \exp \left\{ - \int_0^t (\tilde{\vartheta}(s))' dW(s) - \frac{1}{2} \int_0^t \|\tilde{\vartheta}(s)\|^2 ds \right\}$$

for  $0 \leq t \leq T$ . Here we have solved equation (5.4) for  $\pi(\cdot) \equiv \mu(\cdot)$  and set

$$(6.5) \quad \tilde{\vartheta}(\cdot) := \vartheta(\cdot) - \sigma'(\cdot)\mu(\cdot), \quad \tilde{W}(\cdot) := W(\cdot) + \int_0^\cdot \tilde{\vartheta}(t) dt,$$

whence  $\sigma(\cdot)\tilde{\vartheta}(\cdot) = \beta(\cdot) - \alpha(\cdot)\mu(\cdot)$  from (5.1); we thus re-cast (2.1) as

$$(6.6) \quad dX_i(t) = X_i(t) \left[ \frac{\sum_{j=1}^n \alpha_{ij}(t)X_j(t)}{X_1(t) + \cdots + X_n(t)} dt + \sum_{k=1}^n \sigma_{ik}(t) d\tilde{W}_k(t) \right].$$

On the other hand, we note from (6.4), (6.5) that the reciprocal of the exponential local martingale  $Z(\cdot)X(\cdot)/X(0)$  can be expressed as

$$(6.7) \quad \Lambda(\cdot) := \frac{X(0)}{Z(\cdot)X(\cdot)} = \exp \left\{ \int_0^\cdot (\tilde{\vartheta}(t))' d\tilde{W}(t) - \frac{1}{2} \int_0^\cdot \|\tilde{\vartheta}(t)\|^2 dt \right\};$$

similarly, the reciprocal of the local martingale  $Z(\cdot)X_i(\cdot)/X_i(0)$  is

$$(6.8) \quad \Lambda_i(\cdot) := \frac{X_i(0)}{Z(\cdot)X_i(\cdot)} = \exp \left\{ \int_0^\cdot (\tilde{\vartheta}^{(i)}(t))' d\tilde{W}^{(i)}(t) - \frac{1}{2} \int_0^\cdot \|\tilde{\vartheta}^{(i)}(t)\|^2 dt \right\}$$

where  $\tilde{\vartheta}^{(i)}(\cdot) := \vartheta(\cdot) - \sigma'(\cdot)\mathbf{e}_i$  and  $\tilde{W}^{(i)}(\cdot) := W(\cdot) + \int_0^\cdot \tilde{\vartheta}^{(i)}(t) dt$ .

Comparing (6.7) and (6.8), we observe that  $\mu_i(0)\Lambda(\cdot) = \mu_i(\cdot)\Lambda_i(\cdot)$  and cast the dynamics of (3.3)-(3.4) for the market portfolio  $\mu(\cdot)$  as

$$(6.9) \quad d\mu_i(t) = \mu_i(t) (\mathbf{e}_i - \mu(t))' \sigma(t) d\tilde{W}(t), \quad i = 1, \dots, n.$$

If  $\mathbf{u}(T) = 1$ , i.e.,  $Z(\cdot)X(\cdot)$  is a martingale on  $[0, T]$ , no arbitrage relative to the market is possible on this time-horizon; the ‘reference’ measure

$$(6.10) \quad \tilde{\mathbb{P}}_T(A) := \mathbb{E}[Z(T)X(T)1_A] / X(0), \quad A \in \mathcal{F}(T)$$

is a probability, i.e.,  $\mathbf{u}(T) = \tilde{\mathbb{P}}_T(\Omega) = 1$ ; and under  $\tilde{\mathbb{P}}_T$ , the process  $\tilde{W}(t)$ ,  $0 \leq t \leq T$  in (6.5) is a Brownian motion by the Girsanov theorem, so from (6.9) the market weights  $\mu_1(t), \dots, \mu_n(t)$ ,  $0 \leq t \leq T$  are martingales.

We shall characterize next  $\mathbf{u}(T)$  in terms of the *Föllmer exit measure*, of a “generalized martingale measure”, and of a measure  $\mathbb{Q}$  with respect to which  $\mathbb{P}$  is locally absolutely continuous (eq’s (7.3), (7.6)) and which plays, to a considerable extent, the rôle of  $\tilde{\mathbb{P}}_T$  when  $Z(\cdot)X(\cdot)$  fails to be a  $\mathbb{P}$ -martingale. The processes of (6.4)-(6.8) are important in this effort.

**7. Exit Measure of a Positive Supermartingale.** We shall assume in this section that the process  $Z(\cdot)$  of (5.2) is adapted to  $\mathbb{F} = \{\mathcal{F}(t)\}_{0 \leq t < \infty}$ , and that this filtration is, in turn, the right-continuous version  $\mathcal{F}(t) = \bigcap_{\varepsilon > 0} \mathcal{F}^o(t + \varepsilon)$  of a *standard system*  $\mathbb{F}^o = \{\mathcal{F}^o(t)\}_{0 \leq t < \infty}$ : To wit, each  $(\Omega, \mathcal{F}^o(t))$  is isomorphic to the Borel  $\sigma$ -algebra of some Polish space; and for any decreasing sequence  $\{A_j\}_{j \in \mathbb{N}}$  such that  $A_j$  is an atom of  $\mathcal{F}^o(t_j)$ , for some increasing sequence  $\{t_j\}_{j \in \mathbb{N}} \subset [0, \infty)$ , we have  $\bigcap_{j \in \mathbb{N}} A_j \neq \emptyset$ .

The canonical example is the space  $\Omega$  of right-continuous paths  $\omega : [0, \infty) \rightarrow \mathbb{R}^n \cup \{\Delta\}$ , where  $\Delta$  is an additional ‘‘absorbing point’’; paths stay at  $\Delta$  once they get there, i.e., after  $\mathcal{T}(\omega) = \inf\{t \geq 0 \mid \omega(t) = \Delta\}$ , and are continuous on  $(0, \mathcal{T}(\omega))$ . If  $\mathcal{F}^o(t) = \sigma(\omega(s), 0 \leq s \leq t)$ , then  $\mathbb{F}^o = \{\mathcal{F}^o(t)\}_{0 \leq t < \infty}$  is a standard system; see Föllmer (1972), Appendix.

Under these conditions, we can associate to the  $(\mathbb{P}, \mathbb{F})$ -local martingale  $Z(\cdot)X(\cdot)$  a positive measure  $\mathfrak{P}$  on the predictable  $\sigma$ -algebra of  $[0, \infty) \times \Omega$ ,

$$\mathfrak{P}((T, \infty] \times A) := \mathbb{E} [Z(T)X(T) 1_A] / X(0), \quad A \in \mathcal{F}(T), \quad T \in [0, \infty),$$

by invoking an extension result (Theorem V.4.1 in Parthasarathy (1967), whence the assumptions on the nature of the probability space).

This is the ‘‘exit measure’’ of the supermartingale  $Z(\cdot)X(\cdot)$ , introduced by Föllmer (1972, 1973); see also Delbaen & Schachermayer (1995.a), Föllmer & Gundel (2006). Föllmer (1972) obtained a characterization of the (process-theoretic) properties of supermartingales, such as  $Z(\cdot)X(\cdot)$  here, in terms of the properties of  $\mathfrak{P}$ . It follows from his work that  $Z(\cdot)X(\cdot)$  is a

- *martingale*, if and only if  $\mathfrak{P}$  is concentrated on  $\{\infty\} \times \Omega$ ;
- *potential* (i.e.,  $u(\infty) = 0$ ), if and only if  $\mathfrak{P}$  is concentrated on  $(0, \infty) \times \Omega$ .

**7.1. A Representation of the Föllmer Measure.** From Theorem 4 in Delbaen & Schachermayer (1995.a) and Theorem 1 and Lemma 4 of Pal & Protter (2007), the exponential process  $\Lambda(\cdot)$  of (6.7) is a continuous martingale under some probability measure  $\mathbb{Q}$  on the filtered space  $(\Omega, \mathcal{F})$ ,  $\mathbb{F}$  as above. The measure  $\mathbb{P}$  is locally absolutely continuous with respect to  $\mathbb{Q}$ , with  $d\mathbb{P} = \Lambda(T) d\mathbb{Q}$  on each  $\mathcal{F}(T)$ ; and the process  $\widetilde{W}(\cdot)$  of (6.5) is  $\mathbb{Q}$ -Brownian motion (cf. Ruf (2009), section 5). Thus, from (6.9) *the weights*  $\mu_1(\cdot), \dots, \mu_n(\cdot)$  *are martingales* and satisfy  $\sum_{i=1}^n \mu_i(\cdot) \equiv 1$  a.e., under  $\mathbb{Q}$ .

We consider the first time the process  $\Lambda(\cdot)$  hits the origin:

$$(7.1) \quad \mathcal{T} := \inf\{t \geq 0 \mid \Lambda(t) = 0\} = \inf\{t \geq 0 \mid Z(t)X(t) = \infty\}$$

(infinite, if the set is empty). We have  $\mathbb{P}(\mathcal{T} < \infty) = 0$ , but  $\mathbb{Q}(\mathcal{T} < \infty)$  can be positive, so  $\mathbb{Q}$  may not be absolutely continuous with respect to  $\mathbb{P}$ ; whereas,  $\mathbb{Q}$ -a.e. on  $\{\mathcal{T} < \infty\}$ , we have  $Z(\mathcal{T} + h)X(\mathcal{T} + h) = \infty$ ,  $\forall h \geq 0$  and  $\int_0^{\mathcal{T}} \|\widehat{v}(t)\|^2 dt = \infty$ . Intuitively, the rôle of the absorbing state  $\Delta$  is to

account for events that have zero  $\mathbb{P}$ -measure, but positive  $\mathbb{Q}$ -measure. We also introduce the first times the processes  $\mu_i(\cdot)$  and  $\Lambda_i(\cdot)$  hit the origin:

$$(7.2) \quad \mathcal{T}_i := \inf\{t \geq 0 \mid \mu_i(t) = 0\}, \quad \tilde{\mathcal{T}}_i := \inf\{t \geq 0 \mid \Lambda_i(t) = 0\}.$$

**Proposition 1:** *The quantity of (6.1) can be represented as*

$$(7.3) \quad u(T) = \mathfrak{P}((T, \infty] \times \Omega) = \mathbb{Q}(\mathcal{T} > T).$$

Furthermore, we also have the  $\mathbb{Q}$ -a.e. representations

$$(7.4) \quad \mathcal{T} = \min_{1 \leq i \leq n} \tilde{\mathcal{T}}_i; \quad \text{as well as } \mathcal{T} = \min_{1 \leq i \leq n} \mathcal{T}_i \text{ away from the event } E,$$

where  $E := \{\mathcal{T} < \infty\} \cap \{\mu_1(\mathcal{T}) \cdots \mu_n(\mathcal{T}) > 0\}$ . This event has  $\mathbb{Q}$ -measure equal to zero, if for some real constant  $C > 0$  we have

$$(7.5) \quad \|\vartheta(t, \omega)\|^2 \leq C(1 + \text{Tr}(\alpha(t, \omega))), \quad \forall (t, \omega) \in [0, \infty) \times \Omega.$$

*Proof:* We note  $\mathfrak{P}((T, \infty] \times A) = \mathbb{E}^{\mathbb{P}}(\Lambda^{-1}(T) 1_{\{A \cap \{\mathcal{T} > T\}\}}) = \mathbb{E}^{\mathbb{Q}}(\Lambda(T) \cdot \Lambda^{-1}(T) 1_{A \cap \{\mathcal{T} > T\}}) = \mathbb{Q}(A \cap \{\mathcal{T} > T\})$ ,  $\forall A \in \mathcal{F}(T)$ . With  $A = \Omega$ , we get (7.3). For  $A = \{\mu_1(\mathcal{T}) \cdots \mu_n(\mathcal{T}) = 0\}$ , this gives  $\mathbb{Q}(A \cap \{\mathcal{T} > T\}) = 0$ : all the weights  $\mu_1(\cdot), \dots, \mu_n(\cdot)$  (equivalently, all capitalizations  $X_1(\cdot), \dots, X_n(\cdot)$ ) are strictly positive on  $[0, \mathcal{T})$ ,  $\mathbb{Q}$ -a.e.

Recall  $\mu_i(0)\Lambda(\cdot) \equiv \mu_i(\cdot)\Lambda_i(\cdot)$ ,  $\forall i = 1, \dots, n$  from (6.7), (6.8); this gives  $1/\Lambda(\cdot) = \sum_{i=1}^n (\mu_i(0)/\Lambda_i(\cdot))$  on  $[0, \mathcal{T})$ , and the first equation in (7.4).

On  $\{\mathcal{T} < \infty\} \setminus E$ , for some  $j \in \{1, \dots, n\}$  we shall have  $\mu_j(\mathcal{T}) = 0$ , thus also  $\mathcal{T}_j = \mathcal{T}$  and the second equation in (7.4). On the other hand,  $\mathcal{T}_i = \infty$ ,  $\forall i = 1, \dots, n$ ,  $\mathbb{Q}$ -a.e. on  $\{\mathcal{T} = \infty\}$ , so this equation holds on  $\{\mathcal{T} = \infty\}$ .

Finally, from (6.6), (6.7):  $\int_0^{\mathcal{T}} \text{Tr}(\alpha(t, \omega)) dt < \infty$ ,  $\int_0^{\mathcal{T}} \|\tilde{\vartheta}(t, \omega)\|^2 dt = \infty$  for  $\mathbb{Q}$ -a.e.  $\omega \in E \subseteq \{\mathcal{T} < \infty\}$ . Then (7.5) implies  $\int_0^{\mathcal{T}} \|\vartheta(t, \omega)\|^2 dt < \infty$ , and  $\tilde{\vartheta}(\cdot) = \vartheta(\cdot) - \sigma'(\cdot)\mu(\cdot)$  gives  $\int_0^{\mathcal{T}} \|\tilde{\vartheta}(t, \omega)\|^2 dt < \infty$ , thus  $\mathbb{Q}(E) = 0$ .  $\square$

The equation (7.3) can be thought of as a ‘generalized Wald identity’; cf. Problem 3.5.7 in Karatzas & Shreve (1991). In subsection 9.3 we shall obtain a characterization of the type (7.3) in a Markovian context, in terms of properties of an auxiliary diffusion and with the help of an appropriate partial differential equation. This will enable us to describe the investment strategy that realizes the optimal arbitrage.

**7.2. A Generalized Martingale Measure.** In a similar vein, there exists on the filtered space  $(\Omega, \mathcal{F})$ ,  $\mathbb{F}$  a probability measure  $\hat{\mathbb{Q}}$  under which

$$L(t) := 1/Z(t) = \exp \left\{ \int_0^t \vartheta'(s) d\widehat{W}(s) - \frac{1}{2} \int_0^t \|\vartheta(s)\|^2 ds \right\}, \quad 0 \leq t < \infty$$

is a martingale, and  $d\mathbb{P} = L(T) d\widehat{\mathbb{Q}}$  on each  $\mathcal{F}(T)$ ; whereas  $\widehat{W}(\cdot)$  of (5.3) is  $\widehat{\mathbb{Q}}$ -Brownian motion. Under  $\widehat{\mathbb{Q}}$ , the processes  $X_i(\cdot)$ ,  $i = 1, \dots, n$  are non-negative local (and super-)martingales:  $dX_i(t) = X_i(t) \sum_{k=1}^K \sigma_{ik}(t) d\widehat{W}_k(t)$ . This justifies the appellation “generalized martingale measure” for  $\widehat{\mathbb{Q}}$ .

Defining  $\mathcal{S} := \inf\{t \geq 0 \mid L(t) = 0\}$ , we have  $\mathbb{P}(\mathcal{S} < \infty) = 0$  and  $Z(\cdot)$  is a strict  $\mathbb{P}$ -local martingale if and only if  $\widehat{\mathbb{Q}}(\mathcal{S} < \infty) > 0$  (a potential, if and only if  $\widehat{\mathbb{Q}}(\mathcal{S} < \infty) = 1$ ); and the expression of (6.1), (6.2) is

$$(7.6) \quad u(T) = \mathbb{E}^{\widehat{\mathbb{Q}}}[(X(T)/X(0)) 1_{\{\mathcal{S} > T\}}].$$

This last expression takes the form  $u(T) = 1 - \mathbb{E}^{\widehat{\mathbb{Q}}}[ (X(\mathcal{S})/X(0)) 1_{\{\mathcal{S} \leq T\}} ]$  when  $X(\cdot \wedge T)$  is a  $\widehat{\mathbb{Q}}$ -martingale; from (5.3), this will be the case under the Novikov condition  $\mathbb{E}^{\widehat{\mathbb{Q}}}[\exp\{\frac{1}{2} \int_0^T \mu'(t) \alpha(t) \mu(t) dt\}] < \infty$ . Moreover,  $u(T) = 1$  (no arbitrage relative to the market is possible on  $[0, T]$ ), if and only if:  $X(\cdot \wedge T)$  is a  $\widehat{\mathbb{Q}}$ -martingale, and  $X(\mathcal{S}) 1_{\{\mathcal{S} \leq T\}} = 0$  holds  $\widehat{\mathbb{Q}}$ -a.e.

**8. A Diffusion Model.** We shall assume from now on that  $K = n$ , and that the processes  $\beta_i(\cdot)$ ,  $\sigma_{ik}(\cdot)$ ,  $1 \leq i, k \leq n$  in (2.1) are of the form

$$(8.1) \quad \beta_i(t) = b_i(\mathfrak{X}(t)), \quad \sigma_{ik}(t) = s_{ik}(\mathfrak{X}(t)), \quad 0 \leq t < \infty.$$

Here  $\mathfrak{X}(t) = (X_1(t), \dots, X_n(t))'$  is the vector of capitalizations at time  $t$ , and  $b_i : (0, \infty)^n \rightarrow \mathbb{R}$ ,  $s_{ik} : (0, \infty)^n \rightarrow \mathbb{R}$  are continuous functions. We shall denote by  $\mathbf{b}(\cdot) = (b_1(\cdot), \dots, b_n(\cdot))'$  and  $\mathbf{s}(\cdot) = (s_{ik}(\cdot))_{1 \leq i \leq n, 1 \leq k \leq n}$  the vector and matrix, respectively, of these local rate-of-return and local volatility functions. With this setup, the vector process  $\mathfrak{X}(t)$ ,  $0 \leq t < \infty$  of capitalizations becomes a diffusion, with values in  $(0, \infty)^n$  and dynamics

$$(8.2) \quad dX_i(t) = b_i(\mathfrak{X}(t)) dt + \sum_{k=1}^n s_{ik}(\mathfrak{X}(t)) dW_k(t), \quad i = 1, \dots, n,$$

where for  $\mathbf{x} = (x_1, \dots, x_n)' \in (0, \infty)^n$  we set  $\mathbf{a}_{ij}(\mathbf{x}) := \sum_{k=1}^n s_{ik}(\mathbf{x}) s_{jk}(\mathbf{x})$ ,

$$(8.3) \quad \mathbf{b}_i(\mathbf{x}) := x_i b_i(\mathbf{x}), \quad \mathbf{s}_{ik}(\mathbf{x}) := x_i s_{ik}(\mathbf{x}), \quad \mathbf{a}_{ij}(\mathbf{x}) := x_i x_j a_{ij}(\mathbf{x}).$$

This diffusion  $\mathfrak{X}(\cdot)$  has infinitesimal generator

$$(8.4) \quad \mathcal{L}f := \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \mathbf{a}_{ij}(\mathbf{x}) D_{ij}^2 f + \sum_{i=1}^n \mathbf{b}_i(\mathbf{x}) D_i f.$$

**Assumption C:** For every  $\mathbf{x} \in (0, \infty)^n$ , the matrix  $\mathbf{s}(\mathbf{x}) = (s_{ij}(\mathbf{x}))_{1 \leq i, j \leq n}$  is invertible; the system (8.2) has a unique-in-distribution weak solution, with  $\mathfrak{X}(0) = \mathbf{x}$  and values in  $(0, \infty)^n$ ; and for  $\Theta(\mathbf{x}) := \mathbf{s}^{-1}(\mathbf{x}) \mathbf{b}(\mathbf{x})$ , the following analogue of (2.2), (5.1) holds for each  $T \in (0, \infty)$ :

$$(8.5) \quad \sum_{i=1}^n \int_0^T \left( |\mathbf{b}_i(\mathfrak{X}(t))| + \mathbf{a}_{ii}(\mathfrak{X}(t)) + \Theta_i^2(\mathfrak{X}(t)) \right) dt < \infty, \quad \text{a.s.}$$

It follows from this assumption that the Brownian motion  $W(\cdot)$  is adapted to the augmentation of the filtration  $\mathbb{F}^{\mathfrak{X}}$ , and that  $\vartheta(\cdot) = \Theta(\mathfrak{X}(\cdot))$  is a market-price of risk process as postulated in (5.1). The following conditions from Bass & Perkins (2003), in particular their Theorem 1.2 and Corollary 1.3, are sufficient for the existence of a weak solution for (8.2) which is unique in distribution: the functions  $\mathfrak{s}_{ik}(\cdot)$ ,  $\mathbf{b}_i(\cdot)$  of (8.3) can be extended by continuity on all of  $[0, \infty)^n$ ;  $\mathbf{b}_i(\cdot)$  and  $\mathfrak{h}_{ij}(\mathbf{x}) := \sqrt{x_i x_j} \mathbf{a}_{ij}(\mathbf{x}) > 0$  are Hölder continuous on compact subsets of  $[0, \infty)^n$ ; and we have

$$(8.6) \quad \mathbf{b}_i(\mathbf{x}) \geq 0 \text{ for } x_i = 0; \quad \|\mathbf{b}(\mathbf{x})\| + \|\mathfrak{s}(\mathbf{x})\| \leq C(1 + \|\mathbf{x}\|), \quad \forall \mathbf{x} \in [0, \infty)^n$$

and  $\mathfrak{h}_{ij}(\mathbf{x}) = 0$  for  $i \neq j$ ,  $\mathbf{x} \in \mathcal{O}^n$ , where  $\mathcal{O}^n$  is the boundary of  $[0, \infty)^n$ .

**Remark 2:** The diffusion  $\mathfrak{X}(\cdot)$  of (8.2) takes values in  $(0, \infty)^n$ , if and only if the diffusion  $\Xi(\cdot) = (\Xi_1(\cdot), \dots, \Xi_n(\cdot))'$ ,  $\Xi_i(\cdot) := 1/X_i(\cdot)$ , with dynamics

$$(8.7) \quad d\Xi_i(t) = \mathbf{q}_i(\Xi(t)) dt + \sum_{k=1}^n \mathfrak{r}_{ik}(\Xi(t)) dW_k(t), \quad i = 1, \dots, n$$

and  $\mathfrak{r}_{ik}(\xi) := -\xi_i \mathfrak{s}_{ik}(1/\xi_1, \dots, 1/\xi_n)$ ,  $\mathbf{q}_i(\xi) := \xi_i (\mathbf{a}_{ii} - \mathbf{b}_i)(e^{\xi_1}, \dots, e^{\xi_n})$ , takes values in  $(0, \infty)^n$ . Thus, any conditions guaranteeing the existence of a non-explosive solution to the SDEs of (8.7) for all times, such as linear growth for  $\mathbf{q}_i(\cdot)$  and  $\mathfrak{r}_{ik}(\cdot)$ , also ensure that  $\mathfrak{X}(\cdot)$  takes values in  $(0, \infty)^n$ .

Alternatively, one may invoke results of Friedman (2006), Section 9.4 and Chapter 11, to obtain conditions on  $\mathbf{b}_i(\cdot)$ ,  $\mathfrak{s}_{ik}(\cdot)$  under which the diffusion  $\mathfrak{X}(\cdot)$  of (8.2) never attains any of the faces  $\{x_1 = 0\}, \dots, \{x_n = 0\}$  of  $\mathcal{O}^n$ . In particular, if these functions can be extended by continuity on all of  $[0, \infty)^n$ ; the  $\mathfrak{s}_{ik}(\cdot)$  are continuously differentiable; the matrix  $\mathbf{a}(\cdot)$  degenerates on the faces of the orthant; and the so-called *Fichera drifts*

$$(8.8) \quad \mathfrak{f}_i(\mathbf{x}) := \mathbf{b}_i(\mathbf{x}) - \frac{1}{2} \sum_{j=1}^n D_j \mathbf{a}_{ij}(\mathbf{x})$$

are nonnegative on  $\{x_i = 0\}$ , for each  $i = 1, \dots, n$ ; then  $\mathfrak{X}(\cdot)$  takes values in  $(0, \infty)^n$  (see Theorem 9.4.1 and Corollary 9.4.2 in Friedman (2006)).

**Assumption D:** There exists  $H : (0, \infty)^n \rightarrow \mathbb{R}$  of class  $\mathcal{C}^2$ , such that

$$(8.9) \quad \mathbf{b}(\mathbf{x}) = \mathbf{a}(\mathbf{x}) DH(\mathbf{x}), \quad \forall \mathbf{x} \in (0, \infty)^n.$$

In light of Assumption C, this new requirement amounts essentially to postulating that the vector field  $\mathbf{a}^{-1}(\cdot)\mathbf{b}(\cdot)$  be conservative; it is imposed here for technical reasons (cf. discussion in Remark 3). Under it, we have

$$(8.10) \quad \Theta(\mathbf{x}) = \mathfrak{s}'(\mathbf{x})DH(\mathbf{x}), \quad \text{and} \quad \mathfrak{s}(\mathbf{x})\Theta(\mathbf{x}) = \mathfrak{b}(\mathbf{x}), \quad \mathbf{x} \in (0, \infty)^n.$$

*Throughout the remainder, Assumptions B, C, D will be in force, and  $\mathbb{F} \equiv \mathbb{F}_+^{\mathfrak{X}}$ ; this is a natural choice, and consistent with Assumption A.*

**9. A Parabolic PDE for the function  $U(\tau, \mathbf{x})$ .** The uniqueness in distribution posited in Assumption C implies that  $\mathfrak{X}(\cdot)$  is strongly Markovian; we shall denote by  $\mathbb{P}^{\mathbf{x}}$  the distribution of this process started at  $\mathfrak{X}(0) = \mathbf{x} \in (0, \infty)^n$ . Our objective now is to study

$$(9.1) \quad U(T, \mathbf{x}) := \mathbb{E}^{\mathbb{P}^{\mathbf{x}}} [Z(T)X(T)] / (x_1 + \cdots + x_n),$$

the quantity of (6.1), (6.2) in this diffusion context. We start by observing that with  $H(\cdot)$  as in Assumption D and the notation of (8.4) and (8.10), Itô's rule gives  $H(\mathfrak{X}(T)) - H(\mathfrak{X}(0)) - \int_0^T \mathcal{L}H(\mathfrak{X}(t))dt = \int_0^T \Theta'(\mathfrak{X}(t))dW(t)$ , and the exponential local martingale  $Z(\cdot)$  of (5.2) becomes

$$(9.2) \quad Z(\cdot) = \exp \left\{ H(\mathfrak{X}(0)) - H(\mathfrak{X}(\cdot)) - \int_0^\cdot k(\mathfrak{X}(t)) dt \right\}.$$

In particular,  $Z(\cdot)$  is  $\mathbb{F}^{\mathfrak{X}}$ -adapted. We are setting here

$$(9.3) \quad k(\mathbf{x}) := - \sum_{i=1}^n \sum_{j=1}^n \frac{\mathfrak{a}_{ij}(\mathbf{x})}{2} \left[ D_{ij}^2 H(\mathbf{x}) + D_i H(\mathbf{x}) D_j H(\mathbf{x}) \right],$$

$$(9.4) \quad g(\mathbf{x}) := e^{-H(\mathbf{x})} \sum_{i=1}^n x_i, \quad G(T, \mathbf{x}) := \mathbb{E}^{\mathbb{P}^{\mathbf{x}}} \left[ g(\mathfrak{X}(T)) e^{-\int_0^T k(\mathfrak{X}(t))dt} \right].$$

With this notation, the function of (9.1) becomes  $U(T, \mathbf{x}) = G(T, \mathbf{x})/g(\mathbf{x})$ . A bit more generally, these considerations – coupled with the Markov property of  $\mathfrak{X}(\cdot)$  – lead for any  $0 \leq t \leq T$  to the a.s. identity

$$(9.5) \quad \frac{\mathbb{E}^{\mathbb{P}^{\mathbf{x}}} [X(T)Z(T) | \mathcal{F}(t)]}{X(t)Z(t)} = \frac{G(T-t, \mathbf{y})}{g(\mathbf{y})} \Bigg|_{\mathbf{y}=\mathfrak{X}(t)} = U(T-t, \mathfrak{X}(t)).$$

The following Assumption E will also be imposed from now onwards. *It amounts to assuming that the function  $U(\cdot, \cdot)$  of (9.1) is of class  $\mathcal{C}^{1,2}$ .* Note that the equation (9.6) is satisfied, at least in the support of  $\mathfrak{X}(\cdot)$ , thanks to the assumption  $U(\cdot, \cdot) \in \mathcal{C}^{1,2}((0, \infty) \times (0, \infty)^n)$  and to the martingale property of the process  $G(T-t, \mathfrak{X}(t)) e^{-\int_0^t k(\mathfrak{X}(u))du}$ ,  $0 \leq t \leq T$ .

**Assumption E:** The function  $G(\cdot, \cdot)$  in (9.4) takes values in  $(0, \infty)$ , is continuous on  $[0, \infty) \times (0, \infty)^n$ , of class  $\mathcal{C}^{1,2}$  on  $(0, \infty) \times (0, \infty)^n$ , and solves

$$(9.6) \quad \frac{\partial G}{\partial \tau}(\tau, \mathbf{x}) = \mathcal{L}G(\tau, \mathbf{x}) - k(\mathbf{x})G(\tau, \mathbf{x}), \quad \tau \in (0, \infty), \mathbf{x} \in (0, \infty)^n$$

$$(9.7) \quad G(0, \mathbf{x}) = g(\mathbf{x}), \quad \mathbf{x} \in (0, \infty)^n.$$

This Cauchy problem is exactly the one arising in classical Feynman-Kac theory; see, for instance, sections 5.6, 6.5 in Friedman (2006), section 5.7 in Karatzas & Shreve (1991), Janson & Tysk (2006). From Theorem 1 and Remark in Heath & Schweizer (2000), Assumption E holds if: the functions  $\mathfrak{b}_i(\cdot)$ ,  $\mathfrak{s}_{ik}(\cdot)$  of (8.3) are continuously differentiable on  $(0, \infty)$  and satisfy the growth condition in (8.6); the functions  $\mathfrak{a}_{ij}(\cdot)$  of (8.3) satisfy the non-degeneracy condition (9.14) below; the function  $g(\cdot)$  in (9.4) is Hölder continuous, uniformly on compact subsets of  $(0, \infty)^n$ ; the continuous function  $k(\cdot)$  of (9.3) is bounded from below; and the function  $G(\cdot, \cdot)$  in (9.4) is real-valued and continuous on  $(0, \infty) \times (0, \infty)^n$ . This latter requirement is satisfied, for instance, if the functions  $\mathfrak{r}_{ik}(\cdot)$ ,  $\mathfrak{q}_i(\cdot)$  in (8.7) obey linear growth conditions, and the function  $\mathfrak{g}(\xi) := g(1/\xi_1, \dots, 1/\xi_n)$  has polynomial growth; see Problem 5.3.15 in Karatzas & Shreve (1991), as well as Lemma 2 (and the paragraph preceding it) in Heath & Schweizer (2000).

Sustained computation shows then that the Cauchy problem of (9.6), (9.7) for  $G(\cdot, \cdot)$ , leads to a corresponding Cauchy problem for  $U(\cdot, \cdot)$ , namely

$$(9.8) \quad \frac{\partial U}{\partial \tau}(\tau, \mathbf{x}) = \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \mathfrak{a}_{ij}(\mathbf{x}) D_{ij}^2 U(\tau, \mathbf{x}) + \sum_{i=1}^n \sum_{j=1}^n \frac{\mathfrak{a}_{ij}(\mathbf{x}) D_i U(\tau, \mathbf{x})}{x_1 + \dots + x_n}$$

for  $(\tau, \mathbf{x}) \in (0, \infty) \times (0, \infty)^n$ ; and  $U(0, \mathbf{x}) = 1$  for  $\mathbf{x} \in (0, \infty)^n$ .

9.1. *An Informal Derivation of (9.8).* Rather than including the computations which lead from (9.6) to the equation (9.8), we present here a rather simple, informal argument that we shall find useful also in the next subsection, in a more formal setting. We start by casting (6.4) as

$$\frac{d(X(t)Z(t))}{X(t)Z(t)} = \sum_{k=1}^n \left( \sum_{i=1}^n \mu_i(t) \sigma_{ik}(t) - \vartheta_k(t) \right) dW_k(t) = - \sum_{k=1}^n \tilde{\Theta}_k(\mathfrak{X}(t)) dW_k(t)$$

where, by analogy with (6.5), we have set

$$(9.9) \quad \tilde{\Theta}_k(\mathbf{x}) := \Theta_k(\mathbf{x}) - \sum_{i=1}^n \left( \frac{x_i \mathfrak{s}_{ik}(\mathbf{x})}{x_1 + \dots + x_n} \right), \quad k = 1, \dots, n.$$

On the other hand, assuming that  $U(\cdot, \cdot)$  of (9.1) is of class  $\mathcal{C}^{1,2}$ , we obtain from Itô's rule and with  $R_k(\tau, \mathbf{x}) := \sum_{i=1}^n x_i \mathfrak{s}_{ik}(\mathbf{x}) D_i U(\tau, \mathbf{x})$ ,  $k = 1, \dots, n$ :

$$dU(T-t, \mathfrak{X}(t)) = \left( \mathcal{L}U - \frac{\partial U}{\partial \tau} \right) (T-t, \mathfrak{X}(t)) dt + \sum_{k=1}^n R_k(T-t, \mathfrak{X}(t)) dW_k(t).$$

The product rule of the stochastic calculus applied to the process

$$(9.10) \quad N(t) := X(t)Z(t)U(T-t, \mathfrak{X}(t)) = \mathbb{E}^{\mathbb{P}^{\mathbf{x}}} [X(T)Z(T) | \mathcal{F}(t)]$$

of (9.5), leads then to

$$\begin{aligned} \frac{dN(t)}{X(t)Z(t)} &= dU(T-t, \mathfrak{X}(t)) + U(T-t, \mathfrak{X}(t)) \frac{d(X(t)Z(t))}{X(t)Z(t)} - \sum_{k=1}^n R_k(T-t, \mathfrak{X}(t)) \tilde{\Theta}_k(\mathfrak{X}(t)) dt \\ &= C(T-t, \mathfrak{X}(t)) dt + \sum_{k=1}^n [R_k(T-t, \mathfrak{X}(t)) - U(T-t, \mathfrak{X}(t)) \tilde{\Theta}_k(\mathfrak{X}(t))] dW_k(t). \end{aligned}$$

We have set

$$\begin{aligned} C(\tau, \mathbf{x}) &:= \left( \mathcal{L}U - \frac{\partial U}{\partial \tau} \right) (\tau, \mathbf{x}) - \sum_{k=1}^n R_k(\tau, \mathbf{x}) \tilde{\Theta}_k(\mathbf{x}) \\ &= \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \mathbf{a}_{ij}(\mathbf{x}) D_{ij}^2 U(\tau, \mathbf{x}) + \sum_{i=1}^n \sum_{j=1}^n \frac{\mathbf{a}_{ij}(\mathbf{x}) D_i U(\tau, \mathbf{x})}{x_1 + \dots + x_n} - \frac{\partial U}{\partial \tau}(\tau, \mathbf{x}), \end{aligned}$$

where the last equality is checked easily from (8.4) and (8.10). But the process  $N(\cdot)$  of (9.10) is a martingale, so the term  $C(\tau, \mathbf{x})$  should vanish:

$$(9.11) \quad \frac{dN(t)}{N(t)} = \sum_{k=1}^n \left[ \frac{R_k(T-t, \mathfrak{X}(t))}{U(T-t, \mathfrak{X}(t))} - \tilde{\Theta}_k(\mathfrak{X}(t)) \right] dW_k(t).$$

In other words, the function  $U : [0, \infty) \times (0, \infty)^n \rightarrow (0, 1]$  of (9.3) must satisfy the parabolic partial differential equation (9.8), as postulated earlier.

**Remark 3:** This informal derivation suggests that it may be possible to dispense with Assumptions D, E altogether, if it can be shown from first principles that the function  $U$  of (9.1) is of class  $\mathcal{C}^{1,2}$  on  $(0, \infty) \times (0, \infty)^n$ . Indeed, under suitable conditions, one can rely on techniques from the Malliavin calculus and the Hörmander hypoellipticity theorem (Nualart (1995), pp. 99-124) to show that the  $(n+2)$ -dimensional vector  $(\mathfrak{X}(T), \Upsilon(T), \Xi(T))$  with  $\Upsilon(T) := \int_0^T \Theta(\mathfrak{X}(t))' dW(t)$  and  $\Xi(T) := \int_0^T \|\Theta(\mathfrak{X}(t))\|^2 dt$  has an infinitely differentiable probability density function, for any given  $T \in (0, \infty)$ . This provides the requisite smoothness for the function

$$U(T, \mathbf{x}) = \frac{1}{x_1 + \dots + x_n} \mathbb{E}^{\mathbb{P}^{\mathbf{x}}} \left[ (X_1(T) + \dots + X_n(T)) e^{\Upsilon(T) - (\Xi(T)/2)} \right].$$

The conditions needed for this approach to work are strong; they include the infinite-differentiability of the functions  $\mathfrak{s}_{ik}(\cdot)$ ,  $\Theta_i(\cdot)$ ,  $1 \leq i, k \leq n$ , as well as additional algebraic conditions which, in the present context, are somewhat opaque and not very easy to state or verify. For these reasons we have opted for sticking with Assumptions D, E; these are satisfied in the Examples of section 12, are easy to test, and allow us to represent Föllmer's exit measure via (9.23), (9.24) without involving stochastic integrals.

**9.2. Results and Ramifications.** The equation (9.8) is determined *entirely* from the volatility structure of the model (2.1). Furthermore, the Cauchy problem of (9.8),  $U(0, \cdot) = 1$  admits the trivial solution  $U(\tau, \mathbf{x}) \equiv 1$ ; thus, the existence of arbitrage relative to the market portfolio over a finite time-horizon  $[0, T]$  is tantamount to *failure of uniqueness* for the Cauchy problem of (9.8),  $U(0, \cdot) = 1$  over the strip  $[0, T] \times (0, \infty)^n$ .

**Remark 4:** Assume there exists some  $h > 0$  such that the continuous functions  $a_{ij}(\cdot)$ ,  $1 \leq i, j \leq n$  satisfy either of the conditions

$$(9.12) \quad (x_1 + \cdots + x_n) \sum_{i=1}^n x_i a_{ii}(\mathbf{x}) - \sum_{i=1}^n \sum_{j=1}^n x_i x_j a_{ij}(\mathbf{x}) \geq h (x_1 + \cdots + x_n)^2,$$

$$(9.13) \quad (x_1 \cdots x_n)^{1/n} \left[ \sum_{i=1}^n a_{ii}(\mathbf{x}) - \frac{1}{n} \sum_{i=1}^n \sum_{j=1}^n a_{ij}(\mathbf{x}) \right] \geq h (x_1 + \cdots + x_n),$$

for all  $\mathbf{x} \in (0, \infty)^n$  (we have just re-written (4.2), (4.3) in the present context). Then from the results reviewed in section 4 we deduce that, for all  $T > (2 \log n)/h$  under (9.12), and for all  $T > (2n^{1-(1/n)})/h$  under (9.13), we have:  $U(T, \mathbf{x}) < 1$ ,  $\forall \mathbf{x} \in (0, \infty)^n$ . In particular, under either (9.12) or (9.13), *uniqueness fails for the Cauchy problem of (9.8)*,  $U(0, \cdot) \equiv 1$ .

Whenever uniqueness fails for this problem, it is important to know how to pick the 'right' solution from among all possible solutions, the one which gives the quantity of (9.1). The next result addresses this issue; it implies that  $G(\cdot, \cdot)$  in (9.4) is the *smallest nonnegative*, continuous function, of class  $\mathcal{C}^{1,2}((0, \infty) \times (0, \infty)^n)$ , which satisfies  $(\partial G / \partial \tau) \geq \mathcal{L}G - kG$  and (9.7) (cf. Karatzas & Shreve (1991), Exercise 4.4.7 for a similar situation).

**Theorem 1:** *The function  $U : [0, \infty) \times (0, \infty)^n \rightarrow (0, 1]$  of (9.1) is the smallest non-negative continuous function, of class  $\mathcal{C}^{1,2}$  on  $(0, \infty) \times (0, \infty)^n$ , that satisfies  $U(0, \cdot) \equiv 1$  and (1.1).*

*Proof:* Consider any continuous function  $\tilde{U} : [0, \infty) \times (0, \infty)^n \rightarrow [0, \infty)$  of class  $\mathcal{C}^{1,2}$  on  $(0, \infty) \times (0, \infty)^n$  that satisfies both (1.1) and  $\tilde{U}(0, \cdot) \equiv 1$ , and define as in (9.10) the process  $\tilde{N}(t) := X(t)Z(t)\tilde{U}(T-t, \mathfrak{X}(t))$ ,  $0 \leq t \leq T$ .

Repeating verbatim the arguments in subsection 9.1, we use (1.1) to conclude that the nonnegative process  $\tilde{N}(\cdot)$  is a local supermartingale. Thus  $\tilde{N}(\cdot)$  is bona-fide supermartingale,  $(x_1 + \cdots + x_n) \tilde{U}(T, \mathbf{x}) = \tilde{N}(0) \geq \mathbb{E}^{\mathbb{P}^{\mathbf{x}}}(\tilde{N}(T)) = \mathbb{E}^{\mathbb{P}^{\mathbf{x}}}(X(T)Z(T))$  holds for every  $(T, \mathbf{x}) \in (0, \infty) \times (0, \infty)^n$ , and  $\tilde{U}(T, \mathbf{x}) \geq U(T, \mathbf{x})$  follows from (9.1).  $\square$

**Proposition 2:** *Assume that the continuous functions  $(\mathbf{a}_{ij}(\cdot))_{1 \leq i, j \leq n}$  of (8.3) satisfy the following non-degeneracy condition: for every compact subset  $\mathcal{K}$  of  $(0, \infty)^n$ , there exists a number  $\varepsilon = \varepsilon_{\mathcal{K}} > 0$  such that*

$$(9.14) \quad \sum_{i=1}^n \sum_{j=1}^n \mathbf{a}_{ij}(\mathbf{z}) \xi_i \xi_j \geq \varepsilon \|\xi\|^2, \quad \forall \mathbf{z} \in \mathcal{K}, \xi \in \mathbb{R}^n.$$

Then, if

$$(9.15) \quad U(T, \mathbf{x}) < 1 \quad \text{for some } \mathbf{x} \in (0, \infty)^n$$

holds for some  $T \in (0, \infty)$ , we have

$$(9.16) \quad U(T, \mathbf{x}) < 1, \quad \forall (T, \mathbf{x}) \in (0, \infty) \times (0, \infty)^n.$$

*Proof:* Let us work first under the stronger assumption

$$(9.17) \quad U(T, \mathbf{x}) < 1, \quad \forall \mathbf{x} \in (0, \infty)^n,$$

for some  $T \in (0, \infty)$ . For every  $\tau > 0$ , we consider the set  $\mathcal{S}(\tau) := \{\mathbf{x} \in (0, \infty)^n \mid U(\tau, \mathbf{x}) = 1\}$  and define  $\tau_* := \sup\{\tau \in (0, \infty) \mid \mathcal{S}(\tau) \neq \emptyset\}$  (with  $\tau_* = 0$  if the set is empty). The assumption (9.17) amounts to  $\tau_* < \infty$ , and the claim (9.16) to  $\tau_* = 0$ ; we shall prove this claim by contradiction.

Suppose  $\tau_* > 0$ ; then  $U(\tau_* - \delta, \mathbf{x}_*) = 1$  for any given  $\delta \in (0, \tau_*/2)$ , and some  $\mathbf{x}_* \in (0, \infty)^n$ . For any given  $\mathbf{x} \in (0, \infty)^n$ , consider an open, connected set  $D$  which contains both  $\mathbf{x}$  and  $\mathbf{x}_*$ , and whose closure  $\bar{D}$  is a compact subset of  $(0, \infty)^n$ ; in particular, we have  $\inf\{\|\mathbf{y} - \mathbf{z}\| \mid \mathbf{z} \in \bar{D}, \mathbf{y} \in \mathcal{O}^n\} > 0$ . The function  $U(\cdot, \cdot)$  attains its maximum value over the cylindrical domain  $\mathfrak{E} = \{(\tau, \xi) \mid 0 < \tau < \tau_* + 1, \xi \in D\}$  at the point  $(\tau_* - \delta, \mathbf{x}_*)$ , which lies in the interior of this domain. By assumption then, the operator  $\hat{\mathcal{L}}f = (1/2) \sum_{i=1}^n \sum_{j=1}^n \mathbf{a}_{ij}(\mathbf{x}) D_{ij}^2 f + \sum_{i=1}^n \hat{\mathbf{b}}_i(\mathbf{x}) D_i f$  of (1.2) with

$$(9.18) \quad \hat{\mathbf{b}}_i(\mathbf{x}) := x_i \hat{\mathbf{b}}_i(\mathbf{x}), \quad \hat{\mathbf{b}}_i(\mathbf{x}) := \sum_{j=1}^n \frac{x_j \mathbf{a}_{ij}(\mathbf{x})}{x_1 + \cdots + x_n}, \quad i = 1, \dots, n,$$

is uniformly parabolic with bounded, continuous coefficients on  $\mathfrak{E}$ , so from the maximum principle for parabolic operators (Friedman (2006), Ch. 6):

$$(9.19) \quad U(\tau, \mathbf{x}) = 1, \quad \forall (\tau, \mathbf{x}) \in [0, \tau_* - \delta) \times (0, \infty)^n.$$

Now let us recall the  $\mathbb{P}^{\mathbf{x}}$ -a.s. equality  $\mathbb{E}^{\mathbb{P}^{\mathbf{x}}}[X(T)Z(T) | \mathcal{F}(t)] = U(T - t, \mathfrak{X}(t)) \cdot X(t)Z(t)$  from (9.5); we apply it with  $0 \leq t \leq \tau_* - \delta$ ,  $0 \leq T - t \leq \tau_* - \delta$ , then take expectations with respect to the probability measure  $\mathbb{P}^{\mathbf{x}}$ , and use (9.19) along with (9.1), to obtain for every  $T \in [0, 2(\tau_* - \delta)]$ :

$$U(T, \mathbf{x}) = \frac{\mathbb{E}^{\mathbb{P}^{\mathbf{x}}}[X(T)Z(T)]}{x_1 + \cdots + x_n} = \frac{\mathbb{E}^{\mathbb{P}^{\mathbf{x}}}[X(t)Z(t)]}{x_1 + \cdots + x_n} = U(t, \mathbf{x}) = 1, \quad \forall \mathbf{x} \in (0, \infty)^n.$$

But since  $2(\tau_* - \delta) > \tau_*$ , this contradicts the definition of  $\tau_*$ .

Now we revert to (9.15); as J. Ruf (private communication) observes, yet another application of the maximum principle, as above, leads to (9.17).  $\square$

**Corollary:** *Under the non-degeneracy condition (9.14), and with either (9.12) or (9.13), the inequality (9.16) holds. That is, arbitrage with respect to the market exists then over any time-horizon  $[0, T]$  with  $T \in (0, \infty)$ .*

**9.3. An Auxiliary Diffusion.** Let us consider now the diffusion process  $\mathfrak{Y}(\cdot)$  with infinitesimal generator  $\hat{\mathcal{L}}$  as in (1.2), (9.18) and dynamics

$$(9.20) \quad dY_i(t) = \hat{\mathbf{b}}_i(\mathfrak{Y}(t)) dt + \sum_{k=1}^n \mathfrak{s}_{ik}(\mathfrak{Y}(t)) dW_k(t), \quad i = 1, \dots, n.$$

**Assumption F:** *The system of SDEs (9.20) admits a unique-in-distribution weak solution with values in  $[0, \infty)^n \setminus \{\mathbf{0}\}$ .*

This will be the case, for instance, if the drift functions  $\hat{\mathbf{b}}_i(\cdot)$ ,  $1 \leq i \leq n$  of (9.18) can be extended by continuity on all of  $[0, \infty)^n$  and satisfy the Bass & Perkins (2003) conditions preceding, following and including (8.6). The resulting process  $\mathfrak{Y}(\cdot)$  is then Markovian, and we shall denote by  $\mathbb{Q}^{\mathfrak{Y}}$  its distribution with  $\mathfrak{Y}(0) = \mathbf{y} \in [0, \infty)^n$ . Unlike the original process  $\mathfrak{X}(\cdot)$ , which takes values in  $(0, \infty)^n$ , this new process  $\mathfrak{Y}(\cdot)$  is only guaranteed to take values in the non-negative orthant  $[0, \infty)^n \setminus \{\mathbf{0}\}$ . In particular, with  $\mathbf{x} \in (0, \infty)^n$  the first hitting time

$$(9.21) \quad \mathfrak{T} := \inf\{t \geq 0 \mid \mathfrak{Y}(t) \in \mathcal{O}^n\}$$

of the boundary  $\mathcal{O}^n$  of  $[0, \infty)^n$  may be finite with positive  $\mathbb{Q}^{\mathfrak{X}}$ -probability.

Our next result shows that this possibility amounts to the existence of arbitrage relative to the market, and to the lack of uniqueness for the Cauchy problem of (9.8) and  $U(0, \cdot) \equiv 1$ .

**Theorem 2:** *With the above notation and assumptions, including (9.14), the function  $U : [0, \infty) \times (0, \infty)^n \rightarrow (0, 1]$  of (9.1) admits the representation*

$$(9.22) \quad U(T, \mathbf{x}) = \mathbb{Q}^{\mathfrak{X}}[\mathfrak{T} > T], \quad (T, \mathbf{x}) \in (0, \infty) \times (0, \infty)^n.$$

*Proof:* The function on the right-hand side of (9.22) is space-time harmonic for the diffusion  $\mathfrak{Y}(\cdot)$  on  $(0, \infty) \times (0, \infty)^n$ , so it solves the equation (9.8) there (cf. Thm. 2.7 in Janson & Tysk (2006)). Consider any continuous  $V : [0, \infty) \times (0, \infty)^n \rightarrow [0, \infty)$  of class  $\mathcal{C}^{1,2}$  on  $(0, \infty) \times (0, \infty)^n$ , which satisfies (1.1) and  $V(0, \cdot) \equiv 1$  on  $(0, \infty)^n$ . Then  $V(T-t, \mathfrak{Y}(t))$ ,  $0 \leq t \leq T$  is a non-negative local supermartingale, thus a true supermartingale; we deduce

$$V(T, \mathbf{x}) \geq \mathbb{E}^{\mathbb{Q}^{\mathbf{x}}} \left[ V(T - (T \wedge \mathfrak{T}), \mathfrak{Y}(T \wedge \mathfrak{T})) \right] \geq \mathbb{E}^{\mathbb{Q}^{\mathbf{x}}} \left[ V(0, \mathfrak{Y}(T)) 1_{\{\mathfrak{T} > T\}} \right],$$

thus also  $V(T, \mathbf{x}) \geq \mathbb{Q}^{\mathbf{x}}[\mathfrak{T} > T]$  for every  $(T, \mathbf{x}) \in (0, \infty) \times (0, \infty)^n$ . The claim follows now from Theorem 1.  $\square$

**Corollary:** *Under the assumptions of Theorem 2, for any given  $\mathbf{x} \in (0, \infty)^n$  the  $\mathbb{P}^{\mathbf{x}}$ -supermartingale  $Z(\cdot)X(\cdot)$  is under  $\mathbb{P}^{\mathbf{x}}$  a*

- martingale, if and only if  $\mathbb{Q}^{\mathbf{x}}(\mathfrak{T} < \infty) = 0$ ;
- potential (i.e.,  $\lim_{T \rightarrow \infty} \mathbb{E}^{\mathbb{P}^{\mathbf{x}}}(Z(T)X(T)) = 0$ ), iff  $\mathbb{Q}^{\mathbf{x}}(\mathfrak{T} < \infty) = 1$ ;
- strict local (and super)martingale on any time-horizon  $[0, T]$  with  $T \in (0, \infty)$ , if and only if  $\mathbb{Q}^{\mathbf{x}}(\mathfrak{T} < \infty) > 0$ .

We represent by analogy with (7.3) the exit measure  $\mathfrak{P}^{\mathbf{x}}$  of the supermartingale  $Z(\cdot)X(\cdot)$  with initial configuration  $\mathfrak{X}(0) = \mathbf{x}$ , in the form

$$(9.23) \quad \mathfrak{P}^{\mathbf{x}}((T, \infty] \times \Omega) = U(T, \mathbf{x}) = \mathbb{Q}^{\mathbf{x}}[\mathfrak{T} > T],$$

and from (9.2)-(9.5) we have for  $A \in \mathcal{F}(t)$ ,  $0 \leq t \leq T$ :

$$(9.24) \quad \mathfrak{P}^{\mathbf{x}}((T, \infty] \times A) = \mathbb{E}^{\mathbb{P}^{\mathbf{x}}} \left[ \frac{g(\mathfrak{X}(t))}{g(\mathbf{x})} 1_A \left( \mathbb{Q}^{\mathbf{x}}[\mathfrak{T} > T-t] \right) \Big|_{\mathbf{z}=\mathfrak{X}(t)} e^{-\int_0^t k(\mathfrak{X}(s)) ds} \right].$$

When  $\mathbf{x} \in (0, \infty)^n$  and the quantity of (9.22) is equal to one, the  $\mathbb{Q}^{\mathbf{x}}$ -distribution of the process  $\mathfrak{Y}(t)$ ,  $0 \leq t \leq T$  in (9.20) is the same as the  $\widehat{\mathbb{P}}_T^{\mathbf{x}}$ -distribution of the original stock-price process  $\mathfrak{X}(t)$ ,  $0 \leq t \leq T$ ; this follows by comparing (9.20), (9.18) with (6.6), and denoting by  $\widehat{\mathbb{P}}_T^{\mathbf{x}}$  the probability measure  $\widehat{\mathbb{P}}_T$  of (6.10) with  $\mathfrak{X}(0) = \mathbf{x}$ . We have in this spirit the following result, by analogy with Remark 2.

**Proposition 3:** *Under the assumptions of Theorem 2, suppose that the functions  $\mathfrak{s}_{ik}(\cdot)$  are continuously differentiable on  $(0, \infty)^n$ ; that the matrix  $\mathfrak{a}(\cdot)$  degenerates on  $\mathcal{O}^n$ ; and that the analogues of (8.8), the Fichera drifts (9.25)*

$$\widehat{\mathfrak{f}}_i(\mathbf{x}) := \widehat{\mathfrak{b}}_i(\mathbf{x}) - \frac{1}{2} \sum_{j=1}^n D_j \mathfrak{a}_{ij}(\mathbf{x}) = \sum_{j=1}^n \left( \frac{\mathfrak{a}_{ij}(\mathbf{x})}{x_1 + \cdots + x_n} - \frac{1}{2} D_j \mathfrak{a}_{ij}(\mathbf{x}) \right)$$

for the process  $\mathfrak{Y}(\cdot)$  of (9.20), can be extended by continuity on  $[0, \infty)^n$ .

If  $\widehat{f}_i(\cdot) \geq 0$  holds on each face  $\{x_i = 0\}$ ,  $i = 1, \dots, n$  of the orthant, then we have  $U(\cdot, \cdot) \equiv 1$  in (9.22), and no arbitrage with respect to the market portfolio exists on any time-horizon.

If, on the other hand, we have  $\widehat{f}_i(\cdot) < 0$  on each face  $\{x_i = 0\}$  of the orthant, then  $U(\cdot, \cdot) < 1$  in (9.22) and arbitrage with respect to the market portfolio exists, on every time-horizon  $[0, T]$  with  $T \in (0, \infty)$ .

*Proof:* In light of Theorem 2, the first claim follows from Theorem 9.4.1, Corollary 9.4.2 of Friedman (2006), and the second is a consequence of the support theorem for diffusions, Ikeda & Watanabe (1989) section VI.8.  $\square$

**Remark 5:** (i) The ‘relative weights’  $\nu_i(t) := Y_i(t)/(Y_1(t) + \dots + Y_n(t))$ ,  $i = 1, \dots, n$  have dynamics similar to (6.9):

$$(9.26) \quad d\nu_i(t) = \nu_i(t) (\mathbf{e}_i - \nu(t))' s(\mathfrak{Y}(t)) dW(t).$$

They are thus  $\mathbb{Q}^{\mathbf{x}}$ -martingales with values in  $[0, 1]$  (cf. subsection 6.1); so, when any one of them hits either boundary point of the unit interval, it gets absorbed there. In terms of them, the first hitting time of (9.21) can be expressed as in (7.4):  $\mathfrak{T} = \min_{1 \leq i \leq n} \mathfrak{T}_i$ , where  $\mathfrak{T}_i := \inf\{t \geq 0 \mid \nu_i(t) = 0\}$ .

(ii) The measure  $\mathbb{Q}^{\mathbf{x}}$  corresponds to a *change of drift*, from  $\mathfrak{b}(\cdot)$  in (8.2) to  $\widehat{\mathfrak{b}}(\cdot)$  in (9.18), (9.20); this ensures that, under  $\mathbb{Q}^{\mathbf{x}}$ , the components of the new, ‘fictitious’ market portfolio  $\nu(\cdot)$  are martingales, that  $\nu(\cdot)$  has the numéraire property, and thus and that  $\nu(\cdot)$  cannot be outperformed.

**10. Markovian Market Weights.** Let us assume now the form

$$\mathfrak{b}_i(\mathbf{x}) = \mathfrak{B}_i(x_1/x, \dots, x_n/x), \quad s_{ik}(\mathbf{x}) = \mathfrak{S}_{ik}(x_1/x, \dots, x_n/x)$$

for the functions of (8.1), with  $x := \sum_{j=1}^n x_j$  and suitable continuous functions  $\mathfrak{B}_i(\cdot)$ ,  $\mathfrak{S}_{ik}(\cdot)$  on  $\Delta_+^n$ . For  $\mathfrak{m} = (m_1, \dots, m_n)' \in \Delta_+^n$ , we set  $\mathcal{A}_{ij}(\mathfrak{m}) := \sum_{k=1}^n \mathfrak{S}_{ik}(\mathfrak{m}) \mathfrak{S}_{jk}(\mathfrak{m})$ . In words, we consider instantaneous growth rates and volatilities that depend at time  $t$  only on the current configuration  $\mu(t) = (\mu_1(t), \dots, \mu_n(t))'$  of relative market weights, so the process  $\mu(\cdot)$  of (3.3) is now a diffusion with values in the positive simplex  $\Delta_+^n$  and

$$(10.1) \quad d\mu_i(t) = \mu_i(t) \left[ \Gamma_i(\mu(t)) dt + \sum_{k=1}^n \mathcal{T}_{ik}(\mu(t)) dW_k(t) \right], \quad i = 1, \dots, n,$$

with  $\mathcal{T}_{ik}(\mathfrak{m}) := \mathfrak{S}_{ik}(\mathfrak{m}) - \sum_{j=1}^n m_j \mathfrak{S}_{jk}(\mathfrak{m})$ ,  $\mathcal{P}_{ij}(\mathfrak{m}) := \sum_{k=1}^n \mathcal{T}_{ik}(\mathfrak{m}) \mathcal{T}_{jk}(\mathfrak{m})$ ,

$$\Gamma_i(\mathfrak{m}) := \mathfrak{B}_i(\mathfrak{m}) - \sum_{j=1}^n m_j \mathfrak{B}_j(\mathfrak{m}) - \sum_{j=1}^n m_j \mathcal{A}_{ij}(\mathfrak{m}) + \sum_{j=1}^n \sum_{k=1}^n m_j m_\ell \mathcal{A}_{j\ell}(\mathfrak{m}).$$

In this setup, the function of (9.1) can be expressed in the form  $U(T, \mathbf{x}) = Q(T, x_1/x, \dots, x_n/x)$ , in terms of a function  $Q : (0, \infty) \times \Delta_+^n \rightarrow (0, 1]$  that satisfies the initial condition  $Q(0, \cdot) \equiv 1$  and the equation

$$\frac{\partial Q}{\partial \tau}(\tau, \mathbf{m}) = \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n m_i m_j \mathcal{P}_{ij}(\mathbf{m}) D_{ij}^2 Q(\tau, \mathbf{m}), \quad (\tau, \mathbf{m}) \in (0, \infty) \times \Delta_+^n,$$

which appears on p. 56 of Fernholz (2002) and can be derived from (9.8). On the other hand, by analogy with Theorem 2 and (9.26), the quantity  $Q(T, \mathbf{m})$  is the probability that the process  $\nu(\cdot) = (\nu_1(\cdot), \dots, \nu_n(\cdot))'$  with  $\nu(0) = \mathbf{m} \in \Delta_+^n$  and dynamics (10.2) below, does not hit the boundary of the non-negative simplex  $\Delta^n := \{\mathbf{m} \in [0, 1]^n \mid \sum_{i=1}^n m_i = 1\}$  before  $t = T$ :

$$(10.2) \quad d\nu_i(t) = \nu_i(t) \sum_{k=1}^n \mathcal{T}_{ik}(\nu(t)) dW_k(t), \quad i = 1, \dots, n.$$

**11. The Investment Strategy.** Let us substitute now the expressions of (9.9) into (9.11), to obtain the dynamics of the martingale  $N(\cdot) \equiv Z(\cdot)X(\cdot)U(T - \cdot, \mathfrak{X}(\cdot))$  in (9.10), with  $N(0) = \xi := X(0)U(T, \mathfrak{X}(0))$ :

$$N(t) = \xi + \sum_{k=1}^n \int_0^t N(s) \Psi_k(T - s, \mathfrak{X}(s)) dW_k(s), \quad 0 \leq t \leq T,$$

$$\Psi_k(\tau, \mathbf{x}) := \sum_{i=1}^n s_{ik}(\mathbf{x}) \left( x_i D_i \log U(\tau, \mathbf{x}) + \frac{x_i}{x_1 + \dots + x_n} \right) - \Theta_k(\mathbf{x}).$$

Thus we can identify the “replicating strategy”  $\hat{\pi}(\cdot)$  of (6.2) as

$$(11.1) \quad \hat{\pi}_i(t) = X_i(t) D_i \log U(T - t, \mathfrak{X}(t)) + (X_i(t)/X(t)), \quad i = 1, \dots, n,$$

and its value as  $V^{\xi, \hat{\pi}}(t) = N(t)/Z(t) = X(t)U(T - t, \mathfrak{X}(t))$ ,  $0 \leq t \leq T$ .

**Remark 6:** In the special case of a Markovian model (10.1) for the market weights of  $\mu(\cdot) = (\mu_1(\cdot), \dots, \mu_n(\cdot))'$ , the expression (11.1) takes the form

$$\hat{\pi}_i(t) = \mu_i(t) \left( 1 + D_i \log Q(T - t, \mu(t)) - \sum_{j=1}^n \mu_j(t) D_j \log Q(T - t, \mu(t)) \right)$$

of a “functionally-generated portfolio” in the terminology of Fernholz (2002), p. 56; whereas the value is  $V^{\xi, \hat{\pi}}(t) = X(t)Q(T - t, \mu(t))$ ,  $0 \leq t \leq T$ .

In this case we have  $\sum_{i=1}^n \hat{\pi}_i(\cdot) \equiv 1$ : the strategy that implements the best possible arbitrage relative to the equity market never borrows or lends.

**12. Examples.** We discuss in this section two illustrative examples. Additional examples, in which the investment strategy  $\hat{\pi}(\cdot)$  of (11.1) that realizes the optimal arbitrage can be computed in closed form in dimension  $n = 1$ , can be found in Ruf (2009).

For the first of these examples, take  $n = 1$ ,  $\beta(t) = 1/X^2(t)$  and  $\sigma(t) = 1/X(t)$  in (2.1), where the process  $X(\cdot)$  satisfies  $dX(t) = (1/X(t))dt + dW(t)$  and  $X(0) = 1$ . This is a Bessel process in dimension three – the radial part of a 3-D Brownian motion started at unit distance from the origin – and takes values in  $(0, \infty)$ . We have then  $\vartheta(t) = 1/X(t)$ ,  $Z(t) = 1/X(t)$  for  $0 \leq t < \infty$  in (5.1), (5.2), so  $Z(\cdot)X(\cdot)$  is very clearly a martingale. However,  $Z(\cdot)$  is the prototypical example of a *strict* local martingale – we have  $\mathbb{E}(Z(T)) < 1$  for every  $T \in (0, \infty)$  (e.g., Exercise 3.36, p.168 in Karatzas & Shreve (1991)). This example is taken from Karatzas & Kardaras (2007), p.469, where an arbitrage with respect to the money-market is constructed in closed form. It illustrates that it is possible for  $Z(\cdot)$  to be a strict local martingale and  $Z(\cdot)X(\cdot)$  to be a martingale; in other words, the second and third inequalities in (5.5) fail, while the first stands.

Here we have  $\Theta(x) = 1/x$ ,  $H(x) = \log x$  and  $k(\cdot) \equiv 0$ ,  $g(\cdot) \equiv 1$ ,  $G(\cdot, \cdot) \equiv 1$  in (9.3), (9.4), thus  $U(T, x) \equiv 1$  for all  $T \in [0, \infty)$ ,  $x \in (0, \infty)$ . Arbitrage relative to  $X(\cdot)$  does not exist here, despite the existence of arbitrage relative to the money market and the fact that  $Z(\cdot)$  is a strict local martingale. Note that  $\hat{\mathbf{b}}(x) = 1/x$  in (9.18), so the diffusion of (9.20) is again a Bessel process in dimension three:  $dY(t) = (1/Y(t))dt + dW(t)$ ,  $Y(0) = y > 0$ . This process never hits the origin, so the probability in (9.22) is equal to one, for all  $T \in [0, \infty)$ .

12.1. *The Volatility-Stabilized Model.* Our second example is the model of “stabilization by volatility”, introduced in Fernholz & Karatzas (2005) and studied further by Goia (2009). With  $n \geq 2$ ,  $\zeta \in [0, 1]$  this posits

$$(12.1) \quad \beta_i(t) = (1+\zeta)/(2\mu_i(t)), \quad \sigma_{ik}(t) = \delta_{ik}(\mu_i(t))^{-1/2}; \quad 1 \leq i, k \leq n,$$

that is, rates of return and volatilities which are large for the small stocks and small for the large stocks. The conditions of Bass & Perkins (2003) hold for the resulting system of SDEs, in the notation of (3.2) with  $\kappa := (1+\zeta)/2$ :

$$(12.2) \quad dX_i(t) = \kappa X_i(t) dt + \sqrt{X_i(t) X(t)} dW_i(t), \quad i = 1, \dots, n.$$

The unique-in-distribution solution of (12.2) is expressed in terms of independent Bessel processes  $\mathfrak{R}_1(\cdot), \dots, \mathfrak{R}_n(\cdot)$  in dimension  $4\kappa$  with  $X_i(t) = \mathfrak{R}_i^2(A(t)) > 0$  and  $A(t) := (1/4) \int_0^t X(s) ds$ . In particular,  $\mathfrak{X}(\cdot)$  takes values in  $(0, \infty)^n$ ; for more details on these Lamperti-like descriptions and their

implications, see Fernholz & Karatzas (2005), Goia (2009). The condition (8.5) is satisfied in this example, so Assumption C also holds.

For the model of (12.1), we have  $\Theta_i(\mathbf{x})/\kappa = s_{ii}(\mathbf{x}) = ((x_1 + \dots + x_n)/x_i)^{1/2}$   
 $\mathbf{b}_i(\mathbf{x}) = \kappa(x_1 + \dots + x_n)$ ,  $\mathbf{h}_{ij}(\mathbf{x}) = \delta_{ij}(x_1 + \dots + x_n)$ ,  $\mathbf{a}_{ij}(\mathbf{x}) = x_i \mathbf{h}_{ij}(\mathbf{x})$   
for  $1 \leq i, j \leq n$ . The assumptions of Theorem 2 and of Propositions 1, 3 are all satisfied here, as are (7.5) and (8.9) with  $H(\mathbf{x}) = \kappa \sum_{i=1}^n \log x_i$   
and  $k(\mathbf{x}) = (1 - \zeta^2)(x_1 + \dots + x_n) \sum_{j=1}^n (1/(8x_j))$ . This function  $k(\cdot)$  is  
non-negative, since we have assumed  $0 \leq \zeta \leq 1$ , whereas  $g(\mathbf{x}) = (x_1 + \dots + x_n)(x_1 \dots x_n)^{-\kappa}$ . In particular, with  $\zeta = 1$  we get

$$(12.3) \quad U(T, \mathbf{x}) = \frac{x_1 \dots x_n}{x_1 + \dots + x_n} \mathbb{E}^{\mathbb{P}^{\mathbf{x}}} \left[ \frac{X_1(T) + \dots + X_n(T)}{X_1(T) \dots X_n(T)} \right];$$

see Goia (2009) for a computation of the joint density of  $X_1(T), \dots, X_n(T)$ , which leads then to an explicit computation of  $U(T, \mathbf{x})$  in (12.3) above, and shows that this function is indeed of class  $\mathcal{C}^{1,2}$ .

- With  $\zeta = 1$  one computes  $Z(t) = \prod_{j=1}^n (X_j(0)/X_j(t))$ , therefore  $\Lambda(t) = (X(t)/X(0))^{n-1} \prod_{j=1}^n (\mu_j(t)/\mu_j(0))$  as well as  $\Lambda_i(t) = (X(t)/X(0))^{n-1} \prod_{j \neq i} (\mu_j(t)/\mu_j(0))$  for  $i = 1, \dots, n$ . Both representations in (7.4) hold for the first hitting time of (7.1) in this case; whereas  $\mathcal{S} = \mathcal{T} = \min_{1 \leq i \leq n} \mathcal{T}_i$  as in (7.1)-(7.6), since  $L(t) = (1/Z(t)) = (X(t)/X(0))^n \prod_{j=1}^n (\mu_j(t)/\mu_j(0))$ .
- Both (9.12), (9.13) hold for the example of (12.1) with  $h = n - 1$ , the first as equality; from the Corollary to Proposition 2 and Remark 3, (9.16) holds. We recover the result of Banner & Fernholz (2008) on the existence of arbitrage relative to market of (12.1) over *arbitrary* time-horizons.

The diffusion process  $\mathfrak{Y}(\cdot)$  of (9.20) takes now the form

$$(12.4) \quad dY_i(t) = Y_i(t) dt + \sqrt{Y_i(t)(Y_1(t) + \dots + Y_n(t))} dW_i(t).$$

The conditions of Bass & Perkins (2002) are satisfied again, though one should compare the “weak drift”  $\widehat{\mathbf{b}}_i(\mathbf{x}) = x_i \geq 0$  in (12.4), which vanishes for  $x_i = 0$ , with the “strong drift”  $\mathbf{b}_i(\mathbf{x}) = \kappa(x_1 + \dots + x_n)$  for the the original diffusion  $\mathfrak{X}(\cdot)$  in (12.2), which is strictly positive on  $[0, \infty)^n \setminus \{\mathbf{0}\}$ .

The corresponding Fichera drifts in (9.25), (8.8) are given by  $2\widehat{\mathbf{f}}_i(\mathbf{x}) = x_i - (x_1 + \dots + x_n)$ ,  $2\mathbf{f}_i(\mathbf{x}) = \zeta(x_1 + \dots + x_n) - x_i$ , and  $\mathbf{f}_i(\mathbf{x}) > 0 > \widehat{\mathbf{f}}_i(\mathbf{x})$  hold on  $\{x_i = 0\} \cap \{\sum_{j \neq i} x_j > 0\}$ ; from Remark 2 we verify again that the diffusion  $\mathfrak{X}(\cdot)$  of (12.2) takes values in  $(0, \infty)^n$ .

In contrast, the new diffusion  $\mathfrak{Y}(\cdot)$  of (12.4) lives in  $[0, \infty)^n \setminus \{\mathbf{0}\}$ , and hits the boundary  $\mathcal{O}^n$  of this non-negative orthant with positive probability  $\mathbb{Q}^{\mathbf{x}}[\mathfrak{T} \leq T] = 1 - U(T, \mathbf{x})$  for every  $T \in (0, \infty)$ . The positive

$\mathbb{P}^{\mathbf{x}}$ -supermartingale  $Z(\cdot)X(\cdot)$  is a  $\mathbb{P}^{\mathbf{x}}$ -potential, for every  $\mathbf{x} \in (0, \infty)^n$ . In this case, the three inequalities of (5.5) hold for every  $T \in (0, \infty)$ : the local martingales  $Z(\cdot)$ ,  $Z(\cdot)X(\cdot)$  and  $Z(\cdot)X_i(\cdot)$ ,  $i = 1, \dots, n$  are all strict.

- The model (12.1) can be cast in the form (10.1) for the relative market weights, as a *multivariate Jacobi diffusion process* with dynamics  $d\mu_i(t) = (1 + \zeta)(1 - n\mu_i(t))dt + \sqrt{\mu_i(t)} dW_i(t) - \mu_i(t) \sum_{k=1}^n \sqrt{\mu_k(t)} dW_k(t)$ , or

$$(12.5) \quad d\mu_i(t) = (1 + \zeta)(1 - n\mu_i(t)) dt + \sqrt{\mu_i(t)(1 - \mu_i(t))} dW_i^\sharp(t)$$

with appropriate Brownian motions  $W_1^\sharp(\cdot), \dots, W_n^\sharp(\cdot)$ . Thus, each component  $\mu_i(\cdot)$  is also a diffusion on the unit interval  $(0, 1)$  with local drift  $(1 + \zeta)(1 - ny)$  and local variance  $y(1 - y)$  of Wright-Fisher type. Goia (2009) studies in detail this multivariate diffusion  $\mu(\cdot)$  based on an extension of the Warren-Yor (1999), Gouriéroux-Jasiak (2006) study of skew-products involving Bessel and Jacobi processes.

- From (12.4),  $Y(\cdot) := Y_1(\cdot) + \dots + Y_n(\cdot)$  satisfies the stochastic equation  $dY(t) = Y(t) [dt + dB(t)]$ , where  $B(\cdot) := \sum_{j=1}^n \int_0^\cdot \sqrt{Y_j(t)/Y(t)} dW_j(t)$  is Brownian motion, thus  $Y(\cdot)$  a geometric Brownian motion with drift, under  $\mathbb{Q}^{\mathbf{x}}$ . The process  $\nu(\cdot) = (\nu_1(\cdot), \dots, \nu_n(\cdot))'$  of (10.2) is related to the auxiliary diffusion  $\mathfrak{Y}(\cdot)$  of (12.4) via  $\nu_i(\cdot) = Y_i(\cdot)/Y(\cdot)$ .

The dynamics of these  $\nu_i(\cdot)$ 's are easy to describe in the manner of (9.26), namely,  $d\nu_i(t) = \sqrt{\nu_i(t)} dW_i(t) - \nu_i(t) \sum_{k=1}^n \sqrt{\nu_k(t)} dW_k(t)$ , or in the notation of (12.5):  $d\nu_i(t) = \sqrt{\nu_i(t)(1 - \nu_i(t))} dW_i^\sharp(t)$ . Then the Feller test (e.g. Karatzas & Shreve (1991), pp. 348-350) ensures that each  $\nu_i(\cdot)$  hits one of the endpoints of  $(0, 1)$  in finite expected time. Thus, all but one of the  $Y_i(\cdot)$ 's eventually get absorbed at zero; from that time  $\mathfrak{T}_*$  (with  $\mathbb{E}^{\mathbb{Q}^{\mathbf{x}}}(\mathfrak{T}_*) < \infty$ ) onwards, the only surviving non-zero component  $Y(\cdot)$  behaves like geometric Brownian motion with drift; in particular,  $\mathfrak{Y}(\cdot)$  never hits the origin.

**13. Some Open Questions.** What conditions, if any, on the Markovian covariance structure of section 8 will guarantee that  $\hat{\pi}(\cdot)$  of (11.1) never borrows from the money-market, i.e.,  $\sum_{i=1}^n x_i D_i U(T, \mathbf{x}) \leq 0$ ? That it is a portfolio, i.e., that  $\sum_{i=1}^n x_i D_i U(T, \mathbf{x}) = 0$  holds? (See Remark 6 for a partial answer.) Or better, that  $\hat{\pi}(\cdot)$  of (11.1) is a long-only portfolio, meaning that both this condition and  $D_i (G(T, \mathbf{x}) e^{H(\mathbf{x})}) \geq 0$  hold?

Can an iterative method be constructed, which converges to the minimal solution of the parabolic differential inequality (1.1),  $U(0, \cdot) \equiv 1$  and is numerically implementable (possibly as in Ekström et al. (2008))? How about a Monte-Carlo scheme that computes the quantity  $U(T, \mathbf{x})$  of (9.22)

by generating the paths of the diffusion process  $\mathfrak{Y}(\cdot)$ , then simulating the probability  $\mathbb{Q}^{\mathbf{x}}[\mathfrak{T} > T]$  that  $\mathfrak{Y}(\cdot)$  does not hit the boundary of the non-negative orthant by time  $T$ , when started at  $\mathfrak{Y}(0) = \mathbf{x} \in (0, \infty)^n$ ?

How does  $U(T, \mathbf{x})$  behave as  $T \rightarrow \infty$ ? if it decreases to zero, then at what rate?

## REFERENCES

- BANNER, A. & FERNHOLZ, D. (2008) Short-term arbitrage in volatility-stabilized markets. *Annals of Finance* **4**, 445-454.
- BASS, R. & PERKINS, E. (2003) Degenerate stochastic differential equations with Hölder-continuous coefficients and super-Markov chains. *Trans. Amer. Math. Soc.* **355**, 373-405.
- DELBAEN, F. & SCHACHERMAYER, W. (1995.a) Arbitrage possibilities in Bessel processes and their relations to local martingales. *Probab. Theory Rel. Fields* **102**, 357-366.
- DELBAEN, F. & SCHACHERMAYER, W. (1995.b) The no-arbitrage property under a change of numéraire. *Stochastics & Stochastics Reports* **53**, 213-226.
- EKSTRÖM, E., VON SYDOW, L. & TYSK, J. (2008) Numerical option pricing in the presence of bubbles. *Preprint, Uppsala University*.
- FERNHOLZ, E.R. (2002) *Stochastic Portfolio Theory*. Springer-Verlag, New York.
- FERNHOLZ, E.R. & KARATZAS, I. (2005) Relative arbitrage in volatility-stabilized markets. *Annals of Finance* **1**, 149-177.
- FERNHOLZ, E.R. & KARATZAS, I. (2009) Stochastic Portfolio Theory: A Survey. *Handbook of Numerical Analysis*, volume on *Mathematical Modeling and Numerical Methods in Finance* (A. Bensoussan, ed.), pp. 89-168. Elsevier Publishing Company BV, Amsterdam.
- FERNHOLZ, E.R., KARATZAS, I. & KARDARAS, C. (2005) Diversity and arbitrage in equity markets. *Finance & Stochastics* **31**, 37-53.
- FÖLLMER, H. (1972) The exit measure of a supermartingale. *Zeit. Wahrscheinlichkeitstheorie verw. Gebiete* **21**, 154-166.
- FÖLLMER, H. (1973) On the representation of semimartingales. *Ann. Probab.* **1**, 580-589.
- FÖLLMER, H & GUNDEL, A. (2006) Robust projections in the class of martingale measures. *Illinois Journal of Mathematics* **50**, 439-472.
- FRIEDMAN, A. (2006) *Stochastic Differential Equations & Applications*. Reprint of the 1975/76 edition: Two Volumes Bound as One, Dover Publications, Mineola, NY.
- GOIA, I. (2009) *Bessel and Volatility-Stabilized Processes*. Ph.D. Thesis, Columbia Univ.
- GOURIÉROUX, C. & JASIAK, J. (2006) Multivariate Jacobi process with application to smooth transition. *Journal of Econometrics* **131**, 475-505.
- HEATH, D. & SCHWEIZER, M. (2000) Martingales versus PDEs in finance: an equivalence result with examples. *J. Appl. Probab.* **37**, 947-957.

- HESTON, S.L., LOEWENSTEIN, M. & WILLARD, G.A. (2007) Options and bubbles. *Review of Financial Studies* **20**, 359-390.
- HUGONNIER, J. (2007) Bubbles and multiplicity of equilibria under portfolio constraints. *Preprint, University of Lausanne*.
- HULLEY, H. & PLATEN, E. (2008) Hedging for the long run. *Preprint, University of Technology, Sydney*.
- IKEDA, N. & WATANABE, S. (1989) *Stochastic Differential Equations and Diffusion Processes*. Second Edition. North-Holland, NY and Kodansha Ltd., Tokyo.
- JACOD, J. (1977) A General Representation Theorem for Martingales. *Proceedings of Symposia in Pure Mathematics* **9**, 1-27. American Mathematical Society, Providence, RI.
- JANSON, S. & TYSK, J. (2006) Feynman-Kac formulas for Black-Scholes-type operators. *Bull. London Math. Society* **38**, 269-282.
- KARATZAS, I. & KARDARAS, C. (2007) The numéraire portfolio and arbitrage in semi-martingale markets. *Finance & Stochastics* **11**, 447-493.
- KARATZAS, I. & SHREVE, S.E. (1991) *Brownian Motion and Stochastic Calculus*. Second Edition. Springer-Verlag, NY.
- LOEWENSTEIN, M. & WILLARD, G.A. (2000) Rational equilibrium asset-pricing bubbles in continuous trading models. *Journal of Economic Theory* **91**, 17-58.
- NUALART, D. (1995) *The Malliavin Calculus and Related Topics*. Springer-Verlag, NY.
- PAL, S. & PROTTER, Ph. (2007) Strict local martingales, bubbles, and no early exercise. *Preprint, Cornell University*.
- PARTHASARATHY, K.R. (1967) *Probability Measures on Metric Spaces*. Academic Press, NY.
- RUF, J. (2009) Optimal trading strategies under arbitrage. *Preprint, Columbia University*.
- SIN, C. (1998) Complications with stochastic volatility. *Adv. Appl. Probab.* **30**, 256-268.
- WARREN, J. & YOR, M. (1997) Skew products involving Bessel and Jacobi processes. *Technical Report, Department of Statistics, University of Warwick*.
- WONG, B. & HEYDE, C. (2006) On changes of measure in stochastic volatility problems. *J. Appl. Math. & Stoch. Analysis*, 1-13.
- YOR, M. (2001) *Exponential Functionals of Brownian Motion and Related Processes*. Springer-Verlag, NY.

DEPARTMENT OF COMPUTER SCIENCES  
 UNIVERSITY OF TEXAS AT AUSTIN  
 AUSTIN, TX 78712  
 E-MAIL: fernholz@cs.utexas.edu

INTECH INVESTMENT MANAGEMENT  
 ONE PALMER SQUARE, SUITE 441  
 PRINCETON, NJ 08542  
 E-MAIL: ik@enhanced.com