A remarkable σ-finite measure unifying supremum penalisations for a stable Lévy process

Yuko YANO[(1)(2)]

May 11, 2012

Abstract

The σ-finite measure $\mathcal{P}_{\sup}$ which unifies supremum penalisations for a stable Lévy process is introduced. Silverstein's coinvariant and coharmonic functions for Lévy processes and Chaumont's h-transform processes with respect to these functions are utilized for the construction of $\mathcal{P}_{\sup}$.

Key words

Lévy processes, Stable Lévy processes, Reflected processes,
Penalisation, Path decomposition,
Conditioning to stay negative/positive, Conditioning to hit 0 continuously.

1 Introduction

Roynette-Vallois-Yor ([19] and [20], see also [21] and [22]) have considered the limit laws of Wiener measure weighted by various processes (Γ_t), and they call these studies *Brownian penalisations*. Especially we call the case where the weight process is given by a function of its supremum, i.e., **(S)** $\Gamma_t = f(S_t)$, *supremum penalisation*. Concerning the Brownian supremum penalisations, the authors [20] have obtained the following result. Let $X = ((X_t), (\mathscr{F}_t), \mathbb{W})$ be the canonical representation of a 1-dimensional standard Brownian motion with $\mathbb{W}(X_0 = 0) = 1$ and let $\mathscr{F}_\infty = \sigma(\bigvee_t \mathscr{F}_t)$. Put $S_t = \sup_{s \le t} X_s$. If f is a non-negative Borel function which satisfies

$$\int_0^\infty f(x)\mathrm{d}x = 1, \tag{1.1}$$

then there exists a unique probability law $\mathbb{W}^{(f)}$ on $\mathscr{F}_\infty$ such that

$$\frac{\mathbb{W}[f(S_t)F_s]}{\mathbb{W}[f(S_t)]} \longrightarrow \mathbb{W}^{(f)}[F_s] \quad \text{as} \quad t \to \infty, \tag{1.2}$$

for any fixed $s > 0$ and for any bounded $\mathscr{F}_s$-measurable functional F_s. Moreover the limit measure $\mathbb{W}^{(f)}$ is characterized by

$$\mathbb{W}^{(f)}|_{\mathscr{F}_s} = M_s^{(f)} \cdot \mathbb{W}|_{\mathscr{F}_s}, \tag{1.3}$$

where $(M_s^{(f)}, s \ge 0)$ is a $((\mathscr{F}_s), \mathbb{W})$-martingale which has the form

$$M_s^{(f)} = f(S_s)(S_s - X_s) + \int_{S_s}^\infty f(x)\mathrm{d}x. \tag{1.4}$$

(1) Department of Mathematics, Kyoto Sangyo University. Email: `yyano@cc.kyoto-su.ac.jp`
(2) Supported by Grant-in-Aid for Young Scientists (B) 21740069, 23740073.

We remark that these martingales $(M_s^{(f)})$ which are known as the Azéma-Yor martingales were applied to solve the Skorokhod embedding problem; see [1], [2], also [16] and references therein. In [20] the authors have also obtained the description of the probability measure $\mathbb{W}^{(f)}$ as follows.

Theorem 1.1 (Roynette-Vallois-Yor [20]). *The following holds.*

(i) $\mathbb{W}^{(f)}(S_\infty \in \mathrm{d}x) = f(x)\mathrm{d}x$.

(ii) Let $g = \sup\{t \geq 0 : X_t = S_\infty\}$. *Then* $\mathbb{W}^{(f)}(g < \infty) = 1$ *and, under* $\mathbb{W}^{(f)}$, *we have*

- *(a)* $(X_u, u \leq g)$ *and* $(X_g - X_{g+u}, u \geq 0)$ *are independent;*
- *(b) conditional on* $S_\infty = x$, *the pre-supremum process* $(X_u, u \leq g)$ *is distributed as a Brownian motion starting from* 0 *and stopped at its first hitting time of* x*;*
- *(c) the post-supremum process* $(X_g - X_{g+u}, u \geq 0)$ *is distributed as a* 3*-dimensional Bessel process starting from* 0.

Theorem 1.1 implies that, under the limit measure $\mathbb{W}^{(f)}$, the time g when the process attains its overall supremum is finite, so that the supremum penalisation procedure can be interpreted as looking for probabilities on canonical space, which are close to $\mathbb{W}$, and such that $S_\infty < \infty$ a.s.

Roynette-Vallois-Yor considered Brownian penalisations for many other kinds of weighted processes. For instance, **(L)** $\Gamma_t = f(L_t)$ where L_t denotes the local time of X at the origin, and **(K)** $\Gamma_t = \exp(-\int L(t,x)V(\mathrm{d}x))$ where $L(t,x)$ denotes the local time of X at x; we call the former case *local time penalisation* and the latter case *Kac killing penalisation*. Meanwhile Najnudel-Roynette-Yor [15] have introduced a certain σ-finite measure $\mathcal{W}$ defined as follows:

$$\mathcal{W} = \int_0^\infty \frac{\mathrm{d}u}{\sqrt{2\pi u}} (\Pi^{(u)} \bullet P^{3B}), \tag{1.5}$$

where $\Pi^{(u)}$ denotes the law of Brownian bridge from 0 to 0 of length u and $P^{3B} = (P^{3B,+} + P^{3B,-})/2$ denotes the law of symmetrized 3-dimensional Bessel process; $P^{3B,+}$ is the law of 3-dimensional Bessel process starting from 0, $BES(3)$, whereas $P^{3B,-}$ is the law of $(-BES(3))$. The authors in [15] have shown that the Brownian penalisations including **(S)(L)(K)** can be understood in a unified manner, thanks to this measure $\mathcal{W}$. Especially in the supremum penalisation case, they have shown the following absolute continuity relationship between $\mathcal{W}$ and $\mathbb{W}^{(f)}$:

$$f(S_\infty) \cdot \mathcal{W}^- = \mathbb{W}^{(f)} \quad \text{on} \quad \mathscr{F}_\infty, \tag{1.6}$$

where

$$\begin{aligned} \mathcal{W}^- &= \mathbf{1}_{\{S_\infty < \infty\}} \cdot \mathcal{W} \\ &= \int_0^\infty \frac{\mathrm{d}u}{\sqrt{2\pi u}} \left(\Pi^{(u)} \bullet \frac{P^{3B,-}}{2} \right). \end{aligned} \tag{1.7}$$

As a generalisation of these studies, Yano-Yano-Yor [27] have considered the two kinds of penalisations **(L)** and **(K)** in the case of symmetric α-stable Lévy process with index

$\alpha \in (1,2]$. Let us denote by $((X_t), \mathbb{P})$ such a stable Lévy process with $\mathbb{P}(X_0 = 0) = 1$. The authors have introduced a σ-finite measure $\mathcal{P}$ defined as follows, which is the analogue of $\mathcal{W}$:

$$\mathcal{P} = \int_0^\infty \frac{\Gamma(1/\alpha)}{\alpha\pi} \frac{\mathrm{d}u}{u^{1/\alpha}} (\mathbb{Q}^{(u)} \bullet \mathbb{P}^\times), \tag{1.8}$$

where $\mathbb{Q}^{(u)}$ denotes the law of the stable bridge from 0 to 0 of length u and $\mathbb{P}^\times$ denotes the h-transform process with respect to the harmonic function $|x|^{\alpha-1}$ of the process killed at the first hitting time of 0. We should remark that the process under the measure $\mathbb{P}^\times$ is called *conditioned to avoid* 0, because of the following property obtained by K. Yano [25]: if a functional Z is of the form $Z = f(X_{t_1}, \cdots, X_{t_n})$ for some $0 < t_1 < \cdots < t_n$ and some continuous function $f : \mathbb{R}^n \to \mathbb{R}$ which vanishes at ∞, then one has

$$\mathbb{P}^\times[Z] = \lim_{t\to\infty} \lim_{\varepsilon\to 0+} \mathbb{P}\left[Z \circ \theta_\varepsilon \;\middle|\; \forall u \le t, X_u \circ \theta_\varepsilon \neq 0 \right], \tag{1.9}$$

where $\theta_\cdot$ is the shift operator: $X_u \circ \theta_\cdot = X_{\cdot+u}$. Moreover the following long-time behavior of path under $\mathbb{P}^\times$ is also obtained by K. Yano [26]: if $\alpha \in (1,2)$, then

$$\mathbb{P}^\times \left(\limsup_{t\to\infty} X_t = \limsup_{t\to\infty} (-X_t) = \lim_{t\to\infty} |X_t| = \infty \right) = 1. \tag{1.10}$$

Thus we can see immediately that, under $\mathcal{P}$, $S_\infty = \infty$ a.e. That is, $\mathcal{P}$ cannot unify the supremum penalisations **(S)** in the stable case.

Yano-Yano-Yor [28] have studied the supremum penalisation for a (α, ρ)-stable Lévy process with index $\alpha \in (0,2]$ and positivity parameter $\rho \in (0,1)$. The authors have introduced a generalised Azéma-Yor martingale $(M_s^{(f)})$ which is defined as

$$M_s^{(f)} = f(S_s)(S_s - X_s)^{\alpha\rho} + \alpha\rho \int_{S_s}^\infty f(x)(x - X_s)^{\alpha\rho-1} \mathrm{d}x, \tag{1.11}$$

for any non-negative Borel function f satisfying

$$0 < \int_0^\infty f(x) x^{\alpha\rho-1} \mathrm{d}x < \infty, \tag{1.12}$$

and also introduced the probability measure $\mathbb{P}^{(f)}$ given as

$$\mathbb{P}^{(f)}|_{\mathscr{F}_s} = \frac{M_s^{(f)}}{M_0^{(f)}} \cdot \mathbb{P}|_{\mathscr{F}_s}. \tag{1.13}$$

The authors obtained the following result:

Theorem 1.2 (Yano-Yano-Yor [28]). *Let f be a non-negative function which satisfies either of the following two conditions:*

(i) $f(x) = \mathbf{1}_{\{x \le a\}}$ for some $a > 0$;

(ii) f is absolutely continuous with respect to the Lebesgue measure and satisfies

$$\lim_{x\to\infty} f(x) = 0 \quad \textit{and} \quad 0 < \int_0^\infty |f'(x)| x^{\alpha\rho} \mathrm{d}x < \infty. \tag{1.14}$$

Then it holds that, for any $s > 0$ and any bounded $\mathscr{F}_s$-measurable functional F_s,

$$\frac{\mathbb{P}[f(S_t)F_s]}{\mathbb{P}[f(S_t)]} \longrightarrow \mathbb{P}^{(f)}[F_s] \quad as \quad t \to \infty. \tag{1.15}$$

We remark that the condition (ii) in Theorem 1.2 is stronger than the condition (1.12) because we have

$$\begin{aligned}\int_0^\infty f'(x)x^{\alpha\rho}\mathrm{d}x &= \alpha\rho\int_0^\infty f'(x)\mathrm{d}x\int_0^x y^{\alpha\rho-1}\mathrm{d}y \\ &= \alpha\rho\int_0^\infty y^{\alpha\rho-1}\mathrm{d}y\int_y^\infty f'(x)\mathrm{d}x \\ &= -\alpha\rho\int_0^\infty f(y)y^{\alpha\rho-1}\mathrm{d}y.\end{aligned}$$

One may conjecture that the assumption of Theorem 1.2 can be weakened to the condition (1.12) that is sufficient to define the generalised Azéma-Yor martingale and the measure $\mathbb{P}^{(f)}$; however, this is still an open problem.

In the present paper we introduce a certain σ-finite measure $\mathcal{P}_{\sup}$ by using Chaumont's h-transform processes for Lévy processes (cf. Theorem 5.1 below):

$$\mathcal{P}_{\sup} = \int_0^\infty \mathrm{d}x\psi(x)(\mathbb{P}_{0\nearrow x} \bullet \mathbb{P}_{x\downarrow x}),$$

where ψ is the function stated below in (2.10), $\mathbb{P}_{0\nearrow x}$ denotes the law of the process starting from 0 and conditioned to hit x continuously (in fact, under $\mathbb{P}_{0\nearrow x}$, the process starting from 0 is killed at the first hitting time at x), and $\mathbb{P}_{x\downarrow x}$ denotes the law of the process starting from x and conditioned to stay below level x. $\mathcal{P}_{\sup}$ is another analogue of $\mathcal{W}$ and $\mathcal{P}$, and it is a generalisation of $\mathcal{W}^-$ given in (1.7). We remark that, in the Brownian case, $\mathcal{P}_{\sup}^{\mathrm{BM}}$ is given by the following:

$$\begin{aligned}\mathcal{P}_{\sup}^{\mathrm{BM}} &= \int_0^\infty \mathrm{d}x\left(\mathbb{W}_{0\nearrow x} \bullet P_x^{3B,-}\right) \\ &= \int_0^\infty \mathrm{d}x\int_0^\infty \mathrm{d}u\frac{x}{\sqrt{2\pi u^3}}\mathrm{e}^{-\frac{x^2}{2u}}\left(\mathbb{W}_{0\nearrow x}^{(u)} \bullet P_x^{3B,-}\right),\end{aligned} \tag{1.16}$$

where $\mathbb{W}_{0\nearrow x}$ denotes the law of Brownian motion killed at the first hitting time at x and $\mathbb{W}_{0\nearrow x}^{(u)}(\cdot) = \mathbb{W}_{0\nearrow x}(\cdot|T_{\{x\}} = u)$, and $P_x^{3B,-}$ denotes the law of the translation by x of $(-BES(3))$. The latter equality is obtained from the well-known fact (see, e.g., [11]) that

$$\mathbb{W}(T_{\{x\}} \in \mathrm{d}u) = \mathrm{d}u\frac{x}{\sqrt{2\pi u^3}}\mathrm{e}^{-\frac{x^2}{2u}}. \tag{1.17}$$

We note that the measure $\mathcal{P}_{\sup}^{\mathrm{BM}}$ equals $\mathcal{W}^-$ by the agreement formula obtained by Pitman-Yor [17].



Figure 1. Sample path of $\Pi^{(u)} \bullet P^{3B,-}$



Figure 2. Sample path of $\mathbb{W}^{(u)}_{0\nearrow x} \bullet P^{3B,-}_x$

We then show that the measure $\mathcal{P}_{\sup}$ unifies the supremum penalisations. More precisely, we shall define a probability measure $\mathbb{P}^{(f)}$ as the transformation of the law $\mathbb{P}$ of a Lévy process by the generalised Azéma-Yor martingale defined as (6.2) below. This measure $\mathbb{P}^{(f)}$ is the generalisation of (1.13) for a general Lévy process. We then prove the absolute continuity relationship between $\mathcal{P}_{\sup}$ and $\mathbb{P}^{(f)}$ in the Lévy case, which is the analogue of (1.6) (cf. Theorem 7.3 below):

$$\frac{f(S_\infty)\cdot \mathcal{P}_{\sup}}{\mathcal{P}_{\sup}[f(S_\infty)]} = \mathbb{P}^{(f)} \quad \text{on} \quad \mathscr{F}_\infty.$$

We obtain a detailed description of $\mathbb{P}^{(f)}$ as a consequence of this result (cf. Theorem 7.5 below):

$$\mathbb{P}^{(f)} = \int_0^\infty \mathbb{P}^{(f)}(S_\infty \in \mathrm{d}x)(\mathbb{P}_{0\nearrow x} \bullet \mathbb{P}_{x\downarrow x}).$$

To prove the absolute continuity relationship between $\mathcal{P}_{\sup}$ and $\mathbb{P}^{(f)}$, we shall introduce a path decomposition of the law $\mathbb{P}$ of a Lévy process up to a fixed time t with respect to the position and the time where the process attains its supremum before time t.

The organization of the present paper is as follows. In Section 2 and 3, we recall some preliminary facts about Lévy processes and (α, ρ)-stable Lévy processes, respectively. If a reader needs to see details, he/she may refer to, e.g., [3], [10], [12] and [23]. In Section 4, we review Chaumont's two kinds of h-transform processes for a Lévy process. In Section 5, we establish a path decomposition of the law of a Lévy process at the position and the time where the Lévy process attains its supremum up to a fixed time t. In Section 6, we introduce the generalised Azéma-Yor martingale in the general Lévy case, which is the generalisation of (1.4) and (1.11). A certain probability measure which should appear as the limit measure of the supremum penalisation is also introduced in this section. In Section 7, we introduce the σ-finite measure $\mathcal{P}_{\sup}$ which unifies the supremum penalisations and give some properties of the measure $\mathcal{P}_{\sup}$. In Section 8, we compare $\mathcal{P}_{\sup}$ with $\mathcal{P}$ and give some remarks on these measures.

2 Preliminaries about Lévy processes

Let $\mathscr{D}([0,\infty))$ be the space of càdlàg paths $\omega : [0,\infty) \to \mathbb{R} \cup \{\delta\}$ with lifetime $\zeta(\omega) = \inf\{s : \omega(s) = \delta\}$ where δ is a cemetery point. Let (X_t) denote the coordinate process, $X_t(\omega) = \omega_t$, and let $(\mathscr{F}_t)$ denote its natural filtration with $\mathscr{F}_\infty = \vee_{t\geq 0}\mathscr{F}_t$. Let $\mathbb{P}$ be the law of a Lévy process $X = (X_t, t \geq 0)$ with $\mathbb{P}(X_0 = 0) = 1$ such that

$$\mathbb{P}\left[\exp\{i\lambda X_t\}\right] = \mathrm{e}^{-t\Psi(\lambda)}, \qquad t \geq 0,\ \lambda \in \mathbb{R}, \tag{2.1}$$

where

$$\Psi(\lambda) = i\gamma\lambda + \frac{\sigma^2\lambda^2}{2} + \int_{\mathbb{R}\setminus\{0\}} \left(1 - \mathrm{e}^{i\lambda x} + i\lambda x 1_{\{|x|<1\}}\right) \nu(\mathrm{d}x) \tag{2.2}$$

for some constants γ, σ, and Lévy measure ν on $\mathbb{R} \setminus \{0\}$ which satisfies

$$\int_{\mathbb{R}\setminus\{0\}} (x^2 \wedge 1)\nu(\mathrm{d}x) < \infty. \tag{2.3}$$

We denote by $\mathbb{P}_x$ the law of $X + x$ under $\mathbb{P}$ for every $x \in \mathbb{R}$. Throughout this paper we assume the following absolute continuity condition **(A1)**:

(A1) For each $\alpha > 0$, there exists an integrable function u_α such that

$$\mathbb{P}_x\left[\int_0^\infty \mathrm{e}^{-\alpha t} f(X_t)\mathrm{d}t\right] = \int_{-\infty}^\infty u_\alpha(y) f(x+y)\mathrm{d}y, \tag{2.4}$$

for every non-negative Borel function f.

Let S_t and I_t be respectively the supremum and the infimum processes up to time t, that is, for all $t < \zeta(\omega)$,

$$S_t = \sup\{X_s : 0 \leq s \leq t\} \qquad \text{and} \qquad I_t = \inf\{X_s : 0 \leq s \leq t\}. \tag{2.5}$$

Let T_A denote the first entrance time of a Borel set $A \subset \mathbb{R}$ of X, i.e.,

$$T_A = \inf\{s > 0 : X_s \in A\}. \tag{2.6}$$

Define

$$R = S - X. \tag{2.7}$$

The process $R = (R_t, t \geq 0)$ is called *the reflected process of X at the supremum.* We recall that R is a strong Markov process (Bingham [5], see also [4]). We consider the following condition **(A2)**:

(A2) 0 is regular for $(0, \infty)$ with respect to X under $\mathbb{P}$, i.e., $\mathbb{P}(T_{(0,\infty)} = 0) = 1$.

Then 0 is regular for itself with respect to R, and hence we can define a local time $L = (L_t, t \geq 0)$ at level 0 of R. We denote by τ the right-continuous inverse of L and let $H = X(\tau) = S(\tau)$. We recall that the pair (τ, H) is a bivariate subordinator, called the (upwards) ladder process, in particular, τ and H are separately also subordinators, called the (upwards) ladder time and the (upwards) ladder height process, respectively. Denote by X^* the dual process of X, i.e., $X^* = -X$. Consider

(A2*) 0 is regular for $(-\infty, 0)$ with respect to X under $\mathbb{P}$.

Then we can define a local time L^* at level 0 of $R^* = S^* - X^* = X - I$, and also get the (downwards) ladder time τ^* and the (downwards) ladder height time H^* of R^*.

We denote by E the set of càdlàg paths $e : [0, \infty) \to \mathbb{R} \cup \{\delta\}$ such that

$$e(t) \begin{cases} \in \mathbb{R} \setminus \{0\}, & 0 < t < \zeta_e; \\ = \delta, & t \geq \zeta_e, \end{cases}$$

where

$$\zeta_e = \inf\{t > 0 : e(t) = \delta\}. \tag{2.8}$$

We call E the set of excursions and an element $e \in E$ an excursion path. For $e \in E$, we call ζ_e the lifetime of the excursion e. Set $D = \{l : \tau_l - \tau_{l-} > 0\}$. For each $l \in D$, we set

$$e_l(t) = \begin{cases} R_{t+\tau_{l-}}, & 0 \leq t < \tau_l - \tau_{l-}; \\ \delta, & t \geq \tau_l - \tau_{l-}. \end{cases}$$

By Itô's theorem, the point process $(e_l, l \in D)$ which takes values on E is a Poisson point process, and its characteristic measure $\boldsymbol{n}$ is called *the Itô measure of excursions.* Similarly, we can introduce excursions e^* with respect to R^* and denote by $\boldsymbol{n}^*$ its Ito measure.

We recall the following important formula, see also p. 7 in [4], and Proposition (1.10) in Chapter XII in [18]. Denote by $\mathscr{P}(\mathscr{F}_t)$ the predictable σ-field relative to $(\mathscr{F}_t)$ (cf. p. 47 in [18]), and let $\mathscr{E} = \sigma\{e(t)\}$.

Theorem 2.1 (Compensation formula). *Let $F = F(t, \omega, e)$ be a positive process defined on $[0, \infty) \times \mathscr{D} \times E$, measurable with respect to $\mathscr{P}(\mathscr{F}_t) \otimes \mathscr{E}$ and vanishing at δ. Then one has*

$$\mathbb{P}\left[\sum_{l \in D} F(\tau_{l-}, X, e_l)\right] = \mathbb{P} \otimes \widehat{\boldsymbol{n}}\left[\int_0^\infty \mathrm{d}L_t F(t, X, \widehat{X})\right], \tag{2.9}$$

where the symbol $\widehat{\ }$ means independence.

Under **(A1)** and **(A2)**, there exists a unique coexcessive function ψ for the killed process, i.e., $\mathbb{P}_{-x}[\psi(X_t^*)\mathbf{1}_{\{t<T_{(0,\infty)}\}}] \leq \psi(x)$ for $x \geq 0$, which satisfies

$$\int_0^\infty \psi(y)f(y)\mathrm{d}y = \mathbb{P}\left[\int_0^\infty f(S_{\tau_s})\mathrm{d}s\right] = \mathbb{P}\left[\int_0^\infty f(S_t)\mathrm{d}L_t\right], \tag{2.10}$$

for any non-negative Borel function f on $[0,\infty)$. We remark that ψ is continuous and satisfies that $0 < \psi(x) < \infty$ for $x \in (0,\infty)$. Thanks to Silverstein [24], the function ψ is *coharmonic* on $(0,\infty)$, that is,

$$\mathbb{P}_{-x}\left[\psi(X_{T_M}^*)\mathbf{1}_{\{T_M<T_{(0,\infty)}\}}\right] = \psi(x), \qquad x > 0, \tag{2.11}$$

where M denotes a subinterval of $(-\infty,0)$ whose complement $(-\infty,0)\setminus M$ is open and has compact closure. We assume further that

(A3) $\mathbb{P}_x(T_{(-\infty,0)} < \infty) = 1$ for $x > 0$.

Note that **(A3)** is equivalent to that $I_\infty = -\infty$ $\mathbb{P}$-a.s. Then the function h given by

$$h(x) = \int_0^x \psi(y)\mathrm{d}y = \mathbb{P}\left[\int_0^\infty \mathbf{1}_{\{S_t\leq x\}}\mathrm{d}L_t\right] \tag{2.12}$$

is *coinvariant* by Silverstein [24], that is,

$$\mathbb{P}_{-x}\left[h(X_t^*)\mathbf{1}_{\{t<T_{(0,\infty)}\}}\right] = h(x), \qquad x > 0. \tag{2.13}$$

We remark that the function h is finite, continuous, increasing, and that $h(0) = 0$. We remark that every positive coinvariant function is also coharmonic.

Similarly, under **(A1)** and **(A2*)**, there exists a version of the potential density of the subordinator $(I_{\tau_s^*})_{s\geq 0}$. That is, there exists a unique coexcessive function ψ^* for the killed process, i.e., $\mathbb{P}_x[\psi^*(X_t)\mathbf{1}_{\{t<T_{(-\infty,0)}\}}] \leq \psi^*(x)$ for $x \geq 0$, which satisfies

$$\int_0^\infty \psi^*(y)f(y)\mathrm{d}y = \mathbb{P}\left[\int_0^\infty f(I_{\tau_s^*})\mathrm{d}s\right] = \mathbb{P}\left[\int_0^\infty f(I_t)\mathrm{d}L_t^*\right], \tag{2.14}$$

for any non-negative Borel function f on $(0,\infty)$. Also thanks to Silverstein [24], the function ψ^* is coharmonic on $(0,\infty)$, that is,

$$\mathbb{P}_x\left[\psi^*(X_{T_{M'}})\mathbf{1}_{\{T_{M'}<T_{(-\infty,0)}\}}\right] = \psi^*(x), \qquad x > 0, \tag{2.15}$$

where M' denotes a subinterval of $(0,\infty)$ whose complement $(0,\infty)\setminus M'$ is open and has the compact closure. If we assume further that

(A3*) $\mathbb{P}_{-x}(T_{(0,\infty)} < \infty) = 1$ for $x > 0$.

Note that **(A3*)** is equivalent to that $S_\infty = \infty$ $\mathbb{P}$-a.s. Then the function h^* given by

$$h^*(x) = \int_0^x \psi^*(y)\mathrm{d}y = \mathbb{P}\left[\int_0^\infty \mathbf{1}_{\{I_t\leq x\}}\mathrm{d}L_t^*\right] \tag{2.16}$$

is coinvariant, that is,

$$\mathbb{P}_x\left[h^*(X_t)\mathbf{1}_{\{t<T_{(-\infty,0)}\}}\right] = h^*(x), \qquad x > 0. \tag{2.17}$$

3 Preliminaries about (α, ρ)-stable Lévy processes

Consider a probability measure $\mathbb{P}$ on $\mathscr{D}([0,\infty))$ with respect to which X is a strictly stable Lévy process of index $\alpha \in (0,2]$ with $\mathbb{P}(X_0 = 0) = 1$. That is,

$$\mathbb{P}[\mathrm{e}^{i\lambda X_t}] = \mathrm{e}^{-t\Psi(\lambda)}, \qquad t \geq 0,\ \lambda \in \mathbb{R}, \tag{3.1}$$

where

$$\Psi(\lambda) = \begin{cases} c|\lambda|^\alpha \left(1 - i\beta \mathrm{sgn}(\lambda) \tan \dfrac{\pi\alpha}{2}\right), & \alpha \in (0,1) \cup (1,2), \\ c|\lambda| + di\lambda, & \alpha = 1, \\ c\lambda^2, & \alpha = 2, \end{cases} \tag{3.2}$$

for some constants $c > 0$, $d \in (-\infty, \infty)$ and $\beta \in [-1,1]$. The Lévy measure ν is given by

$$\nu(\mathrm{d}x) = \begin{cases} (c_+ \mathbf{1}_{\{x>0\}} + c_- \mathbf{1}_{\{x<0\}})|x|^{-\alpha-1}\mathrm{d}x, & \alpha \in (0,1) \cup (1,2), \\ \widetilde{c}|x|^{-2}\mathrm{d}x, & \alpha = 1, \\ 0, & \alpha = 2, \end{cases} \tag{3.3}$$

where $\beta = (c_+ - c_-)/(c_+ + c_-)$, and for some constant $\widetilde{c} > 0$. When $c_{+[-]} = 0$, the process is spectrally negative[positive] (or, has no positive[negative] jumps). We remark that the condition **(A1)** is also valid in the stable Lévy case because of the scaling property of X.

Put $\rho = \mathbb{P}(X_t \geq 0)$. By the scaling property of X, ρ does not depend on $t > 0$. We call ρ *the positivity parameter*. It is well-known that the value of ρ for $\alpha \neq 1, 2$ can be represented in terms of the parameter β as

$$\rho = \frac{1}{2} + \frac{1}{\pi\alpha} \arctan\left(\beta \tan \frac{\pi\alpha}{2}\right). \tag{3.4}$$

See Section 2.6 in [29], and p. 218 in [3]. The range of the value of ρ is classified as follows:

$$\rho \begin{cases} \in [0,1] \quad \text{if} \quad \alpha \in (0,1) \\ \quad (\text{when } \rho = 0 \text{ or } 1, \text{the process is a subordinator or a negative subordinator}), \\ \in (0,1) \quad \text{if} \quad \alpha = 1, \\ \in [1 - 1/\alpha, 1/\alpha] \quad \text{if} \quad \alpha \in (1,2) \\ \quad (\text{when } \rho = 1 - 1/\alpha \text{ or } 1/\alpha, \text{ the process is spectrally positive or spectrally negative}), \\ = 1/2 \quad \text{if} \quad \alpha = 2. \end{cases}$$

Assume that

(B) $\rho \in (0,1)$.

Note that **(B)** is equivalent to that $|X|$ is not a subordinator. Then $\alpha\rho \in (0,1]$. We note that the condition **(B)** for the stable Lévy case implies the conditions **(A2)** and **(A2*)**, that is, 0 is regular for both $(0,\infty)$ and $(-\infty,0)$ with respect to X. Therefore we can define the local times L, L^*, etc. for the reflected and dual reflected processes in this case. Moreover the condition **(B)** also implies the conditions **(A3)** and **(A3*)**: More precisely, when $\alpha \in (1,2]$, **(A3)** and **(A3*)** hold since X is strictly stable; when $\alpha \in (0,1]$, they hold because of the condition **(B)**.

Assuming **(B)**, the function h defined in (2.12) is

$$h(x) = Cx^{\alpha\rho}, \qquad x > 0 \tag{3.5}$$

for some constant $C > 0$. This is obtained from the fact that the ladder time process τ is a stable subordinator of index ρ and the ladder height process H is a stable process of index $\alpha\rho$ (see Lemma VIII 1 in [4]). Furthermore, in this case, we have

$$\psi(x) = C\alpha\rho x^{\alpha\rho-1}, \qquad x > 0. \tag{3.6}$$

Similarly, we have

$$h^*(x) = Dx^{\alpha(1-\rho)} \qquad \text{and} \qquad \psi^*(x) = D\alpha(1-\rho)x^{\alpha(1-\rho)-1}, \qquad x > 0 \tag{3.7}$$

for some constant $D > 0$. These constants C and D may depend upon the choice of the local time L and L^*, respectively.

Example 3.1 (Brownian case). When $\alpha = 2$ and $\rho = 1/2$, X is a 1-dimensional Brownian motion up to a multiplicative constant. In this case we have

$$h(x) = x \qquad \text{and} \qquad \psi(x) = 1, \qquad x > 0. \tag{3.8}$$

4 Chaumont's two kinds of conditionings for a Lévy process

In this section we shall review two kinds of conditionings for a Lévy process introduced by Chaumont [7, 6], which are obtained by Doob's h-transform.

Let $X = ((X_t), \mathbb{P})$ be a Lévy process with the conditions **(A1)**, **(A2)** and **(A3)**. The functions ψ and h are stated as (2.10) and (2.12), respectively.

1° The process conditioned to stay negative.

For non-negative $\mathscr{F}_t$-measurable functional F_t, define $(\mathbb{P}_{-x\downarrow 0}, x > 0)$ as

$$\mathbb{P}_{-x\downarrow 0}[F_t(X)] := \frac{1}{h(x)}\mathbb{P}_{-x}\left[h(X_t^*)\mathbf{1}_{\{t<T_{(0,\infty)}\}}F_t(X)\right], \qquad x > 0. \tag{4.1}$$

The family $(\mathbb{P}_{-x\downarrow 0}|_{\mathscr{F}_t}, t \geq 0)$ is proved to be consistent by the coinvariance of the function h and hence $\mathbb{P}_{-x\downarrow 0}$ is well-defined as a probability measure on $\mathscr{F}_\infty$. The process $(X, \mathbb{P}_{-x\downarrow 0})$ is called *the process starting from* $(-x)$ *and conditioned to stay negative* since it has the following property:

Theorem 4.1 ([6, Theorem 1]). *Let* $\boldsymbol{e}$ *be an independent exponential random variable with index* 1*. Then, for any* $x > 0, t \geq 0$ *and any* $\mathscr{F}_t$*-measurable functional* F_t*, it holds that*

$$\lim_{\varepsilon\to 0}\mathbb{P}_{-x}\left[\mathbf{1}_{\{t<\boldsymbol{e}/\varepsilon\}}F_t \;\middle|\; X_s < 0,\ 0 \leq s \leq \boldsymbol{e}/\varepsilon\right] = \mathbb{P}_{-x\downarrow 0}[F_t]. \tag{4.2}$$

It is proved by Chaumont [6] and Chaumont-Doney [9] that $\mathbb{P}_{-x\downarrow 0}$ converges in the Skorokhod sense to $\mathbb{P}_{0\downarrow 0}$ as $x \to 0$. Thus it follows from Theorem 4.1 that, for every $x \geq 0$,

$$\mathbb{P}_{-x\downarrow 0}(X_0 = -x;\ \zeta = \infty;\ X_t < 0 \text{ for all } t > 0;\ \lim_{t\to\infty} X_t = -\infty) = 1. \tag{4.3}$$

Here ζ denotes the lifetime.

Chaumont [6] also showed the absolutely continuity between $\mathbb{P}_{0\downarrow 0}$ and the excursion measure $\boldsymbol{n}$ of the reflected process $R = S - X$ as follows:

Theorem 4.2 ([6, Theorem 3]). *It holds*

$$\mathbb{P}_{0\downarrow 0}[F_t(X)] = \boldsymbol{n}\left[h(X_t)\mathbf{1}_{\{t<\zeta_e\}}F_t(X^*)\right], \tag{4.4}$$

for non-negative $\mathscr{F}_t$-measurable functional F_t.

For $b \le a$, denote by $\mathbb{P}_{b\downarrow a}$ the law of $X + a$ under $\mathbb{P}_{b-a\downarrow 0}$, that is, $(X, \mathbb{P}_{b\downarrow a})$ is the process starting from b and conditioned to stay below level a.

2° The process conditioned to hit 0 continuously.

Define $(\mathbb{P}_{-x\nearrow 0}, x > 0)$ as

$$\mathbb{P}_{-x\nearrow 0}\left[\mathbf{1}_{\{t<\zeta\}}F_t(X)\right] := \frac{1}{\psi(x)}\mathbb{P}_{-x}\left[\psi(X_t^*)\mathbf{1}_{\{t<T_{(0,\infty)}\}}F_t(X)\right], \tag{4.5}$$

for non-negative $\mathscr{F}_t$-measurable functional F_t. The process $(X, \mathbb{P}_{-x\nearrow 0})$ is called *the process starting from* $(-x)$ *and conditioned to hit* 0 *continuously*, or also called *the process conditioned to die at* 0, and has the following property:

Theorem 4.3 ([6, Proposition 2]). *For $x > 0$, it holds that*

$$\mathbb{P}_{-x\nearrow 0}(X_0 = -x;\ \zeta < \infty;\ X_t < 0 \text{ for all } t < \zeta;\ X_{\zeta-} = 0) = 1, \tag{4.6}$$

where ζ denotes the lifetime.

The following result is also shown by Chaumont [6]:

Theorem 4.4 ([6, Proposition 3]). *For any $x > 0, k > 0, t \ge 0$ and any $\mathscr{F}_t$-measurable functional F_t,*

$$\lim_{\varepsilon\to 0}\mathbb{P}_{-x}\left[\mathbf{1}_{\{t<T_{(-k,\infty)}\}}F_t \,\middle|\, S_{T_{(0,\infty)}-} \ge -\varepsilon\right] = \mathbb{P}_{-x\nearrow 0}\left[\mathbf{1}_{\{t<T_{(-k,0)}\}}F_t\right]. \tag{4.7}$$

Denote by $\mathbb{P}_{0\nearrow x}$ the law of $X + x$ under $\mathbb{P}_{-x\nearrow 0}$, that is, $(X, \mathbb{P}_{0\nearrow x})$ is the process starting from 0 and conditioned to hit x continuously. For later use, we rewrite (4.5) by translation to obtain

$$\mathbb{P}_{0\nearrow x}\left[\mathbf{1}_{\{t<\zeta\}}F_t(X)\right] = \frac{1}{\psi(x)}\mathbb{P}\left[\psi(x - X_t)\mathbf{1}_{\{t<T_{(x,\infty)}\}}F_t(X)\right], \tag{4.8}$$

since we have

$$\begin{aligned}
\mathbb{P}_{0\nearrow x}\left[\mathbf{1}_{\{t<\zeta\}}F_t(X)\right] &= \mathbb{P}_{-x\nearrow 0}\left[\mathbf{1}_{\{t<\zeta\}}F_t(X + x)\right] \\
&= \frac{1}{\psi(x)}\mathbb{P}_{-x}\left[\psi(X_t^*)\mathbf{1}_{\{t<T_{(0,\infty)}\}}F_t(X + x)\right] \\
&= \frac{1}{\psi(x)}\mathbb{P}\left[\psi(x + X_t^*)\mathbf{1}_{\{t<T_{(x,\infty)}\}}F_t(X)\right].
\end{aligned}$$

5 Path decomposition at the position and the time where the Lévy process attains its supremum up to time t

Our aim in this section is to prove Theorem 5.1, which consists of a path decomposition with respect to the position and the time where the Lévy process attains its supremum up to time $t > 0$.

Let us denote by $X^{(u)}$ the coordinate process considered up to time u, i.e.,

$$X_t^{(u)} = \begin{cases} X_t, & t < u; \\ \delta, & t \geq u, \end{cases}$$

and denote by $\mathbb{P}_x^{(u)}$ the law of $X^{(u)}$ under $\mathbb{P}_x$. We denote the concatenation between two independent processes $X^{(u)}$ and $\widehat{X}^{(v)}$ by $X^{(u)} \bullet \widehat{X}^{(v)}$, i.e.,

$$(X^{(u)} \bullet \widehat{X}^{(v)})_t = \begin{cases} X_t^{(u)}, & 0 \leq t < u; \\ \widehat{X}_{t-u}^{(v)}, & u \leq t < u+v; \\ \delta, & t \geq u+v. \end{cases}$$

We define the measure $\mathbb{P}_x^{(u)} \bullet \mathbb{P}_y^{(v)}$ as the law of the concatenation $X^{(u)} \bullet \widehat{X}^{(v)}$ between two independent processes $X^{(u)}$ and $\widehat{X}^{(v)}$ where $(X^{(u)}, \widehat{X}^{(v)})$ is considered under the product measure $\mathbb{P}_x^{(u)} \otimes \widehat{\mathbb{P}}_y^{(v)}$.

For $t > 0$, we denote the last time when the process attains its supremum before t by

$$g_t = \sup\{s \leq t : X_s = S_s\}, \tag{5.1}$$

with the convention $\sup \emptyset = 0$.

Theorem 5.1. *Let $X = ((X_t), \mathbb{P})$ be a Lévy process with $\mathbb{P}(X_0 = 0) = 1$ and assume* **(A1)**, *as well as both* **(A2)** *and* **(A2*)**. *Let $F_t(X^{(t)}) = F(t, X_{t\wedge\cdot})$. Then it holds that*

$$\mathbb{P}\left[F_t(X^{(t)})\right] = \int \rho_t(\mathrm{d}x\mathrm{d}u) \left(\mathbb{P}_{0\nearrow x}^{(u)} \bullet \mathbb{M}_x^{(t-u)}\right) \left[F_t(X^{(t)})\right], \tag{5.2}$$

where the integral is taken over $[0, \infty) \times [0, t)$ and

$$\rho_t(\mathrm{d}x\mathrm{d}u) = \mathrm{d}x\psi(x)\mathbb{P}_{0\nearrow x}(\zeta \in \mathrm{d}u)\boldsymbol{n}(\zeta_e > t-u); \tag{5.3}$$

$$\mathbb{P}_{0\nearrow x}^{(u)}(\cdot) = \mathbb{P}_{0\nearrow x}(\cdot|\zeta = u) \quad (\zeta \text{ denotes the lifetime}); \tag{5.4}$$

$$\mathbb{M}_x^{(s)}[F(X)] = \frac{\boldsymbol{n}\left[F(x - X^{(s)}); \zeta_e > s\right]}{\boldsymbol{n}(\zeta_e > s)} \quad (\zeta_e \text{ denotes the lifetime}). \tag{5.5}$$

In other words, the following statements hold:

(i) $\rho_t(\mathrm{d}x\mathrm{d}u)$ gives the joint distribution of S_t and g_t, i.e.,

$$\rho_t(\mathrm{d}x\mathrm{d}u) = \mathbb{P}(S_t \in \mathrm{d}x, g_t \in \mathrm{d}u); \tag{5.6}$$

(ii) given $g_t = u$, the pre-supremum process $(X_s, s \leq u)$ and the post-supremum process $(X_u - X_{u+s}, 0 \leq s \leq t-u)$ are independent under $\mathbb{P}$;

(iii) given $S_t = x$ and $g_t = u$, $(X_s, s \le u)$ under $\mathbb{P}$ is distributed as $\mathbb{P}^{(u)}_{0\nearrow x}$; the process conditioned to hit x continuously, with duration u;

(iv) given $S_t = x$ and $g_t = u$, $(x - X_{u+s}, 0 \le s \le t-u)$ under $\mathbb{P}$ is distributed as the meander $\mathbb{M}^{(t-u)} := \mathbb{M}^{(t-u)}_0$.

Remark 5.2. The fact (ii) in Theorem 5.1 is well-known and can be found in Lemma VI 6 in [4].

Remark 5.3. We can also see that $X_{g_t} = X_{g_t-}$, that is, the process does not jump at g_t; the last hitting time of its supremum up to time t. This fact is guaranteed by the conditions **(A2)** and **(A2*)**, see also [4], p. 160.

Remark 5.4. Theorem 5.1 is obtained independently by Chaumont [8] in his recent work for some purpose different from ours.

Before the proof of Theorem 5.1, we recall the following lemma from Chaumont [7]:

Lemma 5.5 ([7, Lemma 3]). *Let $X = ((X_t), \mathbb{P})$ be a Lévy process with $\mathbb{P}(X_0 = 0) = 1$ satisfying conditions* **(A1)**, **(A2)** *and* **(A2*)**. *Denote by L the local time at 0 of the reflected process $R = S - X$. Let H be a predictable functional. Then it holds that*

$$\mathbb{P}\left[\int_0^\infty H_t(X)\mathrm{d}L_t\right] = \int_0^\infty \mathbb{P}_{-x\nearrow 0}[H_\zeta(X+x)]\psi(x)\mathrm{d}x. \tag{5.7}$$

The proof of Lemma 5.5 for the stable Lévy process is given in [7]. Lemma 5.5 for the general Lévy process is proved in the same way, so we omit the proof.

Proof of Theorem 5.1. We have

$$\int_0^\infty \mathrm{d}t F_t(X^{(t)}) = \sum_{l\in D}\int_0^{\zeta(e_l)} F_{\tau_l-+r}(X^{(\tau_l-)} \bullet (X_{\tau_l-} - e_l))\mathrm{d}r. \tag{5.8}$$

Hence we have

$$\begin{aligned}\mathbb{P}\left[\int_0^\infty \mathrm{d}t F_t(X^{(t)})\right] &= \mathbb{P}\left[\sum_{l\in D}\int_0^{\zeta(e_l)} F_{\tau_l-+r}\left(X^{(\tau_l-)} \bullet (X_{\tau_l-} - e_l)\right)\mathrm{d}r\right] \\ &= \mathbb{P}\otimes\widehat{\boldsymbol{n}}\left[\int_0^\infty \mathrm{d}L_s \int_0^{\widehat{\zeta}_e} F_{s+r}(X^{(s)} \bullet (X_s - \widehat{X}^{(r)}))\mathrm{d}r\right],\end{aligned} \tag{5.9}$$

by the compensation formula (Theorem 2.1). By Lemma 5.5, we have

$$\begin{aligned}(5.9) &= \int_0^\infty \mathrm{d}x\psi(x)\left(\mathbb{P}_{-x\nearrow 0}\otimes\widehat{\boldsymbol{n}}\right)\left[\int_0^\infty F_{\zeta+r}\left((X^{(\zeta)}+x)\bullet(X_\zeta + x - \widehat{X}^{(r)})\right)\mathbf{1}_{\{r<\widehat{\zeta}_e\}}\mathrm{d}r\right] \\ &= \int_0^\infty \mathrm{d}x\psi(x)\left(\mathbb{P}_{-x\nearrow 0}\otimes\widehat{\boldsymbol{n}}\right)\left[\int_0^\infty F_{\zeta+r}\left((X^{(\zeta)}+x)\bullet(x - \widehat{X}^{(r)})\right)\mathbf{1}_{\{r<\widehat{\zeta}_e\}}\mathrm{d}r\right].\end{aligned} \tag{5.10}$$

Here we use the fact that $X_\zeta = 0$. By translation by x of $\mathbb{P}_{-x\nearrow 0}$ and then changing of variable $\zeta + r = u$, we have

$$(5.10) = \int_0^\infty \mathrm{d}x\psi(x)\,(\mathbb{P}_{0\nearrow x}\otimes\widehat{\boldsymbol{n}})\left[\int_0^\infty F_{\zeta+r}\left(X^{(\zeta)}\bullet(x-\widehat{X}^{(r)})\right)\mathbf{1}_{\{r<\widehat{\zeta}_e\}}\mathrm{d}r\right]$$

$$= \int_0^\infty \mathrm{d}x\psi(x)\,(\mathbb{P}_{0\nearrow x}\otimes\widehat{\boldsymbol{n}})\left[\int_0^\infty F_u\left(X^{(\zeta)}\bullet(x-\widehat{X}^{(u-\zeta)})\right)\mathbf{1}_{\{u-\zeta<\widehat{\zeta}_e\}}\mathbf{1}_{\{u>\zeta\}}\mathrm{d}u\right]. \tag{5.11}$$

This identity holds with F_t replaced by $\mathrm{e}^{-qt}F_t$ for any $q>0$, and hence, by uniqueness of the Laplace transform, we obtain

$$\mathbb{P}\left[F_t(X^{(t)})\right] = \int_0^\infty \mathrm{d}x\psi(x)\,(\mathbb{P}_{0\nearrow x}\otimes\widehat{\boldsymbol{n}})\left[F_t\left(X^{(\zeta)}\bullet(x-\widehat{X}^{(t-\zeta)})\right)\mathbf{1}_{\{t-\zeta<\widehat{\zeta}_e\}}\mathbf{1}_{\{t>\zeta\}}\right] \tag{5.12}$$

$$= \int_0^\infty \mathrm{d}x\psi(x)\int_0^t \mathbb{P}_{0\nearrow x}(\zeta\in\mathrm{d}u)\left(\mathbb{P}^{(u)}_{0\nearrow x}\otimes\widehat{\boldsymbol{n}}\right)\left[F_t\left(X^{(u)}\bullet(x-\widehat{X}^{(t-u)})\right)\mathbf{1}_{\{t-u<\widehat{\zeta}_e\}}\right] \tag{5.13}$$

$$= \int_0^\infty \mathrm{d}x\psi(x)\int_0^t \mathbb{P}_{0\nearrow x}(\zeta\in\mathrm{d}u)\boldsymbol{n}(\zeta_e>t-u)\left(\mathbb{P}^{(u)}_{0\nearrow x}\otimes\widehat{\mathbb{M}}^{(t-u)}_x\right)\left[F_t\left(X^{(u)}\bullet\widehat{X}^{(t-u)}\right)\right], \tag{5.14}$$

which completes the proof. □

Remark 5.6. (i) In the (α,ρ)-stable Lévy case with $\alpha\in(0,2]$ and $\rho\in(0,1)$, it is well-known that (see Lemma 3.2 in [13])

$$\boldsymbol{n}(\zeta_e>t) = K\cdot t^{-\rho}, \tag{5.15}$$

where $K>0$ is some constant, and hence we obtain from (3.6) and (5.3) that

$$\mathbb{P}\left(S_t\in\mathrm{d}x, g_t\in\mathrm{d}u\right) = \widetilde{K}\cdot\mathrm{d}x x^{\alpha\rho-1}\mathbb{P}_{0\nearrow x}(\zeta\in\mathrm{d}u)(t-u)^{-\rho}, \tag{5.16}$$

where $\widetilde{K}>0$ is other constant. Furthermore, together with the following well-known fact (see, e.g., Lemma VIII 13 in [4]) that

$$\mathbb{P}(g_t\in\mathrm{d}u) = \frac{1}{\Gamma(1-\rho)\Gamma(\rho)}u^{\rho-1}(t-u)^{-\rho}\mathrm{d}u, \tag{5.17}$$

then we obtain

$$\mathbb{P}(S_t\in\mathrm{d}x|g_t=u)\mathrm{d}u = K\cdot\mathrm{d}x x^{\alpha\rho-1}\mathbb{P}_{0\nearrow x}(\zeta\in\mathrm{d}u)\Gamma(1-\rho)\Gamma(\rho)u^{1-\rho}. \tag{5.18}$$

(ii) In the Brownian case, i.e., $\alpha=2$ and $\rho=1/2$ in (3.2), we note that $(X_t)\overset{\text{law}}{=}(W_{2ct})$ for a 1-dimensional standard Brownian motion (W_t), and we have the following:

$$\mathbb{P}(S_t\in\mathrm{d}x, g_t\in\mathrm{d}u) = \mathrm{d}x\mathrm{d}u\frac{x}{2c\pi\sqrt{u^3(t-u)}}\mathrm{e}^{-\frac{x^2}{4u}}; \tag{5.19}$$

$$\mathbb{P}_{0\nearrow x}(\zeta\in\mathrm{d}u) = \mathbb{P}(T_{\{x\}}\in\mathrm{d}u) = \mathrm{d}u\frac{x}{2c\sqrt{\pi u^3}}\mathrm{e}^{-\frac{x^2}{4u}}, \tag{5.20}$$

because of the following well-known facts (see, e.g., p. 102 and p. 80 in [11], respectively):

$$\mathbb{P}(\widetilde{S}_t \in \mathrm{d}x, \widetilde{g}_t \in \mathrm{d}u) = \mathrm{d}x\mathrm{d}u \frac{x}{\pi\sqrt{u^3(t-u)}} \mathrm{e}^{-\frac{x^2}{2u}}; \tag{5.21}$$

$$\mathbb{P}(\widetilde{T}_{\{x\}} \in \mathrm{d}u) = \mathrm{d}u \frac{x}{\sqrt{2\pi u^3}} \mathrm{e}^{-\frac{x^2}{2u}}, \tag{5.22}$$

where $\widetilde{S}_t = \sup_{s \le t} W_s$, $\widetilde{g}_t = \sup\{s \le t : W_s = \widetilde{S}_t\}$, and $\widetilde{T}_A = \inf\{s > 0 : W_s \in A\}$ for a Borel set $A \subset \mathbb{R}$. Thus we can easily check that the equality (5.16) is valid.

Remark 5.7. Assume moreover **(A3)**. Then, thanks to Bertoin's result; Corollary 3.2 in [3], it holds that

$$\lim_{t\to\infty} \mathbb{M}^{(t)}[F(X)] = \mathbb{P}_{0\downarrow 0}[F(X)], \tag{5.23}$$

where

$$\mathbb{M}^{(t)}[F(X)] = \mathbb{M}_0^{(t)}[F(X)] = \frac{\boldsymbol{n}\left[F(-X^{(t)}); \zeta_e > t\right]}{\boldsymbol{n}(\zeta_e > t)}. \tag{5.24}$$

6 Generalised Azéma-Yor martingales and definition of a probability measure $\mathbb{P}^{(f)}$

Let us introduce a generalisation of (1.4) and (1.11). Let $X = ((X_t), \mathbb{P})$ be a Lévy process with notation given in Section 2 and assume **(A1)**, **(A2)** and **(A3)**. Let ψ and h be the functions given by (2.10) and (2.12), respectively. Let f be a non-negative Borel function on $[0, \infty)$ satisfying

$$(0 <) \int_0^\infty f(x)\psi(x)\mathrm{d}x < \infty. \tag{6.1}$$

We introduce the process $(M_t^{(f)}, t \ge 0)$ by

$$M_t^{(f)} = f(S_t)h(S_t - X_t) + \int_{S_t}^\infty f(x)\psi(x - X_t)\mathrm{d}x. \tag{6.2}$$

Theorem 6.1. *$(M_t^{(f)}, t \ge 0)$ is a $((\mathscr{F}_t), \mathbb{P})$-martingale.*

The proof of Theorem 6.1 is done in the same way as in [28] in the stable Lévy case; the coinvariance of the function h plays a key role. Thus we omit it.

We introduce the probability measure $\mathbb{P}^{(f)}$ on $\mathscr{F}_\infty$ as follows:

$$\mathbb{P}^{(f)}|_{\mathscr{F}_t} = \frac{M_t^{(f)}}{M_0^{(f)}} \cdot \mathbb{P}|_{\mathscr{F}_t}. \tag{6.3}$$

Since $(M_t^{(f)})$ is a martingale, the consistency holds, and hence $\mathbb{P}^{(f)}$ is well-defined.

7 The σ-finite measure which unifies the supremum penalisations

Let us consider a Lévy process $X = ((X_t), \mathbb{P})$ with $\mathbb{P}(X_0 = 0) = 1$. In this section we assume:

(A1), i.e., absolute continuity condition for the resolvent;

(A2) & **(A2*)**. i.e., 0 is regular for both $(0, \infty)$ and $(-\infty, 0)$ with respect to X;

(A3) & **(A3*)**, i.e., $I_\infty = -\infty$ and $S_\infty = \infty$ $\mathbb{P}$-a.s.,

where I_∞ and S_∞ are the overall infimum and supremum of X_t, respectively, i.e., $I_\infty = \inf\{X_t : t \geq 0\}$ and $S_\infty = \sup\{X_t : t \geq 0\}$. Remark again that the condition **(B)** in the (α, ρ)-stable Lévy case implies all the above conditions.

We introduce $\mathcal{P}_{\sup}$ as follows.

Definition 7.1. Define

$$\mathcal{P}_{\sup} = \int_0^\infty \mathrm{d}x \psi(x) (\mathbb{P}_{0\nearrow x} \bullet \mathbb{P}_{x\downarrow x}), \tag{7.1}$$

where $\mathbb{P}_{0\nearrow x}$ denotes the law of $X + x$ under $\mathbb{P}_{-x\nearrow 0}$, i.e., $\mathbb{P}_{0\nearrow x}$ denotes the law of the process starting from 0 and conditioned to hit x continuously, and $\mathbb{P}_{x\downarrow x}$ denotes the law of $X + x$ under $\mathbb{P}_{0\downarrow 0}$, i.e., $\mathbb{P}_{x\downarrow x}$ denotes the law of the process starting from x and conditioned to stay below level x.

Denote

$$g = \sup\{t \geq 0 : X_t = S_\infty\}. \tag{7.2}$$

Theorem 7.2. *The following statements hold:*

(i) $\mathcal{P}_{\sup}(S_\infty \in \mathrm{d}x, g \in \mathrm{d}u) = \mathrm{d}x \psi(x) \mathbb{P}_{0\nearrow x}(\zeta \in \mathrm{d}u)$,
in particular, $\mathcal{P}_{\sup}(S_\infty \in \mathrm{d}x) = \mathrm{d}x \psi(x)$*;*

(ii) $\mathcal{P}_{\sup}$ *is a* σ*-finite measure on* $\mathscr{F}_\infty$*;*

(iii) $\mathcal{P}_{\sup}$ *is singular to* $\mathbb{P}$ *on* $\mathscr{F}_\infty$*.*

(iv) For each $t > 0$ *and* $A \in \mathscr{F}_t$*, it holds that*

$$\mathcal{P}_{\sup}(A) = \begin{cases} 0, & \text{if } \mathbb{P}(A) = 0; \\ \infty, & \text{if } \mathbb{P}(A) > 0. \end{cases} \tag{7.3}$$

Consequently, $\mathcal{P}_{\sup}$ *is not* σ*-finite on* $\mathscr{F}_t$ *for* $t < \infty$*.*

Proof. (i) We have

$$\mathcal{P}_{\sup} = \int_0^\infty \mathrm{d}x \psi(x) \int_0^\infty \mathbb{P}_{0\nearrow x}(\zeta \in \mathrm{d}u) \left(\mathbb{P}^{(u)}_{0\nearrow x} \bullet \mathbb{P}_{x\downarrow x} \right), \tag{7.4}$$

and hence

$$\begin{aligned}\mathcal{P}_{\sup}[F(S_\infty)G(g)] &= \int_0^\infty \mathrm{d}x\psi(x)\int_0^\infty \mathbb{P}_{0\nearrow x}(\zeta\in \mathrm{d}u)\left(\mathbb{P}^{(u)}_{0\nearrow x}\bullet \mathbb{P}_{x\downarrow x}\right)[F(S_\infty)G(g)]\\ &= \int_0^\infty \mathrm{d}x\psi(x)F(x)\int_0^\infty \mathbb{P}_{0\nearrow x}(\zeta\in \mathrm{d}u)G(u),\end{aligned}$$

for any test functions F and G. Thus we obtain the desired result.

(ii) For each $x>0$, $\mathcal{P}_{\sup}(S_\infty<x)=\int_0^x\psi(y)\mathrm{d}y$ is finite, which shows the desired conclusion.

(iii) We have $\mathcal{P}_{\sup}(S_\infty=\infty)=0$ by definition. On the other hand, we have $\mathbb{P}(S_\infty<\infty)=0$ by our assumption **(A3*)**. This implies that $\mathcal{P}_{\sup}$ is singular to $\mathbb{P}$ on $\mathscr{F}_\infty$.

(iv) Suppose that $\mathbb{P}(A)=0$ for $A\in\mathscr{F}_t$. We have

$$\begin{aligned}\mathcal{P}_{\sup}(A) &= \int_0^\infty \mathrm{d}x\psi(x)\left(\mathbb{P}_{0\nearrow x}\bullet\mathbb{P}_{x\downarrow x}\right)(A)\\ &= \int_0^\infty \mathrm{d}x\psi(x)\left(\mathbb{P}_{0\nearrow x}\bullet\mathbb{P}_{x\downarrow x}\right)[\mathbf{1}_A;t<\zeta]+\int_0^\infty \mathrm{d}x\psi(x)\left(\mathbb{P}_{0\nearrow x}\bullet\mathbb{P}_{x\downarrow x}\right)[\mathbf{1}_A;t\geq\zeta]\\ &=: I_1+I_2.\end{aligned}$$

On one hand, we have

$$\begin{aligned}I_1 &= \int_0^\infty \mathrm{d}x\psi(x)\mathbb{P}_{0\nearrow x}[\mathbf{1}_A;t<\zeta]\\ &= \int_0^\infty \mathrm{d}x\mathbb{P}\left[\psi(x-X_t)\mathbf{1}_{\{t<T_{(x,\infty)}\}}\mathbf{1}_A\right]\quad \text{(by (4.8))}\\ &= 0.\end{aligned}$$

On the other hand, we have

$$\begin{aligned}I_2 &= \int_0^\infty \mathrm{d}x\psi(x)\left(\mathbb{P}_{0\nearrow x}\bullet\mathbb{P}_{x\downarrow x}\right)[\mathbf{1}_A(X);t\geq\zeta]\\ &= \int_0^\infty \mathrm{d}x\psi(x)\left(\mathbb{P}_{0\nearrow x}\otimes\widehat{\mathbb{P}}_{0\downarrow 0}\right)\left[\mathbf{1}_A\left(X^{(\zeta)}\bullet\left(x+\widehat{X}^{(t-\zeta)}\right)\right)\mathbf{1}_{\{t\geq\zeta\}}\right]\\ &= \int_0^\infty \mathrm{d}x\psi(x)\left(\mathbb{P}_{0\nearrow x}\otimes\widehat{\boldsymbol{n}}\right)\left[h(\widehat{X}_{t-\zeta})\mathbf{1}_{\{t-\zeta<\widehat{\zeta}_e\}}\mathbf{1}_A\left(X^{(\zeta)}\bullet\left(x-\widehat{X}^{(t-\zeta)}\right)\right)\mathbf{1}_{\{t\geq\zeta\}}\right],\end{aligned}\tag{7.5}$$

by the definition of $\mathbb{P}_{0\downarrow 0}$. Then

$$\begin{aligned}(7.5) &= \int_0^\infty \mathrm{d}x\psi(x)\left(\mathbb{P}_{0\nearrow x}\otimes\widehat{\boldsymbol{n}}\right)\left[h\left(x-(x-\widehat{X}_{t-\zeta})\right)\mathbf{1}_A(X)\mathbf{1}_{\{0\leq t-\zeta<\widehat{\zeta}_e\}}\right]\\ &= \int_0^\infty \mathrm{d}x\psi(x)\left(\mathbb{P}_{0\nearrow x}\otimes\widehat{\boldsymbol{n}}\right)\left[h\left(x-\left(X^{(\zeta)}\bullet(x-\widehat{X}^{(t-\zeta)})\right)_t\right)\mathbf{1}_A(X)\mathbf{1}_{\{0\leq t-\zeta<\widehat{\zeta}_e\}}\right]\\ &= \mathbb{P}\left[h(S_t-X_t)\mathbf{1}_A\right]\quad \text{(by Theorem 5.1)}\\ &= 0.\end{aligned}$$

Thus we obtain $\mathcal{P}_{\sup}(A)=0$.

Conversely, suppose that $\mathbb{P}(A) > 0$ for $A \in \mathscr{F}_t$. Then we see that

$$\begin{aligned}
\mathcal{P}_{\sup}(A) &\geq \int_0^\infty \mathrm{d}x\psi(x)\mathbb{P}_{0\nearrow x}\left[\mathbf{1}_A; t < \zeta\right] \\
&= \int_0^\infty \mathrm{d}x\mathbb{P}\left[\psi(x - X_t)\mathbf{1}_{\{t<T_{(x,\infty)}\}}\mathbf{1}_A\right] \\
&\geq \int_1^\infty \mathrm{d}x\mathbb{P}\left[\psi(x - X_t)\mathbf{1}_{\{t<T_{(1,\infty)}\}}\mathbf{1}_A\right] \\
&= \mathbb{P}\left[\{h(\infty) - h(1 - X_t)\}\,\mathbf{1}_{\{t<T_{(1,\infty)}\}}\mathbf{1}_A\right].
\end{aligned}$$

Since we have

$$h(\infty) = \lim_{x\to\infty} h(x) = \lim_{x\to\infty} \mathbb{P}\left[\int_0^\infty \mathbf{1}_{\{S_t\leq x\}}\mathrm{d}L_t\right] = \mathbb{P}\left[\int_0^\infty \mathrm{d}L_t\right] = \mathbb{P}[L_\infty] = \infty,$$

thus $\mathcal{P}_{\sup}(A) = \infty$. Therefore the proof is completed. □

We shall give some relationships between the measures $\mathcal{P}_{\sup}$, $\mathbb{P}$ and $\mathbb{P}^{(f)}$.

Theorem 7.3. *It holds that*

$$\mathcal{P}_{\sup}\left[f(S_\infty)F_t(X)\right] = \mathbb{P}\left[M_t^{(f)}F_t(X)\right], \tag{7.6}$$

for any $\mathcal{F}_t$-measurable functional $F_t(X)$. Consequently, one has

$$\frac{\mathcal{P}_{\sup}\left[f(S_\infty)F_t(X)\right]}{\mathcal{P}_{\sup}\left[f(S_\infty)\right]} = \mathbb{P}\left[\frac{M_t^{(f)}}{M_0^{(f)}}F_t(X)\right] = \mathbb{P}^{(f)}[F_t(X)], \tag{7.7}$$

and

$$\frac{f(S_\infty)\cdot\mathcal{P}_{\sup}}{\mathcal{P}_{\sup}[f(S_\infty)]} = \mathbb{P}^{(f)} \quad \textit{on} \quad \mathscr{F}_\infty. \tag{7.8}$$

Proof. Recall the computation in the proof of Theorem 7.2 (iv). We have

$$\begin{aligned}
\mathcal{P}_{\sup}\left[f(S_\infty)F_t(X)\right] &= \int_0^\infty \mathrm{d}x\psi(x)(\mathbb{P}_{0\nearrow x}\bullet\mathbb{P}_{x\downarrow x})\left[f(S_\infty)F_t(X)\right] \\
&= \int_0^\infty \mathrm{d}x\psi(x)f(x)(\mathbb{P}_{0\nearrow x}\bullet\mathbb{P}_{x\downarrow x})\left[F_t(X)\right],
\end{aligned} \tag{7.9}$$

since $S_\infty = x$ under the measure $\mathbb{P}_{0\nearrow x}\bullet\mathbb{P}_{x\downarrow x}$. Then

$$\begin{aligned}
(7.9) &= \int_0^\infty \mathrm{d}x\psi(x)f(x)(\mathbb{P}_{0\nearrow x}\bullet\mathbb{P}_{x\downarrow x})\left[F_t(X); t < \zeta\right] \\
&\quad + \int_0^\infty \mathrm{d}x\psi(x)f(x)(\mathbb{P}_{0\nearrow x}\bullet\mathbb{P}_{x\downarrow x})\left[F_t(X); t \geq \zeta\right] \\
&=: I_1 + I_2.
\end{aligned}$$

On one hand, we have

$$I_1 = \int_0^\infty \mathrm{d}x\psi(x)f(x)\mathbb{P}_{0\nearrow x}\left[F_t(X); t<\zeta\right] = \int_0^\infty \mathrm{d}x f(x)\mathbb{P}\left[\psi(x-X_t)\mathbf{1}_{\{t<T_{(x,\infty)}\}}F_t(X)\right]$$
$$= \mathbb{P}\left[F_t(X)\int_0^\infty \mathrm{d}x f(x)\psi(x-X_t)\mathbf{1}_{\{S_t\le x\}}\right]. \tag{7.10}$$

On the other hand, we obtain from the same computation in the proof of (iv) in the previous theorem that

$$\begin{aligned} I_2 &= \int_0^\infty \mathrm{d}x\psi(x)f(x)(\mathbb{P}_{0\nearrow x}\bullet\mathbb{P}_{x\downarrow x})\left[F_t(X); t\ge\zeta\right] \\ &= \int_0^\infty \mathrm{d}x\psi(x)f(x)(\mathbb{P}_{0\nearrow x}\otimes\widehat{\boldsymbol{n}})\left[h(x-X_t)F_t(X)\mathbf{1}_{\{0\le t-\zeta<\widehat{\zeta}_e\}}\right] \\ &= \int_0^\infty \mathrm{d}x\psi(x)(\mathbb{P}_{0\nearrow x}\otimes\widehat{\boldsymbol{n}})\left[f(S_t)h(S_t-X_t)\mathbf{1}_{\{t-\zeta<\widehat{\zeta}_e\}}F_t(X)\mathbf{1}_{\{t\ge\zeta\}}\right]. \end{aligned} \tag{7.11}$$

By Theorem 5.1, we get

$$(7.11) = \mathbb{P}\left[f(S_t)h\left(S_t-X_t\right)F_t\left(X\right)\right]. \tag{7.12}$$

Combining (7.10) and (7.12), we obtain

$$\begin{aligned} \mathcal{P}_{\sup}\left[f(S_\infty)F_t(X)\right] &= \mathbb{P}\left[F_t(X)\int_{S_t}^\infty \mathrm{d}x f(x)\psi(x-X_t)\right] + \mathbb{P}\left[F_t(X)f(S_t)h\left(S_t-X_t\right)\right] \\ &= \mathbb{P}\left[F_t(X)\left\{\int_{S_t}^\infty \mathrm{d}x f(x)\psi(x-X_t) + f(S_t)h\left(S_t-X_t\right)\right\}\right], \end{aligned} \tag{7.13}$$

that is,

$$\mathcal{P}_{\sup}\left[f(S_\infty)F_t(X)\right] = \mathbb{P}\left[M_t^{(f)}F_t(X)\right]. \tag{7.14}$$

Especially, when $t=0$, we have

$$\mathcal{P}_{\sup}\left[f(S_\infty)\right] = \int_0^\infty \mathrm{d}x f(x)\psi(x). \tag{7.15}$$

Therefore we obtain

$$\frac{\mathcal{P}_{\sup}\left[f(S_\infty)F_t(X)\right]}{\mathcal{P}_{\sup}\left[f(S_\infty)\right]} = \mathbb{P}\left[\frac{M_t^{(f)}}{M_0^{(f)}}F_t(X)\right] = \mathbb{P}^{(f)}[F_t(X)]. \tag{7.16}$$

This completes the proof. □

Remark 7.4. Recently Najnudel-Nikeghbali [14] gave a generalization of $\mathcal{W}$. A non-negative submartingale $(X_t, t\ge 0)$ is said to be of the class (Σ) if it can be decomposed as $X_t = N_t + A_t$ where $(N_t, t\ge 0)$ and $(A_t, t\ge 0)$ are $\mathcal{F}_t$-adapted process, (N_t) is a càdlàg martingale, and (A_t) is a continuous increasing process which is carried by the set of zeros with $A_0 = 0$. Then they proved that there exists a σ-finite measure $\mathcal{Q}$ such that

$$\mathcal{Q}\left[F_t; g\le t\right] = \mathbb{P}\left[F_t X_t\right], \tag{7.17}$$

for all non-negative $\mathcal{F}_t$-measurable functional F_t. Here g is the last exit time from 0. It may be quite natural to ask now whether the process $(h(S_t-X_t), t\ge 0)$ is of the class (Σ) or not. However, we have not succeeded in answering the question.

The measure $\mathcal{P}_{\sup}$ does not depend upon f. Recall that $\mathbb{P}^{(f)}$ is the limit measure of supremum penalisation. The measure $\mathcal{P}_{\sup}$ implies the following fact that gives the detailed description of $\mathbb{P}^{(f)}$.

Theorem 7.5. *One has*

$$\mathbb{P}^{(f)} = \int_0^\infty \mathbb{P}^{(f)}(S_\infty \in \mathrm{d}x)(\mathbb{P}_{0\nearrow x} \bullet \mathbb{P}_{x\downarrow x}). \tag{7.18}$$

That is, it holds that, under $\mathbb{P}^{(f)}$,

(i) $\mathbb{P}^{(f)}(S_\infty \in \mathrm{d}x) = \dfrac{1}{M_0^{(f)}}\psi(x)f(x)\mathrm{d}x$ *where* $M_0^{(f)} = \displaystyle\int_0^\infty \psi(x)f(x)\mathrm{d}x$;

(ii) given $g = u$, $(X_s, s \le u)$ *and* $(X_u - X_{u+s}, s \ge 0)$ *are independent;*

(iii) given $S_\infty = x$ *and* $g = u$, $(X_s, s \le u)$ *is distributed as the process conditioned to hit* x *continuously with duration* u;

(iv) given $S_\infty = x$ *and* $g = u$, $(-(x - X_{u+s}), s \ge 0)$ *is distributed as the process conditioned to stay negative.*

Under our assumption in this section, the following result for the martingale $(M_t^{(f)})$ can be proved.

Theorem 7.6. *Let* $X = ((X_t), \mathbb{P})$ *be a Lévy process with* **(A1)**, **(A2)**, **(A2*)**, **(A3)** *and* **(A3*)**, *and let* $M_t^{(f)}$ *be the process given in* (6.2). *Then* $M_t^{(f)}$ *converges to* 0 $\mathbb{P}$*-a.s. as* $t \to \infty$.

Proof. We show that $M_t^{(f)} \to 0$ $\mathbb{P}$-a.s. through the measure $\mathcal{P}_{\sup}$. Since $(M_t^{(f)})$ is a non-negative $\mathbb{P}$-martingale as proved before, there exists a $\mathscr{F}_\infty$-measurable functional $M_\infty^{(f)}$ such that $M_t^{(f)} \to M_\infty^{(f)}$ $\mathbb{P}$-a.s. by the martingale convergence theorem. For $a > 0$,

$$\begin{aligned}
\mathbb{P}\left[M_\infty^{(f)}\right] &= \mathbb{P}\left[M_\infty^{(f)}\mathbf{1}_{\{S_\infty \ge a\}}\right] \quad \text{(by the fact that } \mathbb{P}(S_\infty = \infty) = 1) \\
&\le \liminf_{t\to\infty} \mathbb{P}\left[M_t^{(f)}\mathbf{1}_{\{S_t \ge a\}}\right] \quad \text{(by Fatou's lemma)} \\
&= \liminf_{t\to\infty} \mathcal{P}_{\sup}\left[f(S_\infty)\mathbf{1}_{\{S_t \ge a\}}\right] \quad \text{(by (7.7))} \\
&= \mathcal{P}_{\sup}\left[f(S_\infty)\mathbf{1}_{\{S_\infty \ge a\}}\right]. \quad \text{(by the dominated convergence theorem)}
\end{aligned}$$

Letting $a \to \infty$, then $\mathcal{P}_{\sup}\left[f(S_\infty)\mathbf{1}_{\{S_\infty \ge a\}}\right] \to 0$. Thus $\mathbb{P}[M_\infty^{(f)}] = 0$, and therefore we obtain $\mathbb{P}(M_\infty^{(f)} = 0) = 1$. □

8 Some remarks on $\mathcal{P}$ and $\mathcal{P}_{\sup}$

Let us consider a symmetric (i.e., $\rho = 1/2$) stable Lévy process X with index $\alpha \in (1, 2]$, and recall the σ-finite measure $\mathcal{P}$ which is given in [27] (see also [25]):

$$\mathcal{P} = \int_0^\infty \mathbb{P}[\mathrm{d}L_u^X](\mathbb{Q}^{(u)} \bullet \mathbb{P}^\times), \tag{8.1}$$

where L_t^X denotes the local time at 0 of X itself, $\mathbb{Q}^{(u)}$ denotes the law of the stable bridge from 0 to 0 with length u and $\mathbb{P}^\times$ denotes the h-transform process with respect to the harmonic function $|x|^{\alpha-1}$ of the process killed at the first hitting time of 0. On comparison, it becomes clear that the two σ-finite measures $\mathcal{P}_{\sup}$ and $\mathcal{P}$ are quite different: $\mathcal{P}_{\sup}$ is based on the excursion theory for the reflected process of a Lévy process, whereas $\mathcal{P}$ comes from the excursion theory for a Lévy process itself. We stress that this difference cannot appear in the Brownian case because of the fact that $(S_t, S_t - X_t)_{t\geq 0} \overset{\text{law}}{=} (L_t^X, |X_t|)_{t\geq 0}$ which is known as Lévy's theorem.

Finally, we would like to emphasize the following fact again:

$$\mathcal{P}(S_\infty < \infty) = 0 \quad \text{and} \quad \mathcal{P}_{\sup}(S_\infty = \infty) = 0. \tag{8.2}$$

We mention the relationship between $\mathcal{P}_{\sup}$ and $\mathcal{P}$ as follows:

(i) $\mathcal{P} \perp \mathcal{P}_{\sup}$ on $\mathscr{F}_\infty$;

(ii) if $A \in \mathscr{F}_t$, then

$$\mathcal{P}(A) > 0 \quad \Longleftrightarrow \quad \mathcal{P}_{\sup}(A) > 0, \tag{8.3}$$

and both are infinite.

Acknowledgement. The author would like to thank Professor Marc Yor and Professor Loïc Chaumont for many valuable comments and helpful suggestions. She also thanks the referee for careful reading of the manuscript and suggesting some improvements.

References

[1] J. Azéma and M. Yor, Une solution simple au problème de Skorokhod, *Séminaire de Probabilités, XIII* (Univ. Strasbourg, Strasbourg, 1977/78), Lecture Notes in Math. **721**, Springer, Berlin, 1979, 90–115.

[2] J. Azéma and M. Yor, Le problème de Skorokhod: compléments à "Une solution simple au problème de Skorokhod", *Séminaire de Probabilités, XIII* (Univ. Strasbourg, Strasbourg, 1977/78), Lecture Notes in Math. **721**, Springer, Berlin, 1979, 625–633.

[3] J. Bertoin, Splitting at the infimum and excursions in half-lines for random walks and Lévy processes, *Stoch. Process. Appl.*, **47**, 17–35, 1993.

[4] J. Bertoin, *Lévy processes*, Cambridge Univ. Press, Cambridge, 1996.

[5] N. Bingham, Maxima of sums of random variables and suprema of stable processes, *Z. Wahrsch. Verw. Gebiete*, **26**, 273–296, 1973.

[6] L. Chaumont, Conditionings and path decompositions for Lévy processes, *Stoch. Process. Appl.*, **64**, 39–54, 1996.

[7] L. Chaumont, Excursion normalisée, méandre et pont pour des processus stables, *Bull. Sci. Math.*, **121**, 377–403, 1997.

[8] L. Chaumont, On the law of the supremum of Lévy processes, preprint.

[9] L. Chaumont and R. A. Doney, On Lévy processes conditioned to stay positive, *Electron. J. Probab.*, **10**: no. 28, 948–961 (electronic), 2005; corrections in **13**: no. 1, 1–4 (electronic), 2008.

[10] R. A. Doney, *Fluctuation theory for Lévy processes*, Lectures from the 35th Summer School on Probability Theory held in Saint-Flour, July 6–23, 2005, Lecture Notes in Math., **1897**, Springer, Berlin, 2007.

[11] I. Karatzas and S. E. Shreve, *Brownian Motion and Stochastic Calculus*, Second edition, Springer, Berlin-Heidelberg-New York, 1991.

[12] A. E. Kyprianou, *Introductory lectures on fluctuations of Lévy processes with applications*, Universitext. Springer-Verlag, Berlin, 2006.

[13] D. Monrad and M. L. Silverstein, Stable processes: sample function growth at a local minimum, *Z. Wahrsch. Verw. Gebiete*, **49**, 177–210, 1979.

[14] J. Najnudel, A. Nikeghbali On some properties of a universal sigma-finite measure associated with a remarkable class of submartingales, *Publ. RIMS, Kyoto Univ.*, **47**, 911–936, 2011.

[15] J. Najnudel, B. Roynette and M. Yor, *A global view of Brownian penalisations*, MSJ Memoirs, **19**, Mathematical Society of Japan, Tokyo, 2009.

[16] J. Obłój, The Skorokhod embedding problem and its offsprings, *Probability Surveys*, **1**, 321–392, 2004.

[17] J. Pitman and M. Yor, Decomposition at the maximum for excursions and bridges of one-dimensional diffusions, *Itô's stochastic calculus and probability theory*, 293–310, Springer, Tokyo, 1996.

[18] D. Revuz and M. Yor, *Continuous martingales and Brownian motion*, Third edition, Springer-Verlag, Berlin, 1999.

[19] B. Roynette, P. Vallois and M. Yor, Limiting laws associated with Brownian motion perturbed by normalized exponential weights, I, *Studia. Sci. Math. Hungar.*, **43**, 171–246, 2006.

[20] B. Roynette, P. Vallois and M. Yor, Limiting laws associated with Brownian motion perturbed by its maximum, minimum and local time, II, *Studia. Sci. Math. Hungar.*, **43**, 295–360, 2006.

[21] B. Roynette, P. Vallois and M. Yor, Some penalisations of the Wiener measure, *Jpn. J. Math.*, **1**, 263–290, 2006.

[22] B. Roynette and M. Yor, *Penalising brownian paths*, Lecture Notes in Math., **1969**, Springer, 2009.

[23] K. Sato, *Lévy processes and infinitely divisible distributions*, Translated from the 1990 Japanese original, Revised by the author, Cambridge Studies in Advanced Mathematics, **68**, Cambridge University Press, Cambridge, 1999.

[24] M. L. Silverstein, Classification of coharmonic and coinvariant functions for Lévy processes, *Ann. Probab.*, **8**, 539–575, 1980.

[25] K. Yano, Two kinds of conditionings for stable Lévy processes, to appear in *Proceedings of the 1st MSJ-SI, "Probabilistic Approach to Geometry", Adv. Stud. Pure Math.*, Math. Soc. Japan.

[26] K. Yano, Excursions away from a regular point for one-dimensional symmetric Lévy processes without Gaussian part, *Potential Anal.*, **32**, 305–341, 2010.

[27] K. Yano, Y. Yano and M. Yor, Penalising symmetric stable Lévy paths, *J. Math. Soc. Japan*, **61**, no. 3, 757–798, 2009.

[28] K. Yano, Y. Yano and M. Yor, Penalisation of a stable Lévy process involving its one-sided supremum, to appear in Ann. Inst. H. Poincaré Probab. Statist.

[29] V. M. Zolotarev, *One-dimensional stable distributions*, Translations of Mathematical Monographs, **65**, American Mathematical Society, Providence, RI, 1986.