BEHAVIORS OF ENTROPY ON FINITELY GENERATED GROUPS

JÉRÉMIE BRIEUSSEL

INSTITUT DE MATHÉMATIQUES RUE EMILE-ARGAND 11 CH-2000 NEUCHÂTEL SWITZERLAND

ABSTRACT. A variety of behaviors of entropy functions of random walks on finitely generated groups is presented, showing that for any $\frac{1}{2} \le \alpha \le \beta \le 1$, there is a group Γ with measure μ equidistributed on a finite generating set such that:

$$\liminf \frac{\log H_{\Gamma,\mu}(n)}{\log n} = \alpha \text{ and } \limsup \frac{\log H_{\Gamma,\mu}(n)}{\log n} = \beta.$$

The groups involved are finitely generated subgroups of the group of automorphisms of an extended rooted tree. The return probability and the drift of a simple random walk Y_n on such groups are also evaluated, providing an example of group with return probability satisfying:

$$\liminf \frac{\log |\log P(Y_n =_{\Gamma} 1)|}{\log n} = \frac{1}{3} \text{ and } \limsup \frac{\log |\log P(Y_n =_{\Gamma} 1)|}{\log n} = 1,$$

and drift satisfying:

$$\liminf \frac{\log \mathbb{E}||Y_n||}{\log n} = \frac{1}{2} \text{ and } \limsup \frac{\log \mathbb{E}||Y_n||}{\log n} = 1.$$

1. Introduction

The caracterisation of groups by an asymptotic description of their Cayley graphs may be dated back to Folner's criterion of amenability by quasi-invariant subsets [Fol]. Shortly after, Kesten showed equivalence with the probabilistic criterion that a group Γ is amenable if and only if the return probability $P(Y_n = 1)$ of a simple random walk Y_n on Γ decays exponentialy fast [Kes].

This article focuses on three quantities that partially describe the behavior of the diffusion process of a random walk Y_n with step distribution μ on a group Γ . Namely the entropy function $H_{\Gamma,\mu}(n) = H(\mu^{*n}) = H(Y_n)$, the return probability $P(Y_n = 1)$ and the drift, also called rate of escape, $L_{\Gamma,\mu}(n) = \mathbb{E}_{\mu^{*n}}||\gamma|| = \mathbb{E}||Y_n||$, where ||.|| is a word norm,

$$H(\mu) = -\sum_{\gamma \in \Gamma} \mu(\gamma) \log \mu(\gamma)$$

is the Shannon entropy of the probability measure μ , and μ^{*n} is the *n*-fold convolution of μ , or in other terms the distribution of Y_n . The return probability for

Date: March 29, 2012. Supported by Swiss NSF grant 20-126689.

a finitely supported symmetric law μ is a group invariant by [PSC1], which is not known to be the case for entropy and drift. However, sublinearity of the entropy or of the drift for some measure μ with generating support implies amenability [KV]. In the present paper, the measure μ will always be equidistributed on a canonical finite symmetric generating set of Γ .

The asymptotic behavior of these functions has been precisely established in a number of cases, that mainly include virtually nilpotent groups, linear groups and wreath products ([Var],[PSC2],[PSC3],[Ers1],[Ers2],[Ers5],[Rev]), and a variety of less precise estimates exist for some groups acting on rooted trees ([BV],[Kai],[Ers4], [BKN],[AAV],[AV]).

The object of this article is to present examples of groups that provide new asymptotic behaviors for these probabilistic functions. Entropy, return probability and drift functions will not be precisely computed, but only mild approximations in terms of their exponents.

Definition 1.1. The *lower* and *upper entropy exponents* of a random walk Y_n of law μ on a group Γ are respectively:

$$\underline{h}(\Gamma,\mu) = \liminf \frac{\log H_{\Gamma,\mu}(n)}{\log n} \text{ and } \overline{h}(\Gamma,\mu) = \limsup \frac{\log H_{\Gamma,\mu}(n)}{\log n}.$$

The lower and upper return probability exponents of a random walk Y_n of symmetric finitely supported law μ on a group Γ are respectively:

$$\underline{p}(\Gamma) = \liminf \frac{\log |\log P(Y_n = 1)|}{\log n} \text{ and } \overline{p}(\Gamma) = \limsup \frac{\log |\log P(Y_n = 1)|}{\log n}.$$

The lower and upper drift exponents of a random walk Y_n of law μ on a group Γ are respectively:

$$\underline{\delta}(\Gamma,\mu) = \liminf \frac{\log L_{\Gamma,\mu}(n)}{\log n} \text{ and } \overline{\delta}(\Gamma,\mu) = \limsup \frac{\log L_{\Gamma,\mu}(n)}{\log n}.$$

When equality holds, the entropy exponent of the group Γ with law μ is $h(\Gamma, \mu) = \overline{h}(\Gamma, \mu) = \underline{h}(\Gamma, \mu)$, the return probability exponent of a group Γ is $p(\Gamma) = \underline{p}(\Gamma) = \overline{p}(\Gamma)$ and the drift exponent of the group Γ with law μ is $\delta(\Gamma, \mu) = \overline{\delta}(\Gamma, \mu) = \underline{\delta}(\Gamma, \mu)$. Return probability exponents do not depend on the particular choice of the measure by [PSC1].

Computing the exponents gives moderate information on the function. By [PSC2], the wreath product $\mathbb{Z} \wr \mathbb{Z}$ has return probability $P(Y_n = 1) \approx \exp(-n^{\frac{1}{3}}(\log n)^{\frac{2}{3}})$, and the lamplighter $F \wr \mathbb{Z}$ with finite group F has return probability $P(Y_n = 1) \approx \exp(-n^{\frac{1}{3}})$. Both have return probability exponent $\frac{1}{3}$. Exponent 1 does not imply linearity of entropy or drift, nor exponential decay of return probability, as seen by the exemples in [Ers4].

The groups considered here are directed groups of automorphisms of extended rooted trees. The main construction combines the directed groups of [Bri2] with the notion of boundary permutational extension introduced in [BE] and used in [Bri3] to exhibit various behaviors of growth functions on groups. The entropy exponents

can be computed explicitly in terms of the group construction (see theorem 5.1), which leads to the following corollary:

Theorem 1.2. For any $\frac{1}{2} \leq \alpha \leq \beta \leq 1$, there exists a finitely generated group $\Gamma = \Gamma(\alpha, \beta)$ and a symmetric finitely supported measure μ such that:

$$\underline{h}(\Gamma, \mu) = \alpha \text{ and } \overline{h}(\Gamma, \mu) = \beta.$$

In particular when $\alpha = \beta$, there is a finitely generated group $\Gamma(\beta)$ with measure μ such that:

$$h(\Gamma, \mu) = \beta.$$

Entropy is related to growth because over a finite set, entropy is maximized for equidistribution probability, so that:

$$h_{\Gamma,\mu}(r) = h(\mu^{*n}) \le \log \#supp(\mu^{*n}) = \log b_{\Gamma,S}(r).$$

The groups of theorem 1.2 have sublinear entropy and often exponential growth. However, most of the groups Γ_{ω} of [Bri3] have intermediate growth and are extended directed groups of a binary rooted tree, so by theorem 5.1 they all have entropy exponent $h(\Gamma, \mu) = \frac{1}{2}$.

The return probability and drift exponents of extended directed groups can be estimated from above and from below, but the bounds do not match in general. A specific example provides the:

Theorem 1.3. There exists a finitely generated group Δ and a symmetric finitely supported measure μ' such that:

$$\underline{p}(\Delta) = \frac{1}{3}, \overline{p}(\Delta) = 1 \text{ and } \underline{\delta}(\Delta, \mu') = \frac{1}{2}, \overline{\delta}(\Delta, \mu') = 1.$$

These theorems show that the phenomenon of oscillation studied in [Bri3] for growth function exponents (see also [Bri1] and [KP]) also occurs for entropy, return probability and drift. The existence of a group with such drift exponents was mentionned without proof in [Ers6].

The article is structured as follows. The main construction of extended directed groups of a rooted tree $T_{\bar{d}}$ is described in section 2, where is also presented a side application to the Haagerup property of groups with non-uniform growth. Section 3 presents the basic tool to study these groups which is the rewritting process, leading to the notions of minimal tree and activity, related to inverted orbits of permutational extensions defined in [BE]. Section 4 relates the expected activity of a random walk Y_n to the exponent sequence, which depends only on the tree $T_{\bar{d}}$. At this stage one can prove the main theorem 5.1 on entropy, which implies theorem 1.2 for $\beta < 1$ and permits to derive estimates on the drift. The frequency of oscillation of entropy exponents is also studied in section 5. The main estimates on return probability of extended directed groups are given in theorem 6.1, with a specific example related to the lamplighter group. Section 7 is devoted to another construction adapted from [KP] and similar to [Ers4], which permits to obtain the case $\beta = 1$ in theorem 1.2 and to prove theorem 1.3. A generalisation of the construction of extended directed groups is presented in section 8, followed by some comments and questions in the final section 9.

Aknowledgements. I am most grateful to Gidi Amir and Balint Virag who helped me correct and improve a preliminary non-accurate version of this article. I also wish to thank Laurent Saloff-Coste, Alain Valette and Andrzej Zuk, who supported me during this work. Finally, I thank the anonymous referee for careful reading and many comments.

2. The groups involved

2.1. **Directed groups.** Given a sequence $\bar{d} = (d_j)_{j \geq 0}$ of integers $d_j \geq 2$, the spherically homogeneous rooted tree $T_{\bar{d}}$ is the graph with vertices $v = (i_1 i_2 \dots i_k)$ with i_j in $\{1, 2, \dots, d_{j-1}\}$, including the empty sequence \emptyset called the root, and edges $\{(i_1 i_2 \dots i_k), (i_1 i_2 \dots i_k i_{k+1})\}$. The index k is called the depth or level of v, denoted |v| = k.

The boundary $\partial T_{\bar{d}}$ of the tree $T_{\bar{d}}$ is the collection of infinite sequences $x = (i_1 i_2 \dots)$ with i_j in $\{1, 2, \dots, d_{j-1}\}$.

The group $Aut(T_{\bar{d}})$ of automorphisms of the rooted tree is the group of graph automorphisms that fix the root \emptyset . The following isomorphism is canonical:

(1)
$$Aut(T_{\bar{d}}) \simeq Aut(T_{\sigma\bar{d}}) \wr S_{d_0}.$$

The symbol \wr represents the permutational wreath product $G \wr S_d = (G \times \cdots \times G) \rtimes S_d$ where S_d acts on the direct product of d copies of G by permutation, and σ represents the shift on sequences, so that $\sigma \bar{d} = (d_1, d_2, \ldots)$. As the isomorphism (1) is canonical, we identify an element and its image and write:

$$(2) g = (g_1, \dots, g_{d_0})\sigma$$

with g in $Aut(T_{\bar{d}})$, the g_i in $Aut(T_{\sigma\bar{d}})$ and σ in S_{d_0} . The automorphism g_t represents the action of g on the subtree T_t , isomorphic to $T_{\sigma\bar{d}}$, hanging from vertex t, and the rooted component σ describes how these subtrees $(T_t)_{t=1...d_0}$ are permuted.

With notation (2), for any vertex ty in $T_{\bar{d}}$, one has $g(ty) = \sigma(t)g_t(y)$. If $f = (f_1, \ldots, f_{d_0})\tau$, then $(gf)(ty) = (f \circ g)(ty) = \tau(\sigma(t))f_{\sigma(t)}(g_t(y)) = (\sigma\tau)(t)(g_tf_{\sigma(t)})(y)$, so that $gf = (g_1f_{\sigma(1)}, \ldots, g_{d_0}f_{\sigma(d_0)})\sigma\tau$.

An automorphism g is rooted if $g = (1, ..., 1)\sigma$ for a permutation σ in S_{d_0} . The group of rooted automorphisms of $T_{\bar{d}}$ is obviously isomorphic to S_{d_0} and can be realised canonically as a subgroup or a quotient of $Aut(T_{\bar{d}})$.

The set $H_{\bar{d}}$ of automorphisms directed by the geodesic ray $1^{\infty} = (111...)$ in $T_{\bar{d}}$ is defined recursively. An element h is in $H_{\bar{d}}$ if there exists h' in $H_{\sigma\bar{d}}$ and $\sigma_2, \ldots, \sigma_{d_0}$ rooted in $Aut(T_{\sigma\bar{d}})$ such that:

$$(3) h = (h', \sigma_2, \dots, \sigma_{d_0}).$$

There is a canonical isomorphism of abstract groups: $H_{\bar{d}} \simeq S_{d_1} \times \cdots \times S_{d_1} \times H_{\sigma \bar{d}}$ with $d_0 - 1$ factors S_{d_1} . As a consequence, $H_{\bar{d}}$ is the uncountable but locally finite product:

$$(4) H_{\bar{d}} \simeq S_{d_1} \times \cdots \times S_{d_1} \times S_{d_2} \times \cdots \times S_{d_2} \times \cdots,$$

with $d_{l-1}-1$ factors S_{d_l} , indexed by $\{2,\ldots,d_{l-1}\}$. Under this isomorphism, denote: $h=(\sigma_{1,2},\ldots,\sigma_{1,d_0},\sigma_{2,2},\ldots,\sigma_{2,d_1},\ldots)$ with $\sigma_{k,t}$ in S_{d_k} .

The action of $h \in H_{\bar{d}}$ on the rooted tree $T_{\bar{d}}$ and its boundary $\partial T_{\bar{d}}$ is given by:

$$h(1^{k-1}ti_{k+1}i_{k+2}\dots) = 1^{k-1}t\sigma_{k,t}(i_{k+1})i_{k+2}\dots,$$

where each vertex or boundary element is uniquely written $1^{k-1}ti_{k+1}i_{k+2}...$ with t in $\{2, ..., d_{k-1}\}$ and $k \geq 1$ integer. Notation 1^k is a shortcut for 11...1 with k terms.

Definition 2.1. A group G of automorphisms of $T_{\bar{d}}$ is called *directed* when it admits a generating set of the form $S \cup H$ where S is included in the group S_{d_0} of rooted automorphisms and H is included in $H_{\bar{d}}$. Denote G = G(S, H) such a directed group. Then $G(S_{d_0}, H_{\bar{d}})$ is the (uncountable) full directed group of $T_{\bar{d}}$. Say a group G(S, H) is saturated if $S = S_{d_0}$ is the full group of rooted automorphism and H is a group such that the projection of the equidistribution measure over H onto each factor S_{d_l} in (4) is the equidistribution measure on S_{d_l} . The full directed group is obviously saturated.

Assume the sequence $\bar{d} = (d_i)_i$ is bounded taking finitely many values e_1, \ldots, e_T , then the direct product $H = S_{e_1} \times \cdots \times S_{e_T}$ embeds diagonaly into $S_{d_1} \times \cdots \times S_{d_1} \times \cdots \simeq H_{\bar{d}}$ (where the factors S_{e_t} embeds diagonaly into the subproduct of factors for which $d_l = e_t$). With the obvious identifications, the group $G(S_{d_0}, H)$ is a finitely generated saturated directed group. Note that this precise group is minimal among saturated directed groups of $T_{\bar{d}}$.

2.2. Extended directed groups. For a fixed x, the set $xT_d = \{x, x1, ..., xd\}$ has a structure of rooted tree with root x and one level $\{x1, ..., xd\}$. The extended boundary $E\partial T_{\bar{d}}$ of the rooted tree $T_{\bar{d}}$ is obtained by replacing each boundary point x by a short rooted tree xT_d :

$$E\partial T_{\bar{d}} = \{xT_d | x \in \partial T_{\bar{d}}\} = \{(x; x1, \dots, xd) | x \in \partial T_{\bar{d}}\}.$$

Call extended tree the set $ET_{\bar{d}} = T_{\bar{d}} \sqcup E\partial T_{\bar{d}}$. Its group of automorphisms is the group:

(5)
$$Aut(ET_{\bar{d}}) = S_d \wr_{\partial T_{\bar{d}}} Aut(T_{\bar{d}}) = \{ \varphi : \partial T_{\bar{d}} \to S_d \} \rtimes Aut(T_{\bar{d}}),$$

where the action of the group $Aut(T_{\bar{d}})$ on functions is given by $g.\varphi(x) = \varphi(gx)$ so that $(g_1g_2).\varphi = g_2.(g_1.\varphi)$. The group $Aut(ET_{\bar{d}})$ of automorphisms of extended tree was introduced by Bartholdi and Erschler in [BE] as "permutational wreath product over the boundary". The wreath product isomorphism (1) extends well:

Proposition 2.2. There is a canonical isomorphism:

$$Aut(ET_{\bar{d}}) \simeq Aut(ET_{\sigma\bar{d}}) \wr_{\{1,\dots,d_0\}} S_{d_0}.$$

Proof. Any γ in $Aut(ET_{\bar{d}})$ is decomposed $\gamma = \varphi g$, with $g \in Aut(T_{\bar{d}})$ and $\varphi : \partial T_{\bar{d}} \to S_d$. The classical isomorphism (1) provides a decomposition $g = (g_1, \ldots, g_d)\sigma$. Also the boundary of the tree can be decomposed into d_0 components $\partial T_{\bar{d}} = \partial T_1 \sqcup \cdots \sqcup \partial T_d$ with $T_t \simeq T_{\sigma \bar{d}}$ the tree descended from the first level vertex t. Set $\varphi_t = \varphi|_{\partial T_t}$ the restriction of φ . With these notations, the application Φ realising the canonical isomorphism is given by:

$$\Phi(\gamma) = (\varphi_1 g_1, \dots, \varphi_d g_d) \sigma \in Aut(ET_{\sigma \bar{d}}) \wr_{\{1,\dots,d_0\}} S_{d_0}.$$

In order to prove the proposition, it is sufficient to check that $\Phi(\gamma \gamma') = \Phi(\gamma)\Phi(\gamma')$.

On the one hand, $\gamma \gamma' = \varphi g \varphi' g' = \varphi(g.\varphi') g g' = \psi g g'$, with $\psi = \varphi(g.\varphi')$. As above set $\psi_t = \psi|_{\partial T_t}$, and as classically $g g' = (g_1 g'_{\sigma(1)}, \dots, g_d g'_{\sigma(d)}) \sigma \sigma'$, the embedding is:

$$\Phi(\gamma\gamma') = (\psi_1 g_1 g'_{\sigma(1)}, \dots, \psi_d g_d g'_{\sigma(d)}) \sigma\sigma'.$$

On the other hand:

$$\Phi(\gamma)\Phi(\gamma') = (\varphi_1 g_1, \dots, \varphi_d g_d)\sigma(\varphi'_1 g'_1, \dots, \varphi'_d g'_d)\sigma'
= (\varphi_1 g_1 \varphi'_{\sigma(1)} g'_{\sigma(1)}, \dots, \varphi_d g_d \varphi'_{\sigma(d)} g'_{\sigma(d)})\sigma\sigma'
= (\varphi_1 (g_1, \varphi'_{\sigma(1)}) g_1 g'_{\sigma(1)}, \dots, \varphi_d (g_d, \varphi'_{\sigma(d)}) g_d g'_{\sigma(d)})\sigma\sigma'$$

There remains to check $\psi_t = \varphi_t(g_t.\varphi'_{\sigma(t)})$, and indeed for any $y \in \partial T_t \simeq \partial T_{\sigma \bar{d}}$:

$$\psi_t(y) = \psi(ty) = (\varphi(g.\varphi'))(ty) = \varphi(ty)((g.\varphi')(ty)) = \varphi(ty)\varphi'(g.ty)$$
$$= \varphi(ty)\varphi'(\sigma(t)(g_t.y)) = \varphi_t(y)\varphi'_{\sigma(t)}(g_t.y) = \varphi_t(y)(g_t.\varphi'_{\sigma(t)})(y).$$

Functions over $\partial T_{\bar{d}}$ supported on $1^{\infty} = 1111...$ will play a specific role. For $f \in S_d$, denote $\varphi_f : \partial T_{\bar{d}} \to S_d$ the function $\varphi_f(1^{\infty}) = f$ and $\varphi_f(x) = id$ if $x \neq 1^{\infty}$. Note that for h in $H_{\bar{d}}$, one has $h\varphi_f = \varphi_f h$ in $Aut(ET_{\bar{d}})$ because h fixes the ray 1^{∞} .

Definition 2.3. A group Γ of automorphisms of the extended tree $ET_{\bar{d}}$ is called directed when it admits a generating set of the form $S \cup H \cup F$ where S is rooted, H is included in $H_{\bar{d}}$ and elements of F have the form φ_f for $f \in S_d$. Denote $\Gamma = \Gamma(S, H, F)$ such a directed group. Say a group $\Gamma(S_{d_0}, H, F) < Aut(ET_{\bar{d}})$ is saturated if $G(S_{d_0}, H)$ is saturated. Saturation implies that equidistribution on $H \times F$ projects to equidistribution on each factor S_{d_k} of (4) and on the factor F which justifies the notation $\Gamma(S_{d_0}, H \times F)$ for saturated directed groups.

Unless mentioned otherwise, use for the directed group $\Gamma(S, H, F)$ the set $S \cup H \times F$, where both S and $H \times F$ are finite groups themselves (hence symmetric) in the case of finitely generated saturated directed groups.

Denote H_l the restriction of $H < H_{\bar{d}}$ to levels $\geq l$, i.e. the projection of H to $H_{\sigma^l\bar{d}} = S_{d_{l+1}} \times \cdots \times S_{d_{l+1}} \times \cdots$. Also denote S_l the subgroup of S_{d_l} generated by the projections of H on the $d_{l-1} - 1$ factors S_{d_l} of (4).

Proposition 2.4. Let $\Gamma(S, H, F) < Aut(ET_{\bar{d}})$ be a directed group. There is a canonical embedding:

$$\Gamma(S, H, F) \hookrightarrow \Gamma(S_1, H_1, F) \wr S.$$

More generally:

$$\Gamma(S_l, H_l, F) \hookrightarrow \Gamma(S_{l+1}, H_{l+1}, F) \wr S_l$$
.

Proof. The embedding is clear from proposition 2.2 as the image of generators is given by s = (1, ..., 1)s, $h = (h', \sigma_2, ..., \sigma_d)$ and $\varphi_f = (\varphi_f, 1, ..., 1)$.

Observe that if $\Gamma(S, HF)$ is saturated, then $\Gamma(S_l, H_lF)$ is saturated for all l.

2.3. **Examples.** The class of directed groups of $Aut(T_{\bar{d}})$ contains many examples of groups that have been widely studied in relation with torsion ([Ale],[Gri1],[Gri2], [GS],[BS]), intermediate growth ([Gri1],[Gri2],[BS],[Ers3]), subgroup growth ([Seg]), non-uniform exponential growth ([Wil1],[Wil2],[Bri2]) and amenability ([GZ],[BV], [Bri2],[BKN],[AAV],[AV]).

Theorem 3.6 in [Bri2] states that the full directed group $G(S_{d_0}, H_{\bar{d}})$ is amenable if and only if the sequence \bar{d} is bounded. This result obviously extends to the setting of automorphisms of extended trees. Indeed, the group $\Gamma(S, H, F)$ is a subgroup of $F\wr_{\partial T_{\bar{d}}}G(S, H)$ which is a group extension of a direct sum of finite groups (copies of F) by the group G(S, H), hence is amenable when valency \bar{d} is bounded. On the other hand, G(S, H) is a quotient of $\Gamma(S, H, F)$, so the latter inherits non-amenability when \bar{d} is unbounded.

The notion of automorphisms of extended trees is a reformulation of permutational wreath product, which were introduced in [BE] in order to compute explicit intermediate growth functions (see also [Bri3]). The boundary extension T_d does not need to be a finite tree, i.e. the permutationnal wreath product $F \wr_{\partial T_{\bar{d}}}$ makes sense for any group F. This was used in [BE] to make a stack of extensions of trees and compute the growth function of some finitely generated groups of their automorphisms, namely $b_k(r) \approx e^{r^{\alpha_k}}$, where $\alpha_k \to 1$ with the number k of extensions in the stack.

2.4. Non-uniform growth and Haagerup property. This paragraph illustrates the interest of extended trees by an application that will not be used further in the rest of the article. It can be omitted at first reading.

A group is said to have the Haagerup property if it admits a proper continuous affine action on a Hilbert space (see [CCJJV]). This is for instance the case for free groups and amenable groups. Groups with the Haagerup property have attracted interest as they are known to satisfy the Baum-Connes conjecture.

Denote $A_d < S_d$ the group of alternate permutations, and given a bounded sequence \bar{d} of integers $d \leq d_i \leq D$, define the finite subgroup $A_{\bar{d}} < H_{\bar{d}} < Aut(T_{\bar{d}})$ by the following:

- (1) abstractly $A_{\bar{d}} \simeq \mathcal{A}_d \times \cdots \times \mathcal{A}_D$ with projection on factor d' denoted $pr_{d'}$,
- (2) an element $b \in A_{\bar{d}}$ is realised in $Aut(T_{\bar{d}})$ according to the recursive rule $b = (b', pr_{d_1}(b), 1, \dots, 1)$ for $b' \in A_{\sigma \bar{d}} < Aut(T_{\sigma \bar{d}})$.

Take F to be the free product $\mathcal{A}_5 * \mathcal{A}_5$, and consider the group $\Gamma(\mathcal{A}_{d_0}, A_{\bar{d}}, \mathcal{A}_5 * \mathcal{A}_5) < Aut(ET_{\bar{d}})$ which is a directed group of automorphisms of an extended tree (with infinite extension at the boundary so that $\mathcal{A}_5 * \mathcal{A}_5 < S_{\infty}$).

Proposition 2.5. If $29 \le d_i \le D$, the group $\Gamma(A_{d_0}, A_{\bar{d}}, A_5 * A_5) < Aut(ET_{\bar{d}})$ has non-uniform exponential growth, is non-amenable and has Haagerup property.

Directed groups of the form $G(\mathcal{A}_{\bar{d}}, A_{\bar{d}}) < Aut(T_{\bar{d}})$ were introduced by Wilson as the first examples of groups with non-uniform exponential growth ([Wil2],[Wil1]). For bounded valency and $A_{\bar{d}}$ finite as above these groups are amenable ([Bri2]) hence have Haagerup property. For unbounded valency and $A_{\bar{d}} \simeq \mathcal{A}_5 * \mathcal{A}_5$, these groups are

non-amenable. Zuk asked whether they still have Haagerup property. The groups $\Gamma(\mathcal{A}_{d_0}, A_{\bar{d}}, \mathcal{A}_5 * \mathcal{A}_5)$ resemble the groups $G(\mathcal{A}_{\bar{d}}, A_{\bar{d}})$ for unbounded sequences \bar{d} in the sense that the "free product factor" $\mathcal{A}_5 * \mathcal{A}_5$ is "located at the boundary of the tree". Thus the proposition above hints that Wilson groups $G(\mathcal{A}_{\bar{d}}, A_{\bar{d}})$ of [Wil2] have Haagerup property (but it does not prove it). It provides the first example of groups of non-uniform growth for which Haagerup property does not follow from amenability. No example of group having non-uniform growth but not Haagerup property is known.

Proof. Each of the groups \mathcal{A}_{d_0} , $A_{\bar{d}}$ and $\mathcal{A}_5 * \mathcal{A}_5$ is perfect and generated by finitely many involutions (the number of which depends only on D), so that this is also the case for $\Gamma(\mathcal{A}_{d_0}, A_{\bar{d}}, \mathcal{A}_5 * \mathcal{A}_5)$. Moreover, there is an isomorphism:

$$\Gamma(\mathcal{A}_{d_0}, A_{\bar{d}}, \mathcal{A}_5 * \mathcal{A}_5) \simeq \Gamma(\mathcal{A}_{d_1}, A_{\sigma\bar{d}}, \mathcal{A}_5 * \mathcal{A}_5) \wr \mathcal{A}_{d_0}.$$

(Indeed by proposition 7.2 in [Bri2], Γ contains any commutator ($[b_1, b_2], 1, \ldots, 1$) for b_1, b_2 a pair of elements in one of the generating groups $\mathcal{A}_{d_0}, \mathcal{A}_{\bar{d}}, \mathcal{A}_5 * \mathcal{A}_5$. By perfection, it shows that the embedding of proposition 2.4 is onto.)

This shows that $\Gamma(\mathcal{A}_{d_0}, A_{\bar{d}}, \mathcal{A}_5 * \mathcal{A}_5)$ belongs to a class χ (see [Wil2]) and there exists generating sets with growth exponent tending to 1. Also, the group contains a non-trivial free product, hence is non amenable, and has non-uniform exponential growth.

The group $\Gamma(\mathcal{A}_{d_0}, A_{\bar{d}}, \mathcal{A}_5 * \mathcal{A}_5)$ inherits Haagerup property for it is contained in $(\mathcal{A}_5 * \mathcal{A}_5) \wr_{\partial T_{\bar{d}}} G(\mathcal{A}_{d_0}, A_{\bar{d}})$ which itself has Haagerup property. Indeed, there is a short exact sequence:

$$1 \to (\mathcal{A}_5 * \mathcal{A}_5)^{\partial T_{\bar{d}}} \to (\mathcal{A}_5 * \mathcal{A}_5) \wr_{\partial T_{\bar{d}}} G(\mathcal{A}_{d_0}, A_{\bar{d}}) \to G(\mathcal{A}_{d_0}, A_{\bar{d}}) \to 1,$$

where $(A_5*A_5)^{\partial T_{\bar{d}}}$ has Haagerup property and $G(A_{d_0}, A_{\bar{d}})$ is amenable, which implies Haagerup property for the middle term (example 6.1.6 in [CCJJV]).

In the remaining sections of the present article (as well as in [Bri3]), only finite groups F are considered, which simplifies some arguments and still permits to observe various phenomena.

3. Rewriting process and activity of words

3.1. **Rewriting process.** Given a finitely generated saturated (extended) directed group $\Gamma(S_{d_0}, HF)$ acting on a tree $T_{\bar{d}}$ of bounded valency, a canonical generating set is $S_{d_0} \sqcup FH$. Note that by saturation and bounded valency, both S_{d_0} and HF are finite subgroups of $\Gamma(S_{d_0}, HF)$. In the present article, the disjoint union of these subgroups is taken as generating set. In particular both $1_{S_{d_0}}$ and 1_{HF} are generators, distinct in a "word perspective".

Definition 3.1. Given the generating set $S_{d_0} \sqcup FH$ of a finitely generated saturated group $\Gamma(S_{d_0}, HF)$, a representative word is *alternate* if it has the form $w = s_1 k_1 s_2 k_2 \ldots s_n k_n s_{n+1}$ for some s_i in S_{d_0} and k_i in HF.

Note that any word w in this generating set admits a canonical alternate form obtained by merging packs of successive terms belonging to the same finite group S_{d_0} or HF. For example, the canonical alternate form of $s_1s_2k_3k_3^{-1}1_{S_{d_0}}k_41_{HF}$ is $(s_1s_2)1_{FH}1_{S_{d_0}}k_4$.

The alternate form $w = s_1 k_1 \dots s_n k_n s_{n+1}$ of a word in the generating set $S_d \sqcup FH$ is equivalent to $w = k_1^{\sigma_1} \dots k_n^{\sigma_n} \sigma_{n+1}$, for $\sigma_i = s_1 \dots s_i$, where $k^{\sigma} = \sigma k \sigma^{-1}$ denotes the conjugate of k by σ . Remind that k_j in HF is uniquely decomposed into $k_j = (k'_j, b_{j,2}, \dots, b_{j,d_0})$ with k'_j in H_1F and $b_{j,t}$ in S_{d_1} rooted.

Proposition 3.2 (Rewriting process). With the notations above, the alternate word $w = s_1 k_1 \dots s_n k_n s_{n+1}$ can be algorithmically rewritten in the wreath product as

$$w = (w^1, \dots, w^{d_0})\sigma_{\emptyset}$$

where:

- (1) the rooted permutation is $\sigma_{\emptyset} = s_1 s_2 \dots s_{n+1} = \sigma_{n+1}$,
- (2) the word $w^t = s_1^t k_1^t \dots s_{m_t}^t k_{m_t}^t s_{m_t+1}^t$ is alternate in the generating system $S_{d_1} \sqcup H_1 F$ of the saturated directed group $\Gamma(S_{d_1}, H_1 F)$, of length $m_t \leq \frac{n+1}{2}$ such that $m_1 + \dots + m_{d_0} \leq n$,
- (3) the factors s_i^t depend only on factors $b_{j,t'}$ at times j when $\sigma_j(t) = t'$,
- (4) the factors k_i^t depend only on factors k_j' at times j when $\sigma_j(t) = 1$.

Proof. The factor $k_i^{\sigma_j} = \sigma_j k_j \sigma_j^{-1}$ has image in the wreath product:

$$k_j^{\sigma_j} = (b_{j,\sigma_j(1)}, \dots, k_j', \dots, b_{j,\sigma_j(d_0)}),$$

with k'_j in position $\sigma_j^{-1}(1)$. This shows that w^t in the wreath product image of w is a product of terms $b_{j,\sigma_j(t)}$ at times j when $\sigma_j(t) \neq 1$ and terms k'_j at times j when $\sigma_j(t) = 1$. The word w^t is the canonical alternate form of this product. Each factor k_j furnishes a factor k'_j to exactly one of the coordinates, so the sum of length is $\leq n$. Also by looking at alternate forms, there has to be a factor $k_j^{\sigma_j}$ such that $\sigma_j(t) \neq 1$, giving $b_{j,t'}$ on coordinate t, between two factors such that $\sigma_j(t) = 1$, giving k'_j on coordinate t, so that the length of w^t is at most half the length of w.

Of course, the rewriting process can be iterated, and at each vertex v = ut is associated to the alternate word w another alternate word w^v of length m_v in the generating set $S_{d_{|v|}} \sqcup H_{|v|}F$ of the saturated directed group $\Gamma(S_{d_{|v|}}, H_{|v|})$ defined inductively by $w^v = w^{ut} = (w^u)^t$. Note that the word w^v has length $m_v \leq \frac{n}{2^{|v|}} + 1$, and $\sum_{|v|=l} m_v \leq n$.

3.2. Minimal tree and activity.

Definition 3.3. The minimal tree T(w) of the alternate word w in $S_{d_0} \sqcup HF$ is the minimal regular rooted subtree of $T_{\bar{d}}$ such that $m_v \leq 1$ for any leaf v of T(w). Recall that a subtree T of $T_{\bar{d}}$ is regular if whenever a vertex v belongs to T, either all its descendants vt also belong to T, or none of them does.

The minimal tree T(w) is constructed algorithmically. Indeed it contains the root \emptyset , and if v is in T(w), either $m_v \leq 1$ and no descendant of v belongs to T(w), or




FIGURE 1. Description of the action of a word w via the minimal tree T(w)

 $m_v \geq 2$ and all the sons of v belong to T(w). As m_v decays exponentially fast with generations, the minimal tree T(w) has depth at most $\log_2 n$.

The leaves of the minimal tree T(w) satisfy either $m_v = 0$ in which case they are called *inactive*, or $m_v = 1$ called *active* leaves. The subset A(w) of the boundary (set of leaves) $\partial T(w)$ of the minimal tree T(w) is called the *active* set of the word w. Its size is the activity a(w) = #A(w) of the word w. The activity of a word $w = s_1$ of length 0 (no factor in HF) is a(s) = 0.

Remark 3.4. The minimal tree T(w), as well as the set of active leaves A(w), of a word $w = s_1 k_1 \dots k_n s_{n+1}$ depend only on the word $s_1 k_1 \dots k_n s_{n+1}$ where $k_j = \varphi_{f_i} k_j$, i.e. on the quotient G(S, H) of $\Gamma(S, HF)$.

The notions of minimal tree and active set allow an interesting description of the action of a word w on the tree $T_{\bar{d}}$. Indeed, the action of the automorphism $\gamma =_{\Gamma} w$ in $Aut(ET_{\bar{d}})$ is determined by the following data:

- (1) the minimal tree T(w) and its active set A(w),
- (2) the permutations $\sigma_u \in S_{d_{|u|}}$ attached to vertices $u \in T(w) \backslash A(w)$, (3) the "short words" $w^v = s_1^v k^v s_2^v$ attached to active vertices v.

The description of the short word $w^v = s_1^v k^v s_2^v$ at an active vertex v can be refined into a tree action $s_1^v h_v s_2^v$, together with a boundary function given by $\varphi_{f^v}(x) = f^v$ for $x = v(s_1^v(1))1^{\infty}$ and $\varphi_{f_v}(x) = 1$ otherwise.

A point x in $\partial T_{\bar{d}}$ is called active when it is of the form $x = v(s_1^v(1))1^{\infty}$ for some active leaf v of T(w). If the element $\gamma = w$ in $\Gamma(S, HF)$ has the form $\gamma = \varphi g$ in the permutational wreath product (5), then the support of φ is included in the set $A_{\partial}(w)$ of active boundary points.

3.3. Ascendance forest. Given an alternate word w in $\Gamma(S_{d_0}, HF)$ and the collection $(w^v)_{v \in T(w)}$ of words obtained by rewriting process, each word w^v is a product of factors $w^v = s_1^v k_1^v \dots k_{m_v}^v s_{m_v+1}^v$. Recall that each factor k_i^{vt} in the word w^{vt} is a product of terms $k_j^{\prime v}$, obtained from $(k_j^v)^{\sigma_j^v} = (b_{j,\sigma_j^v(1)}^v, \dots, k_j^{\prime v}, \dots, b_{j,\sigma_j^v(d_{|v|})}^v)$ with $k_j^{\prime v}$ in position t, during the rewriting process of the word w^v for the vertex v, into $w^v = (w^{v1}, \ldots, w^{vd_{|v|}})\sigma_v$.

Consider the graph AF(w) with set of vertices the collection $(k_i^v)_{v,i}$ of factors in $H_{|v|}F$ appearing in the rewritten words $(w^v)_{v \in T(w)}$, where a pair of factors $(k_i^v, k_j^{v'})$ is linked by an edge when v' = vt and the term k_i^{v} appears in k_j^{vt} .

Fact 3.5. The graph AF(w) is a forest, i.e. a graph with no loop. More precisely, AF(w) is a union of trees $(\tau_v)_{v \in A(w)}$ indexed by the active set of w. Moreover, if v is an active leaf, then

$$k^v = \prod_{k_i \in \partial \tau_v} k_j^{(|v|)}$$

is the product ordered with j, where $k_j^{(|v|)}$ is the restriction of $k_j \in HF$ to $H_{|v|}F$. In particular:

$$f_v = \prod_{k_j \in \partial \tau_v} f_j.$$

Proof. Let $v = t_1 \dots t_l$ be an active leaf with $w^v = s_1^v k^v s_2^v$. The term k^v is a product of terms $k_j^{\prime t_1 \dots t_{l-1}}$ in $w^{t_1 \dots t_{l-1}}$ (for those j's where $\sigma_j^{t_1 \dots t_{l-1}}(1) = t_l$), which are the neighbours of the vertex k^v in the graph AF(w). This describes the ball of center k^v and radius 1.

Inductively, each factor $k_i^{t_1...t_{l-r}}$ which represents a vertex in the sphere of center k^v and radius r is a product of terms $k_j^{t_1...t_{l-r-1}}$ in $w^{t_1...t_{l-r-1}}$ (for those j's where $\sigma_j^{t_1...t_{l-r}}(1) = t_{l-r}$), which form the link of $k_i^{t_1...t_{l-r}}$ in the sphere of radius r+1.

Now each factor $k_j^{t_1...t_{\lambda}}$ in a word of level λ contributes to exactly one factor $k_i^{t_1...t_{\lambda}t_{\lambda+1}}$, which rules out the possibility of a loop.

So the connected component of k^v in the graph AF(w) is a tree τ_v , it is also the l-ball of center k^v and the leaves of τ_v form the l-sphere, which is precisely the set of factors k_j of w that lie in τ_v . By construction of AF(w), k^v is the required ordered product.

Remark 3.6. Observe that the ascendance forest AF(w) of a word $w = s_1k_1 \dots k_ns_{n+1}$ depends only on the word $s_1h_1 \dots h_ns_{n+1}$ for $k_j = \varphi_{f_j}h_j$. Indeed given a factor $k_i^{t_1\dots t_\lambda}$, its link to factors in level $\lambda+1$ is determined by the factor $\sigma_i^{t_1\dots t_\lambda}$, which is determined by the factors $s_1^{t_1\dots t_\lambda}, \dots s_i^{t_1\dots t_\lambda}$ themselves determined by the sequence $(h_j^{t_1\dots t_{\lambda-1}})_j$. Thus the link of k_i^v does not depend on the sequence $(f_j)_{j\in\{1,\dots,n\}}$. A consequence of this observation and the preceding fact is that for any function φ with support included in $A_{\partial}(w)$, there exists $(f_j)_{j\in\{1,\dots,n\}}$ such that $\varphi s_1h_1\dots h_ns_{n+1} = r_1 \varphi_{f_1}h_1\dots \varphi_{f_n}h_ns_{n+1}$. Moreover the number of such n-tuples is independent of the function φ . This shows:

Fact 3.7. Given $g = s_1 h_1 \dots h_n s_{n+1}$, for any $\varphi : \partial T_{\bar{d}} \to F$ with support in $A_{\partial}(g)$, one has:

$$\#\{f_1,\ldots,f_n\in F|\varphi g=_\Gamma s_1\varphi_{f_1}h_1\ldots\varphi_{f_n}h_ns_{n+1}\}=\left(\frac{1}{\#F}\right)^{\#A_{\partial}(g)}.$$

3.4. Counting activity. The inverted orbit of a product $w = r_1 \dots r_k$ of automorphisms $r_i \in Aut(T_{\bar{d}})$ is the set $\mathcal{O}(w) = \{1^{\infty}, r_k 1^{\infty}, r_{k-1} r_k 1^{\infty}, \dots, r_1 \dots r_k 1^{\infty}\}$ (see [BE]). The inverted orbit of w^{-1} coincides with the activity set $A_{\partial}(w)$ defined in section 3.2. The notion of activity is classical in the context of automorphisms of rooted trees (see [Nek],[Sid]).

Proposition 3.8. On a finitely generated saturated directed group $\Gamma(S_{d_0}, HF)$ acting on a tree of bounded valency \bar{d} , the activity function a(w) which for $w = s_1\varphi_{f_1}h_1\ldots\varphi_{f_n}h_ns_{n+1}$ counts equivalently:

- (1) the size of the set A(w) of active leaves in the minimal tree $T(w) \subset T_{\bar{d}}$,
- (2) the size of the set $A_{\partial}(w)$ of active boundary points in $\partial T_{\bar{d}}$,
- (3) the number of trees (i.e. connected components) in the ascendance forest AF(w),
- (4) the size of the inverted orbit $\mathcal{O}(s_{n+1}^{-1}h_n^{-1}\dots h_1^{-1}s_1^{-1})$ in the sense of [BE],

satisfies under rewriting process $w = (w^1, \dots, w^{d_0})\sigma_{\emptyset}$:

$$a(w) = a(w^1) + \dots + a(w^{d_0}).$$

Moreover, the constraint $a(w) \leq r$ only permits to describe at most exponentially many elements in $\Gamma(S_{d_0}, HF)$, i.e. there exists a constant C depending only on $D = \max\{d_i\}$ and #HF such that:

$$\#\{\gamma \in \Gamma(S_{d_0}, HF) | \exists w =_{\Gamma} \gamma, a(w) \leq r\} \leq C^r.$$

Note that a(w) is the activity for words in the group $\Gamma(S_{d_0}, HF)$ and $a(w^t)$ is the activity function for words in the group $\Gamma(S_{d_1}, H_1F)$.

Proof. Points (1), (2), (3) are clear from the descriptions above. Point (4) is shown by induction on n. Suppose $s_1k_1 \ldots k_{n-1}s_n = \varphi_n g_n$ for $g_n = s_1h_1 \ldots h_{n-1}s_n$, then $s_1k_1 \ldots k_{n-1}s_n\varphi_{f_n}h_n = \varphi_n(g_n.\varphi_{f_n})g_nh_n$ and the point $g_n^{-1}(1^{\infty})$ which is the support of the function $g_n.\varphi_{f_n}$ is added to the set of active leaves $A_{\partial}(w)$.

The equality on activities under rewriting process is trivial when $n \leq 1$, and if $n \geq 2$ then the minimal tree is not restricted to the root, one has $T(w) = T(w^1) \sqcup \cdots \sqcup T(w^{d_0}) \sqcup \{\emptyset\}$, so that $A(w) = A(w^1) \sqcup \cdots \sqcup A(w^{d_0})$, and equality holds.

In order to prove the exponential bound, first observe that $\#\partial T(w) \leq Da(w)$. It is obvious if $\partial T(w)$ is the root $\{\emptyset\}$ or the first level $\{1,\ldots,d_0\}$, and then clear for arbitrary T(w) by induction on the size #T(w) since $\partial T(w) = \partial T(w^1) \sqcup \cdots \sqcup \partial T(w^{d_0})$.

Now if $a(w) \leq r$, its minimal tree T(w) has a boundary of size $\leq Dr$, so there are $\leq K^r$ possibilities for T(w) (for some K depending only on D). To finish the description of $\gamma =_{\Gamma} w$, one has to choose permutations σ_u at vertices $u \in T(w) \setminus \partial T(w)$, for which there are $\leq D!^{Dr}$ possibilities, and short words $s_1^v k_v s_2^v$ at leaves $v \in \partial T(w)$, for which there are $\leq (D!^2 \# HF)^{Dr}$ possibilities.

4. Random Walks

Given a finitely generated saturated directed group $\Gamma(S_{d_0}, HF)$, consider random alternate words $Y_n = s_1 k_1 s_2 k_2 \dots s_n k_n s_{n+1}$, where s_i in S_{d_0} and k_i in HF are equidistributed and all factors are independent. Such a random alternate product Y_n is the simple random walk on $\Gamma(S_{d_0}, HF)$ for the symmetric generating set $S_{d_0}HFS_{d_0}$ (for the product of two independent equidistributed variables $s_i's_{i+1}$ in S_{d_0} is another equidistributed variable, independent of other factors).

- 4.1. Inheritance of random process through wreath product. Given a random alternate word Y_n , the rewriting process of proposition 3.2 furnishes an image in the wreath product $Y_n = (Y_n^1, \dots, Y_n^{d_0})\tau_n$. Each coordinate Y_n^t is a random process on words in $S_{d_1} \sqcup H_1 F$ by proposition 2.4, which turns out to be a random alternate word of random length.
- **Lemma 4.1.** Denote $Y_n = (Y_n^1, \dots, Y_n^{d_0})\tau_n$ the alternate words obtained by rewriting process of a random alternate word $Y_n = s_1k_1s_2k_2\dots s_nk_ns_{n+1}$ in a finitely generated saturated directed group $\Gamma(S_{d_0}, HF)$. Then:
 - (1) The rooted random permutation is $\tau_n = s_1 \dots s_{n+1}$ hence is equidistributed.
 - (2) The random length m_t of the random product Y_n^t has the law of the sum of n independent Bernoulli variables (u_j) on $\{0,1\}$ with $P(u_j=1)=p_0=\frac{d_0-1}{d_0^2}$.

 In particular, by the law of large numbers $m_t \sim p_0 n$ almost surely, and by the principle of large deviations, for any $\theta > 0$, there exists $c_{\theta} < 1$ such that

$$P\left(\frac{m_t}{n}\notin[p_0-\theta,p_0+\theta]\right)\leq c_\theta^n.$$

(3) For each coordinate t, the conditionned variable $Y_n^t|m_t$ has precisely the law of the random alternate word Y'_{m_t} of length m_t in $S_{d_1} \sqcup H_1F$, i.e.

$$Y_n^t = Y_{m_t}' = s_1^t k_1^t \dots k_{m_t}^t s_{m_t+1}^t$$

where the factors s_j^t and k_j^t are equidistributed in S_{d_1} and H_1F respectively (except s_1^t and $s_{m_t+1}^t$), and all factors $(s_j^t, k_j^t)_j$ are independent.

This lemma is a restating of lemma 4.6 in [Bri1]. A brief proof is given below.

Proof. As in proposition 3.2, write $Y_n = k_1^{\sigma_1} \dots k_n^{\sigma_n} \sigma_{n+1}$, where each factor has image in the wreath product $k_j^{\sigma_j} = (b_{j,\sigma_j(1)}, \dots, k_j', \dots, b_{j,\sigma_j(d_0)})$ with $k_j' \in H_1F$ in position $\sigma_j(t)$ and $b_{j,s} \in S_{d_1}$. As $(\sigma_j)_{j=1}^n$ is a sequence of independent terms equidistributed in S_{d_0} (for $\sigma_j = s_1 \dots s_j$ with $(s_i)_{i=1}^n$ is a sequence of independent terms equidistributed in S_{d_0}), the position sequence $(\sigma_j(t))_{j=1}^n$ is equidistributed in $\{1, \dots, d_0\}$ for any choice of t.

Then for a fixed t, Y_n^t is a product of n terms which are either $b_{j,\sigma_j(t)}$ at times j when $\sigma_j(t) \neq 1$, which happens with probability $\frac{d_0-1}{d_0}$, or k'_j at times j when $\sigma_j(t) = 1$, which happens with probability $\frac{1}{d_0}$. In both cases, the factors are equidistributed in S_{d_1} or H_1F respectively because k_j is equidistributed in HF by saturation. Moreover, all the terms are independent.

Now to obtain an alternate word, the runs of successive terms that belong to the same finite group (either S_{d_1} or H_1F) are merged, the factors $(s_j^t, k_j^t)_j$ are still equidistributed and independent.

There remains to count the number of such runs, given by:

$$2m_t + 1 = 1 + \sum_{j=1}^{n} 1_{\{(\sigma_j(t)=1 \text{ and } \sigma_{j+1}(t)\neq 1) \text{ or } (\sigma_j(t)\neq 1 \text{ and } \sigma_{j+1}(t)=1)\}}.$$

Knowing that $P(\sigma_j(t)=1)=\frac{1}{d_0}$ and $P(\sigma_j(t)\neq 1)=\frac{d_0-1}{d_0}$ independently of previous terms, two successive terms belong to different finite groups with probability $2\frac{d_0-1}{d_0}\frac{1}{d_0}=2p_0$.

Note that m_t depends only on $\sigma_1, \ldots, \sigma_n$, i.e. on s_1, \ldots, s_n , whereas the factors (s_j^t, k_j^t) are determined by k_1, \ldots, k_n . In particular, fixing $\sigma_1, \ldots, \sigma_n$ (hence m_t), and playing with k_1, \ldots, k_n any alternate word Y_n^t of length m_t in $S_{d_1} \sqcup H_1F$ appears with the same probability.

The lemma can be iterated to show that for any vertex v in $T_{\bar{d}}$, the random product Y_n^v obtained by rewriting process of the random alternate word Y_n is also an alternate random word in the group $\Gamma(S_{d_{|v|}}, H_{|v|}F)$, of length $m_v \sim p_0 \dots p_{|v|}n$ almost surely, i.e. the conditioned variable $Y_n^v|m_v = Y_{m_v}^{(|v|)}$ is a random alternate word in $\Gamma(S_{d_{|v|}}, H_{|v|}F)$ of length m_v .

4.2. Exponent sequence associated to valency sequence. Given a bounded sequence $\bar{d} = (d_i)_i$ of integers ≥ 2 , define $\bar{p} = (p_i)_i$ by $p_i = \frac{d_i - 1}{d_i^2}$. Note $d = \min(d_i)$, $D = \max(d_i)$, $p = \max(p_i) = \frac{d - 1}{d^2}$ and $P = \min(p_i) = \frac{D - 1}{D^2}$. Define the exponent function $\beta(n)$ associated to the valency sequence \bar{d} by:

$$k(n) = k_{\bar{d}}(n) = \min\{k | p_0 \dots p_k n \le 1\} \text{ and } \beta(n) = \beta_{\bar{d}}(n) = \frac{\log(d_0 \dots d_{k(n)})}{\log n}.$$

Moreover, given a small $\theta \neq 0$ and an integer N_0 depending only on θ , set:

$$k^{\theta}(n) = \min\{k | (p_0 + \theta) \dots (p_k + \theta)n \le N_0\} \text{ and } \beta^{\theta}(n) = \frac{\log(d_0 \dots d_{k^{\theta}(n)})}{\log n}.$$

For $N_0 = 1$, the function $k^{\theta}(n)$ is increasing with θ , and $\beta^{\theta}(n) \longrightarrow_{\theta \to 0} \beta(n)$ for a fixed n.

Proposition 4.2. There exists a function $\varepsilon(\theta) \longrightarrow_{\theta \to 0} 0$ such that for all n large enough (depending on θ):

$$|\beta(n) - \beta^{\theta}(n)| \le \varepsilon(\theta).$$

Proof. Assume $\theta > 0$ (similar proof for $\theta < 0$). For n large enough, $k^{\theta}(n) \geq k(n)$, and the difference is:

 $|\beta(n) - \beta^{\theta}(n)| = |\log_n(d_0 \dots d_{k(n)}) - \log_n(d_0 \dots d_{k^{\theta}(n)})| = \log_n(d_{k(n)+1} \dots d_{k^{\theta}(n)}),$ hence is bounded by:

$$|\beta(n) - \beta^{\theta}(n)| \le |k^{\theta}(n) - k(n)| \frac{\log D}{\log n}.$$

By definition of k(n) and $k^{\theta}(n)$, there are inequalities:

$$\frac{N_0}{n} \ge (p_0 + \theta) \dots (p_{k^{\theta}(n)} + \theta) > \frac{N_0(p_{k^{\theta}(n)} + \theta)}{n} \ge p_0 \dots p_{k(n)} N_0(P + \theta),$$

so that:

$$(p+\theta)^{k^{\theta}(n)-k(n)} \geq (p_{k(n)+1}+\theta)\dots(p_{k^{\theta}(n)}+\theta) = \frac{(p_0+\theta)\dots(p_{k^{\theta}(n)}+\theta)}{(p_0+\theta)\dots(p_{k(n)}+\theta)}$$

$$\geq \left(\frac{p_0}{p_0+\theta}\right)\dots\left(\frac{p_{k(n)}}{p_{k(n)}+\theta}\right)N_0(P+\theta)$$

$$\geq \left(\frac{P}{P+\theta}\right)^{k(n)}N_0(P+\theta),$$

which shows that for some constant K:

$$|k^{\theta}(n) - k(n)| \le k(n) \left| \frac{\log(\frac{P}{P+\theta})}{\log(p+\theta)} \right| + K.$$

Notice that $\frac{\log n}{|\log P|} \le k(n) \le \frac{\log n}{|\log p|}$ to finally obtain:

$$|\beta^{\theta}(n) - \beta(n)| \le \frac{\log D}{|\log p|} \left| \frac{\log(\frac{P}{P+\theta})}{\log(p+\theta)} \right| + \frac{K'}{\log n}.$$

The proposition follows from the limit $|\log(\frac{P}{P+\theta})| \longrightarrow_{\theta \to 0} 0$.

Fact 4.3. There is a bound on $\beta(n)$ depending only on the bounds $d \leq d_i \leq D$ on the valency of the rooted tree $T_{\bar{d}}$. Namely for some constant C depending only on D:

$$\beta_d \le \beta(n) \le \beta_D + \frac{C}{\log n},$$

where
$$\beta_d = \frac{\log d}{-\log p} = \frac{\log d}{\log(\frac{d^2}{d-1})} = \frac{1}{1 + \frac{\log \frac{d}{d-1}}{\log d}} = \frac{1}{2 - \frac{\log(d-1)}{\log d}}.$$

Note that $\beta_2 = \frac{1}{2}$ and $\beta_d \longrightarrow_{d \to \infty} 1$.

Proof. Note that:

$$\beta(n) = \frac{\log(d_0 \dots d_{k(n)})}{\log n} \ge \frac{\log(d_0 \dots d_{k(n)})}{-\log(p_0 \dots p_{k(n)})},$$

$$\beta(n) \le \frac{\log(d_0 \dots d_{k(n)})}{-\log(p_0 \dots p_{k(n)}) + \log p_{k(n)}} \le \frac{\log(d_0 \dots d_{k(n)})}{-\log(p_0 \dots p_{k(n)})} + \frac{C}{\log n}.$$

As $p_i = \frac{d_i - 1}{d_i^2}$, simply compute:

$$\frac{\log(d_0 \dots d_{k(n)})}{\log(\frac{d_0^2}{d_0 - 1} \dots \frac{d_{k(n)}^2}{d_{k(n)} - 1})} = \frac{1}{1 + \frac{\log(\frac{d_0}{d_0 - 1}) + \dots + \log(\frac{d_{k(n)}}{d_{k(n)} - 1})}{\log d_0 + \dots + \log d_{k(n)}}} \le \frac{1}{1 + \frac{\log(\frac{D}{D - 1})}{\log D}},$$

and similarly for the lower bound, by the inequality on ratio of average:

$$\frac{\log(\frac{D}{D-1})}{\log D} \le \frac{\log(\frac{d_0}{d_0-1}) + \dots + \log(\frac{d_{k(n)}}{d_{k(n)}-1})}{\log d_0 + \dots + \log d_{k(n)}} \le \frac{\log(\frac{d}{d-1})}{\log d}.$$

For a real number x > 0, define $h_{\bar{d}}(x) = d_0 \dots d_{k(x)}$ for the unique integer k(x) such that $\frac{1}{p_0} \dots \frac{1}{p_{k(x)}} \ge x > \frac{1}{p_0} \dots \frac{1}{p_{k(x)-1}}$. In particular, $n^{\beta_{\bar{d}}(n)} = h_{\bar{d}}(n)$ for any integer n.

Fact 4.4. Given a non-decreasing function g(x) such that:

$$dg(x) \le g\left(\frac{d^2}{d-1}x\right) \text{ and } g\left(\frac{D^2}{D-1}x\right) \le Dg(x),$$

there exists a sequence \bar{d} in $\{d, D\}^{\mathbb{N}}$ and a constant C such that:

$$\frac{1}{C}g(x) \le h_{\bar{d}}(x) \le Cg(x).$$

Proof. Set $x_k = \frac{1}{p_0} \dots \frac{1}{p_k}$ so that $h(x_{k+1}) = d_0 \dots d_{k+1} = h(x_k) d_{k+1}$ and $x_{k+1} = \frac{1}{p_{k+1}} x_k$. Assume by induction that d_0, \dots, d_k are constructed. Then:

(1) if $\frac{h(x_k)}{q(x_k)} \geq 1$, set $d_{k+1} = d$, and obtain:

$$\frac{dh(x_k)}{Dg(x_k)} \le \frac{dh(x_k)}{g(\frac{1}{p}x_k)} \le \frac{h(x_{k+1})}{g(x_{k+1})} = \frac{h(\frac{1}{p}x_k)}{g(\frac{1}{p}x_k)} \le \frac{dh(x_k)}{dg(x_k)},$$

(2) if $\frac{h(x_k)}{g(x_k)} < 1$, set $d_{k+1} = D$ and obtain:

$$\frac{Dh(x_k)}{Dg(x_k)} \le \frac{Dh(x_k)}{g(\frac{1}{P}x_k)} = \frac{h(x_{k+1})}{g(x_{k+1})} \le \frac{Dh(x_k)}{g(\frac{1}{p}x_k)} \le \frac{Dh(x_k)}{dg(x_k)}.$$

This shows that $\frac{d}{D} \leq \frac{h(x_k)}{g(x_k)} \leq \frac{D}{d}$. As moreover $\frac{P}{p} \leq \frac{x_k}{x_{k+1}} \leq \frac{p}{P}$ and g is non-decreasing, this proves the fact.

4.3. **Expected activity.** The expectation of activity is ruled by the exponent sequence.

Lemma 4.5. For any $\theta > 0$ and n large enough, there exists $C_{\theta} > 0$ such that:

$$\frac{1}{C_{\theta}} n^{\beta^{-\theta}(n)} \le \mathbb{E}a(Y_n) \le C_{\theta} n^{\beta^{\theta}(n)},$$

where the functions $\beta(n)$, $\beta^{\theta}(n)$ are defined in section 4.2. In particular, for any $\varepsilon > 0$ and n large enough:

$$\left| \frac{\log \mathbb{E}a(Y_n)}{\log n} - \beta(n) \right| \le \varepsilon.$$

For the following proof observe the fact that for any words $a(ww') \ge a(w)$ by proposition 3.8(4), so that the function $\mathbb{E}a(Y_n)$ is non-decreasing with n.

Proof. By proposition 3.8, the activity of Y_n relates to activity on the inherited process by $a_0(Y_n) = a_1(Y_n^1) + \cdots + a_1(Y_n^{d_0})$, where $a_k(w)$ is the activity function on the group $\Gamma(S_{d_k}, H_k F)$. Thus:

$$\mathbb{E}a_0(Y_n) = \sum_{t=1}^{d_0} \mathbb{E}a_1(Y_n^t).$$

Now by lemma 4.1, the conditioned variable $Y_n^t|m_t$ is a random alternate word Y'_{m_t} of random length m_t in the group $\Gamma(S_{d_1}, H_1F)$. Compute by conditioning and large deviation principle:

$$\mathbb{E}a_{1}(Y_{n}^{t}) = \sum_{i=0}^{n} \mathbb{E}(a_{1}(Y_{n}^{t})|m_{t} = i)P(m_{t} = i)$$

$$\leq \sum_{i \leq (p_{0} + \theta)n} P(m_{t} = i)\mathbb{E}a_{1}(Y_{i}) + nP(m_{t} \geq (p_{0} + \theta)n)$$

$$\leq \mathbb{E}a_{1}(Y'_{(p_{0} + \theta)n}) + nc_{\theta}^{n},$$

where $c_{\theta} < 1$ (use that $\mathbb{E}a_1(Y'_n)$ is non-decreasing to bound the sum). This shows that for N_0 large enough so that $d_0N_0c_{\theta}^{N_0} \leq 1$ and $n \geq N_0$:

$$\mathbb{E}a_0(Y_n) \le d_0 \mathbb{E}a_1(Y'_{(p_0+\theta)n}) + 1.$$

Note that c_{θ} hence N_0 depend on θ and on d_0 , but as the valency is bounded they can be chosen uniform for all d_i . This permits to iterate the above inequality to get for all k:

$$\mathbb{E}a_0(Y_n) \le d_0 \dots d_k \mathbb{E}a_{k+1}(Y_{(p_0+\theta)\dots(p_k+\theta)n}^{(k+1)}) + d_0 \dots d_{k-1} + \dots + d_0 + 1,$$

provided n is large enough. Recall that $Y_m^{(k)}$ is a random alternate word of length m in $\Gamma(S_{d_k}, H_k F)$. For $k(n) = \min\{k | (p_0 + \theta) \dots (p_k + \theta)n \leq N_0\}$, this shows:

$$\mathbb{E}a_0(Y_n) \le d_0 \dots d_{k(n)}(\mathbb{E}a_{k^{\theta}(n)+1}(Y_{N_0}^{(k^{\theta}(n)+1)}) + 1) \le n^{\beta^{\theta}(n)}(N_0 + 1),$$

by the trivial estimation $a_k(Y_{N_0}) \leq N_0$ for any k and the definition of $\beta^{\theta}(n)$.

Similarly for the lower bound:

$$\mathbb{E}a_0(Y_n) \ge d_0\mathbb{E}a_1(Y'_{(p_0-\theta)n}) - 1 \ge d_0 \dots d_k\mathbb{E}a_{k+1}(Y^{(k+1)}_{(p_0-\theta)\dots(p_k-\theta)n}) - (d_0 \dots d_{k-1} + \dots + 1),$$

so that for $k = k^{-\theta}(n) = \min\{k | (p_0 - \theta) \dots (p_k - \theta)n \le N_0\}$, one has:

$$\mathbb{E}a_0(Y_n) \ge \frac{1}{2}d_0 \dots d_{k^{-\theta}(n)} = \frac{1}{2}n^{\beta^{-\theta}(n)},$$

by the trivial estimate $\mathbb{E}a_k(Y_N^{(k)}) \geq \frac{3}{2}$ for any k and $N \geq 3$ (indeed, $a_k(Y_N^{(k)}) \geq 2$ as soon as there are two distinct elements among $\{\sigma_1^{-1}(1), \sigma_2^{-1}(1), \sigma_3^{-1}(1)\}$, which happens with probability $\geq \frac{3}{4}$ for any value of d_k).

5. Entropy exponents

5.1. **Main theorem.** Given a valency sequence $\bar{d} = (d_i)_i$ and $p_i = \frac{d_i - 1}{d_i^2}$, recall that the exponent sequence $\beta_{\bar{d}}(n)$ is defined by:

$$\beta_{\bar{d}}(n) = \beta(n) = \frac{\log(d_0 \dots d_{k(n)})}{\log n} \text{ where } k(n) = \min\{k | p_0 \dots p_k n \le 1\}.$$

Theorem 5.1. Let $\Gamma = \Gamma(S_{d_0}, HF)$ be a finitely generated saturated directed subgroup of $Aut(ET_{\bar{d}})$, μ the measure equidistributed on $S_{d_0}HFS_{d_0}$, and $\beta(n)$ the exponent sequence of \bar{d} , then for any $\varepsilon > 0$ and n large enough:

$$\left| \frac{\log H_{\Gamma,\mu}(n)}{\log n} - \beta(n) \right| \le \varepsilon.$$

In particular, $\overline{h}(\Gamma, \mu) = \limsup \beta(n)$ and $\underline{h}(\Gamma, \mu) = \liminf \beta(n)$.

Fact 4.3 ensures that if $d \leq d_i \leq D$ for all i, then:

$$\frac{1}{2} \le \beta_d \le \underline{h}(\Gamma, \mu) \le \overline{h}(\Gamma, \mu) \le \beta_D < 1.$$

Proof. First prove the upper bound, which is a straightforward generalisation of proposition 4.11 in [BKN]. Note the similarity with the upper bound in lemma 4.5. Indeed, under rewriting process $Y_n = (Y_n^1, \ldots, Y_n^{d_0})\tau_n$ one has:

$$H(Y_n) \le H(Y_n^1) + \dots + H(Y_n^{d_0}) + H(\tau_n),$$

where τ_n is equidistributed on S_{d_0} so $H(\tau_n) \leq C$ for a constant C depending only on D.

By lemma 4.1, the law of $Y_n^t|m_t$ under the length condition m_t is the law of a random alternate product Y'_{m_t} in the group $\Gamma(S_{d_1}, H_1F)$, and the length m_t has the binomial law of $\sum_{i=1}^n u_i$ for independent u_i in $\{0,1\}$ with $P(u_i=1)=p_0$, of entropy bounded by $C \log n$ for C depending only on D. This ensures (lemma A.4 in [BKN]):

$$H(Y_n^t) \le \sum_{m=0}^n H(Y_m') P(m_t = m) + H(m_t)$$

 $\le H(Y_{(p_0+\theta)n}') + nP(m_t \ge (p_0+\theta)n) + C\log n,$

by splitting the sum at $m=(p_0+\theta)n$ for an arbitrary $\theta>0$. By large deviation principle, there exists $c_{\theta}<1$ such that $P(m_t\geq (p_0+\theta)n)\leq c_{\theta}^n$. Thus there is N_0 depending only on θ and D such that for $n\geq N_0$ one has (for a slightly larger constant C since $nc_{\theta}^n=o(\log n)$):

$$H(Y_n) \le d_0 H(Y'_{(p_0+\theta)n}) + C \log n.$$

As for expected activity, this permits to integrate $H(n) = \sup_{k\geq 0} \{H(Y_n^{(k)})\} \leq Cn$, where $Y_n^{(k)}$ is a random alternate product in the group $\Gamma(S_{d_k}, H_k F)$ and C is a uniform constant depending only on the sizes of the generating sets $S_{d_k} \sqcup H_k F$, hence only on D and #HF, into:

$$H(n) \le d_0 \dots d_k H((p_0 + \theta) \dots (p_k + \theta)n) + (d_0 \dots d_{k-1} + \dots + 1)C \log n,$$

as long as $k \leq k^{\theta}(n)$, i.e. when $(p_0 + \theta) \dots (p_{k-1} + \theta)n \geq N_0$ (see section 4.2 for definition of $k^{\theta}(n)$ and $\beta^{\theta}(n)$). Thus:

$$H(Y_n) \le H(n) \le d_0 \dots d_{k^{\theta}(n)}(CN_0 + C\log n) = n^{\beta^{\theta}(n)}(CN_0 + C\log n),$$

and $H(Y_n) \leq n^{\beta(n)+2\varepsilon(\theta)}$ for n large enough, by proposition 4.2.

To prove the lower bound, the following fact is useful:

Fact 5.2. For $\gamma = \varphi g$ in $\Gamma(S_{d_0}, HF) = F \wr_{\partial T} G(S_{d_0}, H)$, one has:

$$P(Y_n = \gamma) \le \left(\frac{1}{\#F}\right)^{\#supp(\varphi)}$$
.

Proof of fact 5.2. Denote $Y_n =_{\Gamma} \varphi_n g_n$, and remark that $supp(\varphi_n) \subset A_{\partial}(Y_n)$ of size $a(Y_n)$. Also recall remark 3.6 that $A_{\partial}(Y_n)$ depends only on $g_n = s_1 h_1 \dots h_n s_{n+1}$, and fact 3.7 that given $A_{\partial}(g_n)$, any function $\varphi : \partial T \to F$ with support included in $A_{\partial}(g_n)$ appears with probability $\left(\frac{1}{\#F}\right)^{\#A_{\partial}(g_n)}$. This permits to compute by conditioning on activity:

$$P(Y_n = \gamma) = \sum_{a=1}^n P(Y_n = \gamma | a(Y_n) = a) P(a(Y_n) = a)$$

$$\leq \sum_{a \geq \#supp(\varphi)} P(\varphi_n = \varphi | a(Y_n) = a) P(a(Y_n) = a)$$

$$\leq \sum_{a > \#supp(\varphi)} \left(\frac{1}{\#F}\right)^a P(a(Y_n) = a) \leq \left(\frac{1}{\#F}\right)^{\#supp(\varphi)}$$

This fact guarantees:

$$H(Y_n) = -\sum_{\gamma \in \Gamma} P(Y_n = \gamma) \log P(Y_n = \gamma)$$

$$\geq C \sum_{\gamma \in \Gamma} P(Y_n = \gamma) \# supp(\varphi)$$

$$= C \mathbb{E}_{\mu^{*n}} \# supp(\varphi) = C \mathbb{E} \# supp(\varphi_n),$$

and the expected value of the size of the support of φ_n relates to activity.

More precisely by fact 3.7, given $A_{\partial}(g_n)$, the function $\varphi_n:A_{\partial}(g_n)\to F$ is random, so that:

$$\mathbb{E}[\#supp(\varphi_n)|A_{\partial}(g_n)] = \frac{\#F - 1}{\#F} \#A_{\partial}(g_n).$$

This permits once again to compute by conditioning on activity:

$$\mathbb{E} \# supp(\varphi_n) = \sum_{a=1}^n \mathbb{E} [\# supp(\varphi_n) | \# A(Y_n) = a] P [\# A(Y_n) = a]$$
$$= \sum_{a=1}^n \frac{\# F - 1}{\# F} a P [\# A(Y_n) = a] = \frac{\# F - 1}{\# F} \mathbb{E} a(Y_n).$$

By lemma 4.5, conclude:

$$H(Y_n) \ge C \frac{\#F-1}{\#F} \mathbb{E}a(Y_n) \ge C \frac{\#F-1}{\#F} n^{\beta(n)-\varepsilon}.$$

Note that the proof for the upper bound remains valid for the group $G(S, H) < Aut(T_{\bar{d}})$, i.e. when the group F is trivial, but the lower bound is true only with a non-trivial finite group F (otherwise fact 5.2 is obviously empty).

Remark 5.3. In information theory, the entropy is the "average number of digits" needed to describe some data. In this heuristic point of view, theorem 5.1 is a corollary of lemma 4.5.

Indeed, the activity a(w) of a word w is equivalent to the size of the minimal tree T(w) defined in section 3.2 (recall $a(w) \leq \#\partial T(w) \leq Da(w)$). Moreover, the element $\gamma =_{\Gamma} w$ in Γ is described by figure 1, with σ_u for u non-active vertices, w_v for v active vertices and $\varphi(x)$ at active boundary points x. As each of them is described by $\leq C$ digits for some C depending uniquely on the bound D on valency, the element γ is described with $\leq C \# T(w) \approx a(w)$ digits. Moreover, since any function $\varphi : A_{\partial}(w) \to F$ appears equally likely, one needs at least a(w) digits to describe γ .

Figure 1 is in this sense the "best description" of the element γ in Γ (position in the Cayley graph) represented by the word w (path in the Cayley graph). The loss of information from w to γ is somewhat described by the graph structure of the ascendance forest AF(w) of section 3.3.

5.2. Precise entropy exponent and oscillation phenomena. Theorem 5.1 exhibits a large variety of behaviors for entropy of random walk. In particular, it implies theorem 1.2 for $\frac{1}{2} \le \alpha \le \beta < 1$.

Corollary 5.4. For any $\frac{1}{2} \leq \beta < 1$, there is a valency sequence \bar{d} such that the entropy exponent of the random walk Y_n on a finitely generated saturated directed group $\Gamma(S_{d_0}, HF) < Aut(ET_{\bar{d}})$ is:

$$h(\Gamma, \mu) = \beta.$$

Proof. Take $d \leq D$ such that $\beta_2 \leq \beta_d = \frac{\log d}{\log p} \leq \beta \leq \frac{\log D}{\log P} = \beta_D < 1$. There exist $\lambda \in [0,1]$ such that $\beta = \frac{\log(d^{\lambda}D^{1-\lambda})}{\log(p^{\lambda}P^{1-\lambda})}$. Define the sequence \bar{d} by $d_i \in \{d,D\}$ for all i and $\#\{i \leq n | d_i = d\} = [\lambda n]$. Then the exponent sequence $\beta_{\bar{d}}(n) \to \beta$ and the corollary follows from theorem 5.1.

Remark 5.5. This corollary shows in particular that any saturated directed group of a binary tree has entropy exponent $h(\Gamma,\mu)=\frac{1}{2}$. This is the case of the groups $\Gamma_{\omega}=F\wr_{\partial T_2}G_{\omega}$ for G_{ω} an Aleshin-Grigorchuk group, for which a great variety of growth behaviors are known. For instance, $\Gamma_{(012)^{\infty}}$ has growth function $b_{(012)^{\infty}}(r)\approx e^{r^{\alpha_0}}$ for an explicit $\alpha_0<1$ ([BE]), and for any $\alpha\in[\alpha_0,1]$ there is a group $G_{\omega(\alpha)}$ with growth such that $\lim\frac{\log\log b_{\omega(\alpha)}(r)}{\log r}=\alpha$ ([Bri3]). Considering the entropy, all these growth behaviors collapse to a unique entropy exponent.

Corollary 5.6. For any $\frac{1}{2} \leq \alpha < \beta < 1$, there is a valency sequence \bar{d} such that the lower and upper entropy exponents of the random walk Y_n on a finitely generated saturated directed group $\Gamma(S_{d_0}, HF) < Aut(ET_{\bar{d}})$ are:

$$\underline{h}(\Gamma,\mu) = \alpha \ \ and \ \overline{h}(\Gamma,\mu) = \beta.$$

Proof. By theorem 5.1, it is sufficient to construct an appropriate exponent sequence in order to prove the corollary. Take $d \leq D$ such that $\beta_d \leq \alpha < \beta \leq \beta_D$. Construct a sequence \bar{d} such that $d_i \in \{d, D\}$ for all i according to the following rules.

Recall the definition (see section 4.2) of $\beta(n)$ as satisfying $n^{\beta(n)} = d_0 \dots d_{k(n)}$ where k(n) as the unique integer such that $\frac{p_{k(n)}}{n} < p_0 \dots p_{k(n)} \leq \frac{1}{n}$. In order to ease the reading of the present proof, use the shortcut notation $p_0 \dots p_{k(n)} \approx \frac{1}{n}$. There exists a constant C depending uniquely on D such that the following statements are true if all relations $x \approx y$ below are replaced by $\frac{y}{C} \leq x \leq Cy$.

Suppose k(n) is such that $p_0 \dots p_{k(n)} \approx \frac{1}{n}$ and $d_0 \dots d_{k(n)} \approx n^{\alpha}$ for some n (hence $|\beta(n) - \alpha| \leq \frac{C}{\log n}$), then set $d_{k(n)+1} = \dots = d_{k(n)+l} = D$ for l such that for some minimal integer m:

$$\frac{P^l}{n} \approx p_0 \dots p_{k(n)} P^l = p_0 \dots p_{k(n)+l} \approx \frac{1}{m},$$

$$n^{\alpha} D^l \approx d_0 \dots d_{k(n)} D^l = d_0 \dots d_{k(n)+l} \approx m^{\beta}.$$

This forces $n^{\alpha}D^{l} \approx \left(\frac{n}{P^{l}}\right)^{\beta}$, hence $l = \frac{\beta - \alpha}{\log(P^{\beta}D)} \log n + o(\log n)$.

Suppose l(m) is such that $p_0 \dots p_{l(m)} \approx \frac{1}{m}$ and $d_0 \dots d_{l(m)} \approx m^{\beta}$ for some m (hence $|\beta(m) - \beta| \leq \frac{C}{\log m}$), then set $d_{l(m)+1} = \dots = d_{l(m)+k} = d$ for k such that for some minimal integer n:

$$\frac{p^k}{m} \approx p_0 \dots p_{l(m)} p^k = p_0 \dots p_{l(m)+k} \approx \frac{1}{n},$$

$$m^{\beta} d^k \approx d_0 \dots d_{l(m)} d^k = d_0 \dots d_{l(m)+k} \approx n^{\alpha}.$$

This forces $m^{\beta}d^k \approx \left(\frac{m}{p^k}\right)^{\alpha}$, hence $k = \frac{\alpha - \beta}{\log(p^{\alpha}d)}\log m + o(\log m)$.

This process produces a sequence \bar{d} such that $\alpha - \frac{C}{\log n} \leq \beta(n) \leq \beta + \frac{C}{\log n}$ for all n. Moreover, $d_0 \dots d_{k(n)} \approx n^{\alpha}$ for infinitely many n and $d_0 \dots d_{l(m)} \approx m^{\beta}$ for infinitely many m. Hence $\underline{h}(\Gamma, \mu) = \alpha$, and $\overline{h}(\Gamma, \mu) = \beta$.

Remark 5.7. The above two corollaries are obtained by taking particular instances of exponent functions $\beta(n)$. Theorem 5.1 provides a variety of behaviors for entropy functions. For instance similarly to theorems 7.2 on growth functions in [Gri1] and [Bri3], there exists uncountable antichains of entropy functions with $\underline{h}(\Gamma,\mu) = \alpha < \beta = \overline{h}(\Gamma,\mu)$ for any given $\frac{1}{2} \leq \alpha < \beta < 1$, as easily proved by playing with exponent functions.

5.3. **Frequency of oscillations.** Corollary 5.6 shows that at different scales, the entropy exponent of a random walk can take different values. In order to study the difference between such scales, given $\frac{1}{2} \le \alpha \le \beta \le 1$ and a function H(n), introduce the following quantities, called respectively *upper* and *lower pseudo period exponents*

of the function H(n) for α and β :

$$u_H(\alpha,\beta) = \inf\{\nu | \exists N_0, \forall n \geq N_0, \text{if } H(n) \leq n^{\alpha}, \text{ then } \exists n \leq m \leq n^{\nu}, H(m) \geq m^{\beta}\}, l_H(\alpha,\beta) = \inf\{\lambda | \exists N_0, \forall m \geq N_0, \text{if } H(m) \geq m^{\beta}, \text{ then } \exists m \leq n \leq m^{\lambda}, H(n) \leq n^{\alpha}\}.$$

For $H = H_{\Gamma,\mu}$ entropy of a finitely supported measure μ on a group Γ , write $u_H(\alpha,\beta) = u_{\Gamma,\mu}(\alpha,\beta)$ and $l_H(\alpha,\beta) = l_{\Gamma,\mu}(\alpha,\beta)$.

Note that by playing with the sequence \bar{d} in the proof of corollary 5.6, one can easily produce examples of random walks with arbitrarily large pseudo period exponents, i.e. low frequency. To study how high the frequency may be, i.e. how small the pseudo periods, introduce the functions:

 $u(\alpha, \beta) = \inf\{u_{\Gamma,\mu}(\alpha, \beta) | \mu \text{ finitely supported symmetric measure on a group } \Gamma\},\ l(\alpha, \beta) = \inf\{l_{\Gamma,\mu}(\alpha, \beta) | \mu \text{ finitely supported symmetric measure on a group } \Gamma\}.$

The finitely generated saturated directed groups of theorem 5.1 will provide the upper bounds in the following.

Theorem 5.8. For $\frac{1}{2} < \alpha \le \beta < 1$,

$$u(\alpha, \beta) = \frac{\alpha - 1}{\beta - 1}$$
, and $\frac{\beta}{\alpha} \le l(\alpha, \beta) \le \frac{\beta - \frac{1}{2}}{\alpha - \frac{1}{2}}$.

Proof. The lower bounds follow from elementary properties of entropy. For a submultiplicative function $H(n) \leq n^{\alpha}$ implies $H(kn) \leq kn^{\alpha}$, so that if $H(kn) \geq (kn)^{\beta}$, then $(kn)^{\beta} \leq kn^{\alpha}$, so $kn \geq n^{\frac{1-\alpha}{1-\beta}}$ and $u_H(\alpha,\beta) \geq \frac{1-\alpha}{1-\beta}$. For an increasing function $H(m) \geq m^{\beta}$ and $H(n) \leq n^{\alpha}$ imply $n \geq m^{\frac{\beta}{\alpha}}$, so $l_H(\alpha,\beta) \geq \frac{\beta}{\alpha}$.

Concerning the upper pseudo period exponents, the proof of corollary 5.6 shows that given n, one can take $m \leq C \frac{n}{P^l}$ for $l \leq (\frac{\beta - \alpha}{\log(P^\beta D)} + \varepsilon) \log n$, where C depends only on D and $\varepsilon > 0$ is arbitrary, i.e. $m \leq n^{\nu}$ for:

$$\nu = 1 - \frac{(\beta - \alpha)\log P}{\log(P^{\beta}D)} - \varepsilon\log P = 1 - \frac{\beta - \alpha}{\beta + \frac{\log D}{\log(\frac{D-1}{P^2})}} + \varepsilon|\log P|.$$

As D tends to infinity:

$$\frac{\log D}{\log(\frac{D-1}{D^2})} = \frac{\log D}{\log(D-1) - 2\log D} = \frac{1}{\frac{\log(D-1)}{\log D} - 2} \longrightarrow -1,$$

which proves:

$$u(\alpha, \beta) \le 1 - \frac{\beta - \alpha}{\beta - 1} = \frac{\alpha - 1}{\beta - 1}.$$

Similarly for the lower pseudo period exponent, the proof of corollary 5.6 shows that given m, one can take $n \leq C \frac{m}{p^k}$ for $k = (\frac{\alpha - \beta}{\log(p^{\alpha}d)} + \varepsilon) \log m$, i.e. $n \leq m^{\lambda}$ for:

$$\lambda = 1 - \frac{(\alpha - \beta) \log p}{\log(p^{\alpha}d)} - \varepsilon \log p = 1 - \frac{\alpha - \beta}{\alpha + \frac{\log d}{\log(\frac{d-1}{d^2})}} + \varepsilon |\log p|.$$

Taking d=2, this proves:

$$l(\alpha, \beta) \le 1 + \frac{\beta - \alpha}{\alpha - \frac{1}{2}} = \frac{\beta - \frac{1}{2}}{\alpha - \frac{1}{2}}.$$

Remark 5.9. The value $u(\alpha, \beta) = \frac{\alpha-1}{\beta-1}$ is tightly related to subadditivity of entropy, which implies in particular that $\overline{h}(\Gamma, \mu) \leq 1$ for any group Γ with finitely supported measure μ . Also theorem 5.1 shows that the upper bound $l(\alpha, \beta) \leq \frac{\beta-\frac{1}{2}}{\alpha-\frac{1}{2}}$ is optimal among saturated directed groups with the measure μ equidistributed on $S_{d_0}HFS_{d_0}$. It is unclear whether this bound is optimal in general. It could be related to question 9.3 on lower bound $\underline{h}(\Gamma, \mu) \geq \frac{1}{2}$.

5.4. **Drift of the random walk.** Theorem 5.1 provides estimates on the drift $L_{\Gamma,\mu}(n) = \mathbb{E}||Y_n||$ of the random walk Y_n of step distribution μ equidistributed on $S_{d_0}HFS_{d_0}$, where ||.|| is the word norm for some (arbitrary) generating set.

Corollary 5.10. For any $\varepsilon > 0$ and n large enough:

$$\beta(n) - \varepsilon \le \frac{\log L_{\Gamma,\mu}(n)}{\log n} \le \frac{1 + \beta(n)}{2} + \varepsilon.$$

Proof. Lemma 6 and 7 in [Ers1] show that there are $c_1, c_2, c_3 > 0$ with:

$$c_1 H_{\Gamma,\mu}(n) - c_2 \le L_{\Gamma,\mu}(n) \le c_3 \sqrt{n(H_{\Gamma,\mu}(n) + \log n)}.$$

Combine with theorem 5.1.

6. Return Probability

6.1. General estimates. Given a valency sequence $\bar{d} = (d_i)_i$, set:

$$l(n) = l_{\bar{d}}(n) = \max\{l | \frac{d_0^3}{d_0 - 1} \dots \frac{d_l^3}{d_l - 1} = \frac{d_0}{p_0} \dots \frac{d_l}{p_l} \le n\},$$

and define the auxiliary exponent sequence:

$$\beta'(n) = \beta'_{\bar{d}}(n) = \frac{\log(d_0 \dots d_{l(n)})}{\log n}.$$

In particular, $p_0 \dots p_{l(n)} n \approx d_0 \dots d_{l(n)} = n^{\beta'(n)}$.

Theorem 6.1. For any $\varepsilon > 0$ the return probability of the simple random walk Y_n with generating set $S_{d_0}HFS_{d_0}$ on the saturated directed group $\Gamma(S_{d_0}, HF) < Aut(ET_{\bar{d}})$ satisfies for n large enough:

$$\beta'(n) - \varepsilon \le \frac{\log(-\log P(Y_n = id))}{\log n} \le \beta(n) + \varepsilon.$$

In particular, $\underline{p}(\Gamma) \ge \liminf \beta'(n)$ and $\overline{p}(\Gamma) \le \limsup \beta(n)$.

For instance in the case of constant valency d, one has:

$$\beta'_d = \frac{1}{3 - \frac{\log(d-1)}{\log d}} \le \frac{\log(-\log P(Y_n = id))}{\log n} \le \frac{1}{2 - \frac{\log(d-1)}{\log d}} = \beta_d.$$

Note that $\beta_2' = \frac{1}{3}$ and $\beta_d' \longrightarrow_{d \to \infty} \frac{1}{2}$ for lower bounds, compared with $\beta_2 = \frac{1}{2}$ and $\beta_d \longrightarrow_{d \to \infty} 1$ for upper bounds.

Proof. Proposition 4.5 implies that for n large enough:

$$P(a(Y_n) \ge n^{\beta(n)+\varepsilon}) \le \frac{\mathbb{E}a(Y_n)}{n^{\beta(n)+\varepsilon}} \le \frac{n^{\beta(n)+\frac{\varepsilon}{2}}}{n^{\beta(n)+\varepsilon}} = n^{-\frac{\varepsilon}{2}} \longrightarrow 0.$$

Using the fact that for fixed n, the function $P(Y_n = \gamma)$ is maximal at $\gamma = id$ by symmetry of the random walk, and the inequality of proposition 3.8 deduce:

$$1 \longleftarrow P(a(Y_n) \le n^{\beta(n)+\varepsilon}) = \sum_{\{\gamma \mid \exists w = \Gamma^{\gamma}, a(w) \le n^{\beta(n)+\varepsilon}\}} P(Y_n = \gamma) \le P(Y_n = id) C^{n^{\beta(n)+\varepsilon}}.$$

This proves the upper bound.

Recall that given a word Y_n of length n, the rewriting process provides for each vertex $v \in T_{\bar{d}}$ a word Y_n^v of random length m_v (proposition 3.2). Given $\theta > 0$ small enough so that for any n large enough $l(n) \leq k^{-\theta}(n)$ (defined in section 4.2 by $(p_0 - \theta) \dots (p_{k^{-\theta}(n)} - \theta)n \approx N_0$ for an arbitrary $N_0 \geq 1$, so that the inequality holds for large n as soon as $\frac{d_i-1}{d_i^2} - \theta > \frac{d_i-1}{d_i^3}$ for all i), observe the:

Fact 6.2. For a word Y_n with $a(Y_n) < n^{\beta'(n)}$, there exists a vertex v in $T_{\bar{d}}$ such that:

- (1) |v| < l(n),
- (2) there is t such that $m_{vt} < (p_{|v|} \theta)m_v$,
- (3) $m_v \ge (p_0 \theta) \dots (p_{|v|-1} \theta)n$.

Proof of fact 6.2. By contradiction assume that for all $|v| \leq l(n)$ and t one has $m_{vt} \geq (p_{|v|} - \theta)m_v$. Then by induction on |v|, for all v in level l(n):

$$m_v \ge (p_0 - \theta) \dots (p_{l(n)-1} - \theta)n \ge \frac{N_0}{C(p_{l(n)} - \theta) \dots (p_{k-\theta(n)} - \theta)} \ge 1,$$

so that $a(Y_n^v) \geq 1$. Then $a(Y_n) \geq \sum_{|v|=l(n)} a(Y_n^v) \geq d_0 \dots d_{l(n)} = n^{\beta'(n)}$, which is a contradiction. This shows the existence of a vertex v satisfying (1) and (2). Such a vertex that is closest to the root also satisfies (3).

This fact 6.2 guarantees that:

$$P(a(Y_n) \le n^{\beta'(n)}) \le \sum_{|v| \le l(n)} P[\exists t, m_{vt} \le (p_{|v|} - \theta)m_v \text{ and } m_v \ge (p_0 - \theta) \dots (p_{|v|-1} - \theta)n].$$

Now $P[m_{vt} \leq (p_{|v|} - \theta)m_v \text{ and } m_v] = P[m_{vt} \leq (p_{|v|} - \theta)m_v|m_v]P(m_v)$, and by proposition 4.1 there is $c_{\theta} < 1$ with $P[m_{vt} \leq (p_{|v|} - \theta)m_v|m_v] \leq c_{\theta}^{m_v}$. Then:

(6)
$$P(a(Y_n) \le n^{\beta'(n)}) \le \sum_{|v| \le l(n)} c_{\theta}^{(p_0 - \theta) \dots (p_{l(n) - 1} - \theta)n} = n^{\beta'(n)} c_{\theta}^{(p_0 - \theta) \dots (p_{l(n) - 1} - \theta)n},$$

because there are $d_0 \dots d_{l(n)} = n^{\beta'(n)}$ vertices such that $|v| \leq l(n)$. Compute by conditioning on activity, recalling that fact 3.7 ensures $P[\varphi_n = id|a(Y_n) = a] = (\frac{1}{\#F})^a$:

$$P(Y_n = id) \le P(\varphi_n = id) = \sum_{a=0}^n P[\varphi_n = id | a(Y_n) = a] P(a(Y_n) = a),$$

= $\sum_{a=0}^n \left(\frac{1}{\#F}\right)^a P(a(Y_n) = a).$

The decay of $\left(\frac{1}{\#F}\right)^a$ with a permits to split the sum between $a < n^{\beta'(n)}$ and $a \ge n^{\beta'(n)}$. Obtain:

$$P(Y_n = id) \leq P(a(Y_n) < n^{\beta'(n)}) + \left(\frac{1}{\#F}\right)^{n^{\beta'(n)}} P(a(Y_n) \geq n^{\beta'(n)}),$$

$$\leq n^{\beta'(n)} c_{\theta}^{(p_0 - \theta) \dots (p_{l(n) - 1} - \theta)n} + \left(\frac{1}{\#F}\right)^{n^{\beta'(n)}},$$

by inequality (6). Now there is a function $\varepsilon(\theta) \longrightarrow_{\theta} 0$ such that $(p_0 - \theta) \dots (p_{l(n)-1} - \theta)n \ge n^{\beta'(n)-\varepsilon(\theta)}$ (cf proposition 4.2), so:

$$P(Y_n = id) \le \exp(-cn^{\beta'(n) - \varepsilon(\theta)}),$$

which proves the lower bound.

As for theorem 5.1, the upper bound is valid for G(S, H), i.e. for $F = \{1\}$, but the lower bound is valid only with a non-trivial finite group F.

Remark 6.3. Fact 6.2 shows that if the activity is small, then there is at least one edge along which the word length is contracted more than expected, with $m_{vt} \leq (p_{|v|} - \theta)m_v$ instead of $m_{vt} = p_{|v|}m_v$. In fact, in order to have $a(Y_n) < n^{\beta'(n)}$, such a strong contraction must occur at many edges, so that inequality (6) does not seem optimal. The example below shows that the lower bound of theorem 6.1 is tight, and thus inequality (6) is essentially optimal in general. It might however be improved for particular instances of saturated directed groups.

6.2. A specific example. Consider the particular case of a binary tree and specific generators $s=(1,1)\sigma$ with σ the non-trivial permutation in S_2 and h=(h,s). The group $\langle s,h\rangle < Aut(T_2)$ is a well-known automata group isomorphic to the infinite dihedral group $D_{\infty}=\langle s,h|s^2=h^2=1\rangle$.

Proposition 6.4. A random walk Z_n on the extended directed group $\Gamma(S, HF) = F \wr_{\partial T_2} D_{\infty} < Aut(ET_2)$ with F abelian finite with step distribution equidistributed on SHFHS satisfies:

$$\lim \frac{\log |\log P(Z_n = id)|}{\log n} = \frac{1}{3} \text{ and } \lim \frac{\log \mathbb{E}||Z_n||}{\log n} = \frac{1}{2}.$$

In this particular case, the lower bounds of theorem 6.1 and corollary 5.10 are tight. Note that this specific group played a crucial role in the construction of

antichains of growth functions in [Gri1] and in the construction of groups with oscillating growth in [Bri1] (chapter 2).

Proof. Theorem 6.1 and corollary 5.10 ensure:

$$\liminf \frac{\log |\log P(Z_n = id)|}{\log n} \ge \frac{1}{3} \text{ and } \lim \frac{\log \mathbb{E}||Z_n||}{\log n} \ge \frac{1}{2}.$$

To get an upper bound, compare with the usual wreath product $F_{\partial_{\infty}}D_{\infty}$, for which the return probability satisfies $P(\tilde{X}_n=id)\approx e^{-n^{\frac{1}{3}}}$ (theorem 3.5 in [PSC2], noting that \mathbb{Z} and D_{∞} with their usual generating sets have the same Cayley graph) and the drift $L_{F_{\partial_{\infty}}D_{\infty}}(n)\approx n^{\frac{1}{2}}$ (by lemma 3 in [Ers2]).

Precisely, consider the usual wreath product (lamplighter):

$$F \wr_{D_{\infty}} D_{\infty} = \{\Phi : D_{\infty} \to F | supp(\Phi) \text{ is finite} \} \rtimes D_{\infty},$$

with the action $d.\Phi(x) = \Phi(xd)$. Denote $S = \langle s \rangle \simeq S_2$ and $H = \langle h \rangle \simeq S_2$, so that $D_{\infty} \simeq G(S,H) < Aut(T_2)$, and denote F the subgroup $\{\Phi: D_{\infty} \to F | \forall x \neq 1_{D_{\infty}}, \Phi(x) = 1_F\}$ in $F \wr_{D_{\infty}} D_{\infty}$. Let \tilde{X}_n be the random walk with alternate successive increments equidistributed in the finite symmetric sets HSH and F (this is, up to negligible first and last factors, the random walk with step distribution μ equidistributed on the finite symmetric set SHFHS):

$$\tilde{X}_n = h_1 s_1 h'_1 f_1 h_2 s_2 h'_2 f_2 \dots h_n s_n h'_n f_n.$$

It induces in particular a random walk X_n on the base group D_{∞} given by:

$$X_n = h_1 s_1 h'_1 h_2 s_2 h'_2 \dots h_n s_n h'_n = r_1 r_2 \dots r_n,$$

with $r_i = h_i s_i h_i'$. The value of \tilde{X}_n in $F \wr_{D_{\infty}} D_{\infty}$ is given by the value of X_n in D_{∞} together with a function $\Phi_n : D_{\infty} \to F$ the support of which is included in $\{r_1^{-1}, (r_1 r_2)^{-1}, \ldots, (r_1 r_2 \ldots r_n)^{-1}\}$, since at time i the lamp in position $(r_1 r_2 \ldots r_i)^{-1}$ is modified.

With obvious identification of S, H, F, denote Z_n the similar random walk on $F \wr_{\partial T_2} G(S, H)$:

$$Z_n = h_1 s_1 h'_1 f_1 h_2 s_2 h'_2 f_2 \dots h_n s_n h'_n f_n.$$

The value of Z_n is given by the value of X_n in $G(S, H) \simeq D_{\infty}$ and a function $\varphi_n : \partial T_2 \to F$ with support included in $\{1^{\infty}r_1^{-1}, 1^{\infty}(r_1r_2)^{-1}, \dots, 1^{\infty}(r_1r_2\dots r_n)^{-1}\}$ by proposition 3.8.

Now the Schreier graph $1^{\infty}G(S,H)$ can easily be described as a semi-line. If w is a reduced representative word in $D_{\infty}=\langle s,h|s^2=h^2=1\rangle$ then $1^{\infty}hw=1^{\infty}w$ and $1^{\infty}w\neq 1^{\infty}w'$ if $w\neq w'$ and both w and w' start with s, so there is a canonical 2-covering application $c:Cay(D_{\infty},\{s,h\}) \to 1^{\infty}G(S,H)$ with c(w)=c(hw). This implies:

$$\varphi_n(x) = \sum_{y \in c^{-1}(x)} \Phi_n(y)$$

because F is abelian, so that the order of increments does not influence the sum. This shows that if $\tilde{X}_n = 1$ in $F \wr_{D_{\infty}} D_{\infty}$ then $Z_n = 1$ in $F \wr_{\partial T_2} G(S, H)$, hence $P(Z_n = 1) \geq P(\tilde{X}_n = 1)$ and the group $F \wr_{\partial T_2} G(S, H)$ is a quotient of $F \wr_{D_{\infty}} D_{\infty}$ with identification of the generators, so $||\tilde{X}_n|| \leq ||Z_n||$.

7. Higher order oscillations

This section aims at proving theorem 1.3 and treating the case $\beta=1$ in theorem 1.2. The following construction is designed to obtain groups $\Delta(S', H'F')$ that ressemble $\Gamma(S, HF)$ at some scales and non-amenable groups at other scales. They are still directed groups of a rooted tree but of unbounded valency \bar{d} (in the cases of interest here) and not saturated.

The construction generalises [KP], where a group $\Delta(S', H'_{\omega})$ is constructed given an Aleshin-Grigorchuk group $\Gamma(S, H_{\omega})$ and additional data. Theorem 6.1 and corollary 5.10 permit to show that some of the groups $\Delta = \Delta(S', H'_{\omega})$ or almost equivalently some of the piecewise automatic groups of [Ers4] satisfy:

$$\underline{p}(\Delta) \leq \frac{1}{2}, \overline{p}(\Delta) = 1 \text{ and } \underline{\delta}(\Delta) \leq \frac{3}{4}, \overline{\delta}(\Delta) = 1.$$

The description of the construction in [KP] is more algebraic, whereas the point of view adopted here is in terms of automorphisms of an ambiant tree $T_{\bar{e}}$.

7.1. **Definition of** $\Delta(S', H'F')$. Given a bounded sequence \bar{d} and a saturated directed finitely generated group $\Gamma = \Gamma(S, HF) < Aut(ET_{\bar{d}})$, construct a modification Δ of this group, acting on an extended spherically homogeneous rooted tree $ET_{\bar{e}}$ for another sequence $\bar{e} = (e_l)_{l \in \mathbb{N}}$, with $e_l = d_l + d'_l$ for some $d'_l \geq 0$, possibly $d'_l = \infty$. Note that there is a canonical inclusion $ET_{\bar{d}} \subset ET_{\bar{e}}$, hence a canonical embedding:

$$Aut(ET_{\bar{d}}) \hookrightarrow Aut(ET_{\bar{e}}).$$

Corresponding to the group $\Gamma(S, HF)$, determined by the finite groups S, H, F and the portraits of their elements, i.e. their realisation in $Aut(ET_{\bar{d}})$, construct a new group $\Delta(S', H'F') < Aut(ET_{\bar{e}})$, where $S \simeq S'$, $H \simeq H'$ and $F \simeq F'$ as abstract groups. Define the generators via the wreath product isomorphism of proposition 2.2.

(1) The element s' in S' corresponding to $s \in S \simeq S'$ has the form:

$$s' = (1, \dots, 1)s'$$

where s' is a permutation of $\{1,\ldots,d_0,d_0+1,\ldots,d_0+d_0'\}$ respecting the decomposition $\{1,\ldots,d_0\}\sqcup\{d_0+1,\ldots,d_0+d_0'\}$, so that $s'|_{\{1,\ldots,d_0\}}=s\in S=S_{d_0}< S_{e_0}$ (canonical inclusion) and moreover $s''=s's^{-1}=s'|_{\{d_0+1,\ldots,d_0+d_0'\}}$ is chosen so that $s\mapsto s''$ is a morphism of groups from $S\simeq S'$ into the permutation group $S_{\{d_0+1,\ldots,d_0+d_0'\}}\simeq S_{d_0'}$. Denote S'' its image.

(2) The element h' in H' corresponding to $h = (h_1, \sigma_2, \ldots, \sigma_{d_0}) \in H \simeq H'$ has the form:

$$h' = (h'_1, \sigma'_2, \dots, \sigma'_{d_0}, 1, \dots, 1)h''$$

where $\sigma'_2, \ldots, \sigma'_{d_0} \in S'_1$ are defined as above, corresponding to $\sigma_2, \ldots, \sigma_{d_0} \in S_1$, $h'_1 \in H'_1$ corresponds to $h_1 \in H_1$, and h'' is a permutation in $S_{d_0+d'_0}$ with support included in $\{d_0+1,\ldots,d_0+d'_0\}$ so that $h\mapsto h''$ is a morphism of groups with image $H'' < S_{d'_0}$.

Elements of H' do not act at the boundary $\partial T_{\bar{e}}$.




FIGURE 2. The tree $T_{\bar{e}}$, for $d_0 = d_1 = 2$ and $e_0 = e_1 = 4$, with the subtree $T_{\bar{d}}$ in plain edges, dashed edges where $\langle S_l'', H_l'' F_l'' \rangle \langle S_{d_l'}$ act, and dotted edges where the action of Δ is trivial.

(3) The element f' in F' corresponding to $\varphi_f \in F \simeq F'$ has the form:

$$f' = (f'_1, 1, \dots, 1)f'',$$

where $f_1' \in F'$ corresponds to $\varphi_f \in \Gamma(S_1, H_1F)$ and f'' is a permutation in $S_{d_0+d_0'}$ with support included in $\{d_0+1,\ldots,d_0+d_0'\}$ so that $f \mapsto f''$ is a morphism of groups with image $F'' < S_{d_0'}$.

The element f' acts at the boundary $\partial T_{\bar{e}}$ by $\varphi(x) = 1_F$ if $x \neq 1^{\infty}$ and $\varphi(1^{\infty}) = f$.

(4) Any two elements f'' and h'' in $S_{\{d_0+1,\dots,d_0+d'_0\}}$ commute.

Note that similarly to the definition of $h \in Aut(T_{\bar{d}})$ in section 2.1, the definitions of h' and f' are recursive for they involve the generators h'_1 and f'_1 of the group $\Delta(S'_1, H'_1F'_1) < Aut(ET_{\sigma\bar{e}})$ associated to the saturated directed group $\Gamma(S_1, H_1F) < Aut(ET_{\bar{e}})$.

Condition (4) implies by recursion that at any level l the elements h_l'' and f_l'' in $S_{\{d_l+1,\ldots,d_l+d_l'\}}$ commute, which implies that the subgroup $\langle \{s_l''\}, \{h_l''\}, \{f_l''\} \rangle < S_{d_l'}$ is a quotient of the free product of finite groups $S_l'*(H_l'\times F_l') \simeq S_l*(H_l\times F)$, possibly an infinite quotient in the case $d_l' = \infty$. Denote $S_l'' = \{s_l''\}, H_l'' = \{h_l''\}, F_l'' = \{f_l''\}$. They are subgroups of $S_{\{d_l+1,\ldots,d_l+d_l'\}}$.

By recursion, the action of the generators s', h', f' is well defined on the whole tree $T_{\bar{e}}$. Moreover, only generators f' act non trivially at the boundary $\partial T_{\bar{e}}$, thus the action of s', h', f' on $ET_{\bar{e}}$ is well defined.

To summarise, the group $\Delta(S', H'F')$ is defined by $\Gamma(S, HF)$ and a collection of groups $\langle S''_l, H''_l F''_l \rangle < S_{d'_l}$ that are quotients of the free products of finite groups $S_l * H_l F$ by identification of generators.

7.2. Elementary properties of $\Delta(S', H'F')$. Note that $\Delta(S', H'F')$ is directed but not saturated. As a shortcut, write $\Gamma_l = \Gamma(S_l, H_lF) < Aut(ET_{\sigma^l\bar{d}})$ and $\Delta_l = \Delta(S'_l, H'_lF'_l) < Aut(ET_{\sigma^l\bar{d}})$.

Properties 7.1. (1) The canonical isomorphism of proposition 2.2 induces a canonical embedding:

$$\Delta(S', H', F') \hookrightarrow \Delta(S'_1, H'_1, F'_1) \wr \langle S', H'', F'' \rangle.$$

More generally, $\Delta_l \hookrightarrow \Delta_{l+1} \wr \langle S'_l, H''_l, F''_l \rangle$ for any l.

- (2) Any two elements $h' \in H'$ and $f' \in F'$ commute.
- (3) The group $\Gamma(S, H, F) < Aut(ET_{\bar{d}})$ is a quotient of the group $\Delta(S', H', F') < Aut(ET_{\bar{e}})$.

Property (2) shows that $\langle H', F' \rangle \simeq H' \times F'$. Thus as canonical generating set of $\Delta(S', H'F')$, use $S' \sqcup H'F'$.

Proof. (1) is obvious by construction. For (2), compute (recall the support of f'' and h'' is in $\{d_0 + 1, \ldots, d_0 + d'_0\}$):

$$[h', f'] = ([h'_1, f'_1], [\sigma'_2, 1], \dots, [\sigma'_{d_0}, 1], 1, \dots, 1)[h'', f''] = ([h'_1, f'_1], 1, \dots, 1).$$

By recursion, this shows that [h', f'] has a trivial action on $T_{\bar{e}}$. Moreover, the support of the corresponding function $\partial T_{\bar{e}} \to F$ is contained in 1^{∞} where the value is $[1, f] = 1_F$. This shows $[h', f'] =_{\Delta} 1$.

For (3), observe that all the permutations involved in the description of S', H', F' respect the decomposition $\{1, \ldots, d_l\} \sqcup \{d_l + 1, \ldots, d_l + d'_l\}$ for all $l \in \mathbb{N}$, so that the subset $ET_{\bar{d}} \subset ET_{\bar{e}}$ is stable under the action of $\Delta(S', H', F')$. The quotient action is precisely given by $\Gamma(S, HF)$.

Observe that if $d_l = 0$ for all $l \ge 1$ then $\Delta(S', H'F') \simeq \Gamma(S, HF) \times \langle S'', H''F'' \rangle$. In particular, if $d'_l = 0$ for all $l \in \mathbb{N}$ then $\Delta(S', H'F') = \Gamma(S, HF)$.

A word w(S, HF) is an ordered product of generators s in S, h in H and φ_f in F of the group Γ . By replacing s by s', h by h' and φ_f by f', one naturally obtains a word w(S', H'F') in the generators of Δ . The rewriting process of proposition 3.2 applies to the groups $\Delta(S', H'F')$. More precisely:

Proposition 7.2 (Rewriting process for groups $\Delta(S', H'F')$). If the rewriting process in the group $\Gamma \hookrightarrow \Gamma_1 \wr S_{d_0}$ provides:

$$w(S, HF) = (w^{1}(S_{1}, H_{1}F), \dots, w^{d_{0}}(S_{1}, H_{1}F))\sigma_{\emptyset},$$

then the rewriting process in the group $\Delta \hookrightarrow \Delta_1 \wr S'$ provides:

$$w(S', H'F') = (w^{1}(S'_{1}, H'_{1}F'_{1}), \dots, w^{d_{0}}(S'_{1}, H'_{1}F'_{1}), 1, \dots, 1)w(S', H''F''),$$

where $w(S', H''F'') = \sigma_{\emptyset}w(S'', H''F'')$ with $\sigma_{\emptyset} \in S_{\{1,...,d_0\}}$ and w(S'', H''F'') takes values in $S_{\{d_0+1,...,d_0+d'_0\}}$.

Proof. Denote $w(S, HF) = s_1 h_1 \varphi_{f_1} s_2 \dots h_n \varphi_{f_n} s_{n+1}$. Then

$$w(S', H'F') = s'_1 h'_1 f'_1 s'_2, \dots h'_n f'_n s'_{n+1},$$

where $s \mapsto s'$, $h \mapsto h'$ and $\varphi_f \mapsto f'$ are given in the definition of generators of Δ . These forms are equivalent to:

$$w(S, HF) = (h_1 \varphi_{f_1})^{\sigma_1} \dots (h_n \varphi_{f_n})^{\sigma_n} \sigma_{n+1}, \quad \text{for } \sigma_i = s_1 \dots s_i, w(S', H'F') = (h'_1 f'_1)^{\sigma'_1} \dots (h_n f'_n)^{\sigma'_n} \sigma'_{n+1}, \quad \text{for } \sigma'_i = s'_1 \dots s'_i.$$

For $h\varphi_f = (h_1\varphi_f, b_2, \dots, b_{d_0})$, one has:

$$(h\varphi_f)^{\sigma} = (b_{\sigma^{-1}(1)}, \dots, h_1\varphi_f, \dots, b_{\sigma^{-1}(d_0)})$$

with $h_1\varphi_f$ in position $\sigma(1)$.

Correspondingly, for $h'f' = (h'_1f'_1, b'_2, \dots, b'_{d_0}, 1, \dots, 1)h''f''$, one has:

$$(h'f')^{\sigma'} = (b'_{\sigma^{-1}(1)}, \dots, b'_1f'_1, \dots, b'_{\sigma^{-1}(d_0)}, 1, \dots, 1)(h''f'')^{\sigma''},$$

with $h'_1 f'_1$ in position $\sigma^{-1}(1) = \sigma'^{-1}(1)$, because $\sigma' = \sigma \sigma''$ with $\sigma \in S_{\{1,\dots,d_0\}}$ and $\sigma'' \in S_{\{d_0+1,\dots,d_0+d'_0\}}$.

As $(h''f'')^{\sigma''}$ acts trivially on $\{1,\ldots,d_0\}$, one can compute products (words) componentwise, which proves the proposition.

Remark 7.3. Proposition 7.2 shows in particular that the minimal tree description of section 3.2 and figure 1 for a word w(S, HF) in $\Gamma(S, HF)$ remains valid for the word w(S', H'F') in $\Delta(S', H'F')$ with the same tree $T(w) < T_{\bar{d}} < T_{\bar{e}}$, but for a vertex u in level l, the permutation σ_u now takes values in $S_l \times \langle S_l'', H_l''F_l'' \rangle < S_{e_l}$.

7.3. **Localisation.** A ball of given radius in the Cayley graph of $\Delta(S', H'F')$ depends only on the description of the action on a neighbourhood of the root of the tree.

Proposition 7.4. The ball $B_{\Delta}(R)$ in the Cayley graph of $\Delta(S', H'F')$ with respect to the generating set $S' \sqcup H'F'$ depends only on the $L = 1 + \log_2 R$ first levels in the recursive description of the generators of $\Delta(S', H'F')$ and the 2R + 1-balls in the groups $\langle S''_l, H''_lF''_l \rangle < S_{d'_l}$ for $l \leq L$.

Proof. The ball $B_{\Delta}(R)$ can be drawn if one can test (algorithmically) the oracle $w =_{\Delta} 1$ for any given word w in $S' \sqcup H'F'$ of length $\leq r = 2R + 1$. To test such an oracle, use the following algorithm.

First test the permutation value of w at the root $\Phi(w(s',h'f')) = w(s',h''f'') =_{S_{e_0}} 1_{S_{e_0}}$, where $\Phi: Aut(ET_{\bar{e}}) \to S_{e_0}$ is the root evaluation. This test depends only on the zero level in $\Delta(S',H'F')$ and the |w|-ball in $\langle S''_0,H''_0F''_0\rangle < S_{d'_0}$.

If $\Phi(w) \neq_{S_{e_0}} 1$ then $w \neq_{\Delta} 1$. If $\Phi(w) =_{S_{e_0}} 1$, then w fixes all vertices in the first level of $T_{\bar{e}}$. By propositions 3.2 and 7.2 of rewriting process, one has $w = (w^1, \ldots, w^{d_0}, 1, \ldots, 1)$ in the wreath product with $|w^t| \leq \frac{|w|+1}{2} \leq \frac{r+1}{2}$.

Test for each t in $\{1, \ldots, e_0\}$, the permutation value at the root $\Phi(w^t(s_1', h_1'f_1')) = w(s_1', h_1''f_1'') =_{S_{e_1}} 1_{S_{e_1}}$, which depends only on the zero level in $\Delta(S_1', H_1'F_1')$ hence on the first level in $\Delta(S_1', H_1'F_1')$ and the $|w^t|$ -ball in $\langle S_1'', H_1''F_1'' \rangle < S_{d_1'}$.

If $\Phi(w^t) \neq_{S_{e_1}} 1$, then $w \neq_{\Delta} 1$. If $\Phi(w^t) =_{S_{e_0}} 1$ for all t, then w fixes all vertices in the two first levels of $T_{\bar{e}}$. By propositions 3.2 and 7.2 of rewriting process, $w^t = (w^{t_1}, \dots, w^{t_{e_0}}, 1, \dots, 1)$ in the wreath product with $|w^{t_s}| \leq \frac{|w^t|+1}{2} < \frac{r}{4} + 1$.

Continue the process and test the value at the root of the words $w^{t_1...t_l}$ while their length is ≥ 1 . This test depends only on the l first levels in $\Delta(S', H'F')$ and the $|w^{t_1...t_l}|$ -ball in $\langle S''_l, H''_l F''_l \rangle \langle S_{d'_l}$. If $\Phi(w^{t_1...t_l}) \neq_{S_{e_l}} 1$ for some $t_1...t_l$, then $w \neq_{\Delta} 1$.

For $L=\log_2 r$, one has $|w^{t_1\dots t_L}|<\frac{r}{2^L}+1=2$, so $w^{t_1\dots t_L}$ is a generator in $\Delta_L(S'_L,H'_LF'_L)$. This implies that if $\Phi(w^{t_1\dots t_l})=_{S_{e_l}}1$ for all $t_1\dots t_l,\ l\leq L-1$ and $w^{t_1\dots t_L}=_{\Delta_L}1$, then $w=_{\Delta}1$.

The algorithm allows to test the oracle $w =_{\Delta} 1$ using only the data in the $L = \log_2 R$ first levels of $\Delta(S', H'F')$ and the $\frac{r}{2^l}$ -ball in the group $\langle S''_l, H''_l F''_l \rangle < S_{d'_l}$ for $l \leq \log_2 r$.

7.4. Asymptotic properties. The asymptotic description of $\Delta(S', H'F')$ is well understood in two extreme cases.

Proposition 7.5. [Low asymptotic] If $d'_l = 0$ for $l \ge L + 1$ and d'_l finite for $l \le L$, the quotient homomorphism (of restriction of the action to $ET_{\bar{d}} \subset ET_{\bar{e}}$):

$$\begin{array}{ccc} f: \Delta(S',H'F') & \to & \Gamma(S,HF) \\ s' & \mapsto & s \\ h' & \mapsto & h \\ f' & \mapsto & \varphi_f \end{array}$$

has finite kernel.

In particular, for the random walks Y_n in $\Gamma(S, HF)$ of law μ equidistributed on SHFS and Y'_n in $\Delta(S', H'F')$ of law μ' equidistributed on S'H'F'S', there exists C, K > 0 such that for all n:

- (1) $P(Y'_n =_{\Delta} 1) \le P(Y_n =_{\Gamma} 1) \le CP(Y'_n =_{\Delta} 1),$
- (2) $L_{\Gamma,\mu}(n) \leq L_{\Delta,\mu'}(n) \leq L_{\Gamma,\mu}(n) + K$,
- (3) $H_{\Gamma,\mu}(n) \le H_{\Delta,\mu'}(n) \le H_{\Gamma,\mu}(n) + \log C$.

Proof. An element δ in the kernel ker $f = \{\delta \in \Delta | \delta|_{ET_{\bar{d}}} = 1\}$ is described by its action on $ET_{\bar{e}} \setminus ET_{\bar{d}}$. By rewriting process, $\delta = (\delta_1, \ldots, \delta_{d_0}, 1, \ldots, 1)\delta''$ with $\delta'' \in S_{d_0+1,\ldots,d_0+d'_0} < S_{e_0}$, for which there are $\leq \#\langle S'', H''F'' \rangle$ choices. To describe δ , there remains to describe $\delta_1, \ldots, \delta_{d_0}$ that belong to the kernel $\ker(f_1 : \Delta_1 \to \Gamma_1)$.

For each $t \in \{1, \ldots, d_0\}$, the element δ_t has the form $\delta_t = (\delta_{t1}, \ldots, \delta_{td_1}, 1, \ldots, 1)\delta_t''$ with $\delta_t'' \in S_{d_1+1,\ldots,d_1+d_1'} < S_{e_1}$, for which there are $\leq \#\langle S_1'', H_1'' F_1'' \rangle$ choices.

By induction, the element $\delta \in \ker f$ is described by:

$$\{\delta''_{t_1...t_l}|t_i\in\{1,\ldots,d_i\},l\leq L\},\$$

for which there are $C \leq \#\langle S'', H''F'' \rangle (\#\langle S_1'', H_1''F_1'' \rangle)^{d_0} \dots (\#\langle S_L'', H_L''F_L'' \rangle)^{d_0 \dots d_{L-1}}$ possible choices.

Denote $f^{-1}(1_{\Gamma}) = \{\delta_1, \dots, \delta_C\}$, then $f^{-1}(\gamma) = \{\delta_1, \dots, \delta_C\}$ when $f(\gamma) = \delta$ and:

$$\mu^{*n}(\gamma) = P(Y_n =_{\Gamma} \gamma) = \sum_{i=1}^{C} P(Y_n' =_{\Delta} \delta \delta_i) = \sum_{\delta' \in f^{-1}(\gamma)} \mu'^{*n}(\delta').$$

For $\gamma = 1$, this guarantees (1):

$$P(Y_n' =_{\Delta} 1) \le P(Y_n =_{\Gamma} 1) \le CP(Y_n' =_{\Delta} 1),$$

because $P(Y_n =_{\Delta} \delta)$ is maximal for $\delta = 1_{\Delta}$.

One also has $||\delta|| \leq ||f(\delta)|| + K$ for $K = \max\{||\delta_1||, \dots, ||\delta_C||\}$. Indeed, if $w(S, HF) = \gamma = f(\delta)$, then $w(S', H'F') = \delta \delta_i$ for some i and when $\delta_i = w_i(S', H'F')$ of length $\leq K$, one has $ww_i^{-1}(S', H'F') = \delta$. This shows (2):

$$\mathbb{E}_{\mu^{*n}}||\gamma|| \leq \mathbb{E}_{\mu^{*n}}||\delta|| \leq \mathbb{E}_{\mu^{*n}}||\gamma|| + K.$$

Now fix n, and define for any γ in Γ the measure with support in $f^{-1}(\gamma) \subset \Delta$:

$$\nu_{\gamma}(\delta) = \begin{cases} \frac{\mu'^{*n}(\delta)}{\sum_{\delta' \in f^{-1}(\gamma)} \mu'^{*n}(\delta')} & \text{if } f(\delta) = \gamma, \\ 0 & \text{if } f(\delta) \neq \gamma. \end{cases}$$

Then the measure μ'^{*n} decomposes as:

$$\mu'^{*n} = \sum_{\gamma \in \Gamma} \mu^{*n}(\gamma) \nu_{\gamma},$$

and by lemma A.4 in [BKN] on conditionnal entropy:

$$H(\mu'^{*n}) \le \sum_{\gamma \in \Gamma} \mu^{*n}(\gamma) H(\nu_{\gamma}) + H(\mu^{*n}).$$

The support of ν_{γ} has size $\leq C$ so $H(\nu_{\gamma}) \leq \log C$, which shows (3):

$$H(\mu^{*n}) \le H(\mu'^{*n}) \le \log C + H(\mu^{*n}).$$

Proposition 7.6. [High asymptotic] If there exists l such that $d'_l = \infty$ and

$$S_{\infty} > \langle S_l'', H_l'' F_l'' \rangle \simeq S_l * (H_l \times F_l)$$

is non-amenable, then $\Delta(S', H'F') < Aut(ET_{\bar{e}})$ is non-amenable.

In particular, for the random walk Y_n of law μ equidistributed on the finite generating set S'H'F'S', there exists c > 0 such that for n large enough:

- (1) $P(Y_n =_{\Delta} 1) \leq e^{-cn}$
- (2) $L_{\Delta,\mu'}(n) \geq cn$,
- (3) $H_{\Delta,\mu'}(n) \geq cn$.

The proof will use the following:

Fact 7.7. Given $\gamma_1 \in \Delta_1$, there exists $\gamma \in \Delta$ such that $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_{e_0})id_{S'}$ for some $\gamma_2, \dots, \gamma_{e_0}$ in Δ_1 .

Note that this fact implies that Δ is infinite, for it contains γs for all s in S' and then $\#\Delta \geq \#S'\#\Delta_1 \geq \#S'\#S_1'\#\Delta_2 \geq \ldots$ This is in particular an elementary proof that directed saturated groups are infinite.

Proof of fact 7.7. Let $\gamma_1 = x_1 \dots x_r$ be a representative word in $S'_1 \sqcup H'_1 F'$. By saturation of H, there exists for each x_i , an element h'_i in H' such that $h'_i = (*, \dots, x_i, \dots, *)$ (where * marks some unknown value) with x_i in position 1 if $x_i \in H'_1 F'$ and x_i in some position between 2 and d_0 if $x_i \in S'_1$. Now by saturation of Γ , $S = S_{d_0}$ so S' acts transitively on $\{1, \dots, d_0\}$, and there exists $s_i \in S'$ such that $y_i = s_i h_i s_i^{-1} = (x_i, *, \dots, *)$. Then $\gamma = y_1 \dots y_r = (x_1 \dots x_r, *, \dots, *) = (\gamma_1, \gamma_2, \dots, \gamma_{e_0})$. (Note that in fact $\gamma_{d_0+1} = \dots = \gamma_{e_0} = 1$.)

Proof of proposition 7.6. The fact shows that the composition

$$St_1(\Delta) \hookrightarrow \Delta_1 \times \ldots \Delta_1 \xrightarrow{pr_1} \Delta_1$$

is surjective, so that if Δ_1 is non-amenable, so is $St_1(\Delta)$ which is a subgroup of Δ , and thus Δ is non-amenable. Iterating the process shows that if Δ_l is non-amenable, so is Δ . Consequence (1) follows by Kesten theorem [Kes], (2) and (3) by a theorem of Kaimanovich-Vershik [KV].

7.5. **High order oscillations.** The following theorems are similar to theorem 7.1 in [Bri3] on oscillation of growth functions (see also chapter 2 in [Bri1] and [KP]). The entropy function of the groups Δ involved is not precisely evaluated, but only some (rare) values of the function. The idea is to use alternatingly localisation and asymptotic evaluation to obtain a group with low entropy at some scales and high entropy at other scales.

Theorem 7.8. Let $\Gamma(S, HF) < Aut(ET_{\bar{d}})$ be a saturated directed group with measure μ equidistributed on SHFS. Let $h_1(n), h_2(n)$ be functions such that $\frac{h_1(n)}{H_{\Gamma,\mu}(n)} \to \infty$ and $\frac{h_2(n)}{n} \to 0$. Then there exists a group $\Delta(S', H'F') < Aut(ET_{\bar{e}})$ such that the entropy for the measure μ' equidistributed on S'H'F'S' satisfies:

- (1) $H_{\Gamma,\mu}(n) \leq H_{\Delta,\mu'}(n) \leq Cn$ for all n and a constant C,
- (2) $H_{\Delta,\mu'}(n_i) \leq h_1(n_i)$ for an infinite sequence (n_i) ,
- (3) $H_{\Delta,\mu'}(m_i) \geq h_2(m_i)$ for an infinite sequence (m_i) .

Corollary 7.9. For any $\frac{1}{2} \le \alpha \le 1$, there exists a finitely generated group Δ and a finitely supported symmetric measure μ' such that:

$$\underline{h}(\Delta, \mu') = \alpha \text{ and } \overline{h}(\Delta, \mu') = 1.$$

Corollaries 5.4, 5.6 and 7.9 imply theorem 1.2.

Proof of corollary 7.9. If $\alpha = 1$, take Δ any non-amenable group. If $\alpha < 1$, apply theorem 7.8 with $h(\Gamma, \mu) = \alpha$ (exists by corollary 5.4), $h_1(n) = H_{\Gamma,\mu}(n) \log n$ and $h_2(n) = \frac{n}{\log n}$.

Proof of theorem 7.8. The first condition is trivially satisfied since $\Gamma(S, HF)$ is a quotient of $\Delta(S', H'F')$.

Construct the group $\Delta(S', H'F') < Aut(ET_{\bar{e}})$ with a sequence $(l_i)_i$ rapidly increasing such that $d'_l = 0$ when $l \notin \{l_i\}_i$ and $(d'_{l_i})_i$ is rapidly increasing such that the group $\langle S''_{l_i}, H''_{l_i}F''_{l_i} \rangle < S_{d'_{l_i}}$ is a big finite quotient of the free product $S''_{l_i} * H''_{l_i}F''_{l_i}$.

Roughly, as (l_i) is rapidly increasing, there are scales at which an observer has the impression that $\Delta(S', H'F')$ resembles the group $\Gamma(S, HF)$ and has low asymptotic. As d'_{l_i} is big, there are scales at which an observer has the impression that $\Delta(S', H'F')$ contains a free group and has high asymptotic.

More precisely, assume by induction that parameters l_j, d'_{l_j} and integers $m_{j-1} \le n_j \le m_j$ are constructed for j < i together with an integer $k_{i-1} = 1 + \log_2 r_{i-1} \ge l_{i-1}$ such that $\{H(m)|m \le m_{i-1}\}$ depends only on $B(r_{i-1})$. By localisation 7.4, this ball

hence the values of the m_{i-1} first values of the entropy function depend uniquely on l, d'_l and the groups $\langle S''_l, H''_l F''_l \rangle < S_{d'_l}$ for $l \leq k_{i-1}$.

Now construct $l_i, d'_{l_i}, n_i, m_i, k_i$ by describing the sequence of groups $\langle S''_l, H''_l F''_l \rangle < S_{d'_l}$ for $k_{i-1} < l \le k_i$.

The group $\underline{\Delta}_i(\Gamma', H'F')$ where $d'_l = 0$ for all $l \geq k_i$ has low asymptotic by proposition 7.5, so:

(7)
$$H_{\Delta_i,\mu'}(n) \le H_{\Gamma,\mu}(n) + \log C$$

for some C, and in particular, there exists n_i such that:

$$H_{\underline{\Delta}_i,\mu'}(n_i) \leq h_1(n_i).$$

By localisation 7.4 this value of entropy depends only on a ball of radius R_i in the Cayley graph of $\underline{\Delta}_i(\Gamma', H'F')$ which depends only on the $L_i = 1 + \log_2 R_i$ first levels.

Now fix $l_i = L_i + 1$ and let $\overline{\Delta}_i(S', H'F')$ be the group with d'_l and $\langle S''_l, H''_l F''_l \rangle < S_{d'_l}$ as above for $l \leq l_i - 1$ and fix (momentarily) $d'_{l_i} = \infty$, with:

$$\langle S_{l_i}'', H_{l_i}'' F_{l_i}'' \rangle \simeq S_{l_i}'' * H_{l_i}'' F_{l_i}'' \simeq S_l * H_l F < S_{\infty},$$

and $d'_l = 0$ for $l > l_i$. The group $\overline{\Delta}_i(S', H'F')$ is non-amenable by proposition 7.6 of high asymptotic, so:

(8)
$$H_{\overline{\Delta}_{i},\mu'}(m) \ge cm$$

for some c > 0 and in particular, there exists m_i such that:

$$H_{\overline{\Delta}_i,\mu'}(m_i) \geq h_2(m_i).$$

Now by localisation 7.4, the m_i first values of the entropy function depend only on a ball of radius r_i in the Cayley graph of $\overline{\Delta}_i(S', H'F')$, which depends only on the $k_i = 1 + \log_2 r_i$ first levels, and the balls of radius $2r_i + 1$ in $\langle S_l'', H_l''F_l'' \rangle < S_{d_l'}$ for $l \leq k_i$.

In particular, the values $\{H(m)|m \leq m_i\}$ are the same if $\langle S_{l_i}'', H_{l_i}'' F_{l_i}'' \rangle$ is any group coinciding with the free product $S_{l_i}'' * H_{l_i}'' F_{l_i}'' \simeq S_l * H_l F$ in a ball of radius $2r_i + 1$. As a free product of finite groups is residually finite, there exists such a group which is finite of size d_{l_i}' .

The parameters $l_i, d'_{l_i}, k_i \geq l_i$, the finite groups $\langle S''_l, H''_l F''_l \rangle < S_{d'_l}$ for $l \leq k_i$ and the integers n_i, m_i are now constructed for all i by induction.

The sequence $(d'_{l_i})_i$ of positive integers and the finite groups $\langle S''_{l_i}, H''_{l_i} F''_{l_i} \rangle < S_{d'_{l_i}}$ permit to define a group Δ for which $H_{\Delta,\mu'}(n_i) \leq h_1(n_i)$ and $H_{\Delta,\mu'}(m_i) \geq h_2(m_i)$ for all i, because the balls of radius r_i in Δ and in $\overline{\Delta}_i$ coincide.

In the point of view of information theory of remark 5.3, the minimal tree description remains valid. However, the number of digits needed to describe σ_u is not anymore bounded independently of the level l, for now $d_{l_i} \to \infty$, which explains the higher values taken by the entropy.

Theorem 7.10. Let $\Gamma(S, HF) < Aut(ET_{\bar{d}})$ be a saturated directed group with random walk Y_n of law μ equidistributed on SHFS. Let $p_1(n), p_2(n)$ be functions such

that $\frac{p_1(n)}{P(Y_n=\Gamma^1)} \to 0$ and that for any c>0 and n large, $p_2(n) \geq e^{-cn}$. Then there exists a group $\Delta(S', H'F') < Aut(ET_{\bar{e}})$ such that the return probability for the random walk Y'_n with law μ' equidistributed on S'H'F'S' satisfies:

- (1) $P(Y_n =_{\Gamma} 1) \ge P(Y'_n =_{\Delta} 1) \ge e^{-Cn}$ for all n and a constant C, (2) $P(Y'_{n_i} =_{\Delta} 1) \ge p_1(n_i)$ for an infinite sequence (n_i) , (3) $P(Y'_{n_i} =_{\Delta} 1) \le p_2(m_i)$ for an infinite sequence (m_i) .

Theorem 7.11. Let $\Gamma(S, HF) < Aut(ET_{\bar{d}})$ be a saturated directed group with measure μ equidistributed on SHFS. Let $L_1(n), L_2(n)$ be functions such that $\frac{L_1(n)}{L_{\Gamma,\mu}(n)} \to$ ∞ and $\frac{L_2(n)}{n} \to 0$. Then there exists a group $\Delta(S', H'F') < Aut(ET_{\bar{e}})$ such that the drift for the measure μ' equidistributed on S'H'F'S' satisfies:

- (1) $L_{\Gamma,\mu}(n) \leq L_{\Delta,\mu'}(n) \leq Cn$ for all n and a constant C,
- (2) $L_{\Delta,\mu'}(n_i) \leq L_1(n_i)$ for an infinite sequence (n_i) ,
- (3) $L_{\Delta,\mu'}(m_i) \geq L_2(m_i)$ for an infinite sequence (m_i) .

Proof. The proof of theorem 7.8 applies to theorems 7.10 and 7.11, with (a priori) different choices of parameters l_i, d'_{l_i} and integers n_i, m_i , obtained by replacing inequality (7) by (proposition 7.5 of low asymptotic):

(9)
$$P(Y_n =_{\Delta} 1) \ge \frac{1}{C} P(Y_n =_{\Gamma} 1) \ge p_1(n)$$
, or $L_{\Delta,\mu'}(n) \le L_{\Gamma,\mu}(n) + K \le L_1(n)$,

for n large enough, which permits to find n_i , and replacing inequality (8) by (proposition 7.6 of high asymptotic):

(10)
$$P(Y_m =_{\Delta} 1) \le e^{-cm} \le p_2(m)$$
, or $L_{\Delta,\mu'}(m) \ge cm \ge L_2(m)$, for m large enough, which permits to find m_i .

Corollary 7.12. [theorem 1.3] There exists a finitely generated group Δ and a finite symmetric measure μ' such that the return probability of the random walk Y'_n of law μ' satisfies:

$$\underline{p}(\Delta) = \frac{1}{3}, \overline{p}(\Delta) = 1 \text{ and } \underline{\delta}(\Delta, \mu') = \frac{1}{2}, \overline{\delta}(\Delta, \mu') = 1.$$

Proof. Take $\Gamma(S, HF) = F \wr_{\partial T_2} D_{\infty} < Aut(ET_2)$ to be the directed saturated group of proposition 6.4, for which $e^{-n^{\frac{1}{3}-\varepsilon}} \geq P(Y_n =_{\Gamma} 1) \geq e^{-n^{\frac{1}{3}}}$ and $L_{\Gamma,\mu}(n) \approx n^{\frac{1}{2}}$. Take $p_1(n) = e^{-n^{\frac{1}{3}} \log n}$, $p_2(n) = e^{-\frac{n}{\log n}}$, $L_1(n) = n^{\frac{1}{2}} \log n$ and $L_2(n) = \frac{n}{\log n}$. Apply theorems 7.10 and 7.11.

8. Generalisation

The definition of a saturated directed group $\Gamma(S, HF)$ acting on an extended tree $ET_{\bar{d}}$ for a bounded sequence d can be slightly generalised, with adaptation of theorems 5.1 and 6.1.

By S_d denote a finite group acting faithfully and transitively on $\{1,\ldots,d\}$ (not necessarily the full group of permutation). Replace definition (3) in section 2.1 by:

$$h = (h_1, \sigma_2, \ldots, \sigma_{d_0})\sigma_h$$

with σ_h in $Fix_{S_{d_0}}(1)$, a permutation of $\{1,\ldots,d_0\}$ that fixes 1. By recursion, $\sigma_{h_l}(1) = 1$ for all l, hence h fixes the infinite ray 1^{∞} , and thus commutes with φ_f . The group $H_{\bar{d}}$ directed by the ray 1^{∞} is the uncountable locally finite group:

$$H_{\bar{d}} = AT_0 \times AT_1 \times \dots$$
, where $AT_l = S_{d_{l+1}} \wr S_{d_l-1} = (\underbrace{S_{d_{l+1}} \times \dots \times S_{d_{l+1}}}_{d_l-1 \text{ factors}}) \rtimes S_{d_l-1}$.

Given another sequence $\bar{c} = (c_l)_l$ of integers such that $1 \leq c_l \leq d_l - 1$, define the subgroup PT_l by:

$$PT_{l} = \underbrace{S_{d_{l+1}} \times \cdots \times S_{d_{l+1}}}_{c_{l} \text{ factors}} \times \{1\} \times \cdots \times \{1\} < S_{d_{l+1}} \wr S_{d_{l-1}} = AT_{l},$$

with c_l factors $S_{d_{l+1}}$ when $c_l < d_l - 1$, and $PT_l = AT_l$ if $c_l = d_l - 1$. The hypothesis of saturation of a group G(S, H) can be relaxed as relative saturation with respect to \bar{c} by requiring that $S = S_{d_0}$ acts transitively on $\{1, \ldots, d_0\}$ and the subgroup $H < H_{\bar{d}}$ is included in:

$$H < PT_0 \times PT_1 \times \dots$$

with the equidistribution measure on H projecting to equidistribution measure on each factor PT_l .

Given the sequences $d = (d_l)_l$ and $\bar{c} = (c_l)_l$ of integers with $1 \le c_l \le d_l - 1$, define a new sequence $\bar{p'} = (p'_l)_l$ by $p'_l = \frac{c_l}{(c_l+1)d_l}$. and set:

$$\beta_{\bar{d},\bar{c}}(n) = \frac{\log(d_0...d_{k(n)})}{\log n}, \quad \text{where } k(n) = k_{\bar{d},\bar{c}}(n) = \min\{k|p_0'\dots p_k'n \leq 1\},\\ \beta_{\bar{d},\bar{c}}'(n) = \frac{\log(d_0...d_{l(n)})}{\log n}, \quad \text{where } l(n) = l_{\bar{d},\bar{c}}(n) = \max\{l|\frac{d_0}{p_0'}\dots\frac{d_l}{p_l'} \leq n\}.$$

With these notations, theorems 5.1 and 6.1 generalise to:

Theorem 8.1. Given bounded sequences \bar{d} and \bar{c} , a relatively saturated directed group $\Gamma(S, HF)$ and the measure μ of equidistribution on SHFS, one has for arbitrary $\varepsilon > 0$ and n large:

- $(1) \left| \frac{\log H_{\Gamma,\mu}(n)}{\log n} \beta_{\bar{d},\bar{c}}(n) \right| \leq \varepsilon,$ $(2) \beta_{\bar{d},\bar{c}}(n) \varepsilon \leq \frac{\log L_{\Gamma,\mu}(n)}{\log n} \leq \frac{1 + \beta_{\bar{d},\bar{c}}(n)}{2} + \varepsilon,$ $(3) \beta'_{\bar{d},\bar{c}}(n) \varepsilon \leq \frac{\log \log P(Y_n = \Gamma^1)}{\log n} \leq \beta_{\bar{d},\bar{c}}(n) + \varepsilon.$

For constant sequences $d_l = d$ and $c_l = c \leq d - 1$, the sequences $\beta_{\bar{d},\bar{c}}(n)$ and $\beta'_{\bar{d},\bar{c}}(n)$ have limits respectively:

$$\beta_{d,c} = \frac{\log d}{\log(\frac{d(c+1)}{c})} = \frac{1}{1 + \frac{\log(\frac{c+1}{c})}{\log d}}, \quad \text{so } \beta_{d,1} = \frac{1}{1 + \frac{\log 2}{\log d}},$$
$$\beta'_{d,c} = \frac{\log d}{\log(\frac{d^2(c+1)}{c})} = \frac{1}{2 + \frac{\log(\frac{c+1}{c})}{\log d}}, \quad \text{so } \beta'_{d,1} = \frac{1}{2 + \frac{\log 2}{\log d}}.$$

(1) In the case of constant valency d, for $H = AT(d) = S_d$ Examples 8.2. S_{d-1} diagonally embedded into $AT_0 \times AT_1 \times \dots$, obtain the mother group $G(S_d, AT(d))$ of polynomial automata of degree 0 (see [AAV], [BKN]). With an extension F at the tree boundary, one has for $\Gamma(d) = \Gamma(S_d, AT(d)F) <$ $Aut(ET_d)$ the estimates:

$$h(\Gamma(d), \mu) = \beta_d$$
 and $\beta'_d \leq \underline{p}(\Gamma(d), \mu) \leq \overline{p}(\Gamma(d), \mu) \leq \beta_d$.

(2) For the spinal groups $G_{\omega}(q)$ of the article [BS] acting on a tree of constant valency q = d. With an extension F at the tree boundary, one has for $\Gamma_{\omega}(q) = F \wr_{\partial T} G_{\omega}(q)$ the estimates:

$$h(\Gamma_{\omega}(q), \mu) = \beta_{q,1} \text{ and } \beta'_{q,1} \leq \underline{p}(\Gamma_{\omega}(q), \mu) \leq \overline{p}(\Gamma_{\omega}(q), \mu) \leq \beta_{q,1}.$$

Proof of theorem 8.1. Notation of proposition 3.2 and lemma 4.1 becomes:

$$k_j = (k'_j, b_{j,2}, \dots, b_{j,c_0+1}, 1, \dots, 1)\sigma_{h_j},$$

and one should consider $k_j^{\sigma_j}$ for $\sigma_j = s_1 \sigma_{h_1} s_2 \dots s_{j-1} \sigma_{h_{j-1}} s_j$. The sequence $(\sigma_j)_j$ consists of independent terms equidistributed in S_{d_0} . As its action is transitive on $\{1, \dots, d_0\}$, for any fixed t, the sequence $(\sigma_j(t))_j$ is a sequence of independent equidistributed elements of $\{1, \dots, d_0\}$.

This ensures that Y_n^t is a product of n terms that are:

- (1) either $b_{j,\sigma_j(t)}$ equidistributed in S_{d_1} at times j when $\sigma_j(t) \in \{2,\ldots,c_0+1\}$,
- (2) or k'_j equidistributed in H_1F at times j when $\sigma_j(t) = 1$,
- (3) or trivial factors 1 at times j when $\sigma_j(t) \in \{c_0 + 2, \dots, d_0\}$.

The number of non-trivial terms is $n_t \sim \frac{c_0+1}{d_0}n$ almost surely. Now merging packs of terms in the same finite group S_{d_1} (with probability $\frac{c_0}{c_0+1}$ among non-trivial terms) or H_1F (with probability $\frac{1}{c_0+1}$ among non-trivial terms), the length of Y_n^t is almost surely:

$$m_t \sim \frac{c_0}{(c_0+1)^2} n_t \sim \frac{c_0}{(c_0+1)d_0} n.$$

This shows that lemma 4.1 is true with $p'_0 = \frac{c_0}{(c_0+1)d_0}$ instead of $p_0 = \frac{d_0-1}{d_0^2}$.

Then lemma 4.5 generalises as:

$$\left| \frac{\log \mathbb{E}a(Y_n)}{\log n} - \beta_{\bar{d},\bar{c}}(n) \right| \le \varepsilon,$$

for $\varepsilon > 0$ and n large. Theorems 5.1 and 6.1 and corollary 5.10 follow straightforwardly. Note that the condition on θ in fact 6.2 becomes $p'_i - \theta > \frac{p'_i}{d_i}$.

9. Comments and Questions

9.1. Analogies between growth and entropy for directed groups. Analogies between growth and entropy for directed groups are two fold.

First, there is an analogy between the computation of growth exponents (as $\liminf \underline{\alpha}(\Gamma)$, $\liminf \underline{\alpha}(\Gamma)$, $\inf \overline{\alpha}(\Gamma)$ or $\liminf \underline{\alpha}(\Gamma)$ or $\inf \frac{\log \log b_{\Gamma}(r)}{\log r}$) in [Bri3] and the computation of entropy exponents in theorem 5.1. For entropy, the computation is based on the contraction by a factor p_0 of the word length under rewriting process of random alternate words in $\Gamma(S, HF)$. For growth in the extended Aleshin-Grigorchuk group $\Gamma_{(012)^{\infty}} = F \wr G_{(012)^{\infty}}$, the computation is based on the contraction by a factor $\frac{\eta}{2}$ in the wreath product for reduced representative words (see [BE], and lemma 5.4 in [Bri3]).

The contraction factor for entropy should only hold for random alternate words, whereas the contraction factor for growth has to hold for any alternate (pre-reduced) word, which heuristically explains why $\frac{1}{2} = h(\Gamma_{(012)^{\infty}}, \mu) < \alpha(\Gamma_{(012)^{\infty}}) \approx 0.76$. This inequality is a well-known property of Shannon entropy that $H(\mu) \leq \log \#supp(\mu)$ with equality for an equidistributed measure.

There is a second analogy at the level of parameter space. For a fixed bound D on the valency \bar{d} , the space of saturated directed groups is (partially) parametrised by the Cantor set $\{2, \ldots, D\}^{\mathbb{N}}$. The entropy exponent $\beta_{\bar{d}}(n)$ is computed in theorem 5.1 in terms of the sequence \bar{d} and the contraction factors $(p_i)_i$ as $n^{\beta_{\bar{d}}(n)} = d_0 \ldots d_{k(n)}$ where $p_0 \ldots p_{k(n)} \approx \frac{1}{n}$. By fact 4.4, any function g(n) such that $dg(n) \leq g(\frac{d^2}{d-1}n)$ and $g(\frac{D^2}{D-1}n) \leq Dg(n)$ is the entropy of some finitely generated group (with the approximation of theorem 5.1).

The space of extended Aleshin-Grigorchuk groups Γ_{ω} is also parametrized by a Cantor set $\{0,1,2\}^{\mathbb{N}}$. Though the growth function for a given sequence ω is not known, Bartholdi and Erschler have shown recently in [BE3] that any function $e^{g(n)}$ with $g(2n) \leq 2g(n) \leq g(\eta_+ n)$ for $\eta_+ \approx 2.46$ explicit is the growth function of some group (also compare corollary 4.2 in [BE3] with definition of exponent sequence at section 4.2).

9.2. Comparison between growth, entropy, return probability and drift. Among finitely generated groups with symmetric finitely supported measure, it is a natural question to classify the pairs $(\underline{\alpha}(\Gamma), \overline{\alpha}(\Gamma))$, $(\underline{h}(\Gamma, \mu), \overline{h}(\Gamma, \mu))$, $(\underline{p}(\Gamma), \overline{p}(\Gamma))$ and $(\underline{\delta}(\Gamma, \mu), \overline{\delta}(\Gamma, \mu))$ in the triangle $0 \le \alpha \le \beta \le 1$. Comparing theorem 1.3 with theorem 1.2 and the main result in [Bri3] raises the following two questions.

Question 9.1. Given a point (α, β) in the triangle $\frac{1}{3} \le \alpha \le \beta \le 1$, does there exist a finitely generated group Γ the return probability exponents of which satisfy:

$$(p(\Gamma), \overline{p}(\Gamma)) = (\alpha, \beta)$$
?

Question 9.2. Given a point (α, β) in the triangle $\frac{1}{2} \leq \alpha \leq \beta \leq 1$, does there exist a finitely generated group Γ together with a (symmetric finitely supported) measure μ , the drift exponents of which satisfy:

$$(\underline{\delta}(\Gamma, \mu), \overline{\delta}(\Gamma, \mu)) = (\alpha, \beta)$$
?

One approach would be to improve theorem 6.1 by understanding how a particular choice of H affects the return probability. Another approach is by the technics developed in [KP], that could lead to strengthening of theorems 7.8 and 7.10.

Another natural question is to know if there are such pairs outside the above mentioned triangles besides the pair (0,0), obtained by virtually nilpotent groups for growth, entropy and return probability, by finite groups for drift. By [LP], the number $\frac{1}{2}$ is a lower bound on the drift exponent of infinite groups. It raises the:

Question 9.3. Does there exist a group Γ and a measure μ such that:

$$\begin{array}{ll} 0<\underline{h}(\Gamma,\mu)<\frac{1}{2},\\ \text{or} & 0<\underline{p}(\Gamma)<\frac{1}{3},\\ \text{or} & 0<\underline{\alpha}(\Gamma)<\alpha(\Gamma_{(012)^{\infty}})\approx 0.76 \end{array}$$

By [CGP], groups of exponential growth have return probability exponents $\geq \frac{1}{3}$. By theorem 6.1, this is also a lower bound for many groups with intermediate growth. A conjecture of Grigorchuk asserts that a finitely generated group Γ is virtually nilpotent when its growth function satisfies $b_{\Gamma}(r) \leq e^{r^{\frac{1}{2}-\varepsilon}}$. If this were the case, then $\overline{p}(\Gamma) < \frac{1}{5}$ would imply virtual nilpotency by [CGP]. The bound $\frac{1}{2}$ for entropy corresponds to some simple random walk on the lamplighter group or on an extended directed Aleshin-Grigorchuk group (see remark 5.5).

Finally, one may wonder how these asymptotic quantities relate with each other. For instance entropy is bounded by logarithm of growth so $h(\Gamma, \mu) \leq \alpha(\Gamma)$. Also corollary 7.4 in [CGP] implies that:

$$\frac{\underline{\alpha}(\Gamma)}{2 + \underline{\alpha}(\Gamma)} \le \underline{p}(\Gamma) \text{ and } \overline{p}(\Gamma) \le \frac{\overline{\alpha}(\Gamma)}{2 - \overline{\alpha}(\Gamma)}.$$

Naively, one expects groups with low growth to have low return probability exponents and vice-versa. However, taking Γ the lamplighter on \mathbb{Z} and $\Gamma' = F \wr G_{\omega}(q)$ an extension of a spinal group $G_{\omega}(q)$ of the article [BS], one has (note that theorem 6.1 in [BS] still applies with boundary extension F and see exemple 8.2 (2)):

$$\overline{\alpha}(\Gamma') < \alpha(\Gamma) = 1 \text{ but } \frac{1}{3} = p(\Gamma) < \frac{1}{2 + \frac{\log 2}{\log q}} \le \underline{p}(\Gamma').$$

This raises the:

Question 9.4. For Γ finitely generated group with finitely supported symmetric measure μ , what are the possible values of the 8-tuple:

$$(\underline{\alpha}(\Gamma), \overline{\alpha}(\Gamma), \underline{p}(\Gamma), \overline{p}(\Gamma), \underline{h}(\Gamma, \mu), \overline{h}(\Gamma, \mu), \underline{\delta}(\Gamma, \mu), \overline{\delta}(\Gamma, \mu)) ?$$

References

[Ale] Aleshin S., Finite automata and Burnside's problem for periodic groups, Math. Notes 11 (1972), 319-328.

[AAV] Amir G., Angel O., Virag B., Amenability of linear-activity automaton groups, arXiv: 0905.2007v1 (2009).

[AV] Amir G., Virag B., Positive speed for high-degree automaton groups, arXiv: 1102.4979v1 (2011).

[BE] Bartholdi L., Erschler A., Growth of permutational extensions, arXiv: 1011.5266v1 (2010).

[BE2] Bartholdi L., Erschler A., Poisson-Furstenberg boundary and growth of groups, arXiv: 1107.5499v1 (2011).

[BE3] Bartholdi L., Erschler A., Groups of given intermediate word growth, arXiv: 1110.3650v1 (2011).

[BKN] Bartholdi L., Kaimanovich V. A., Nekrashevych V. V., On amenability of automata groups, Duke Math. J. 154 (2010), no. 3, 575-598.

[BS] Bartholdi L., Sunik Z., On the word and period growth of some groups of tree automorphisms, Comm. Algebra 29 (2001), no. 11, 4923-4964.

[BV] Bartholdi L., Virag B., Amenability via random walks, Duke Math. J. 130 (2005), no. 1, 39-56.

[Bri1] Brieussel J., Croissance et moyennabilité de certains groupes d'automorphismes d'un arbre enraciné, Thèse de doctorat, Université D. Diderot Paris 7, (2008). Available at http://www.institut.math.jussieu.fr/theses/2008/brieussel/.

[Bri2] Brieussel J., Amenability and non-uniform growth of some directed automorphism groups of a rooted tree, Math. Z. 263 (2009), no. 2, 265-293.

- [Bri3] Brieussel J., Growth behaviors in the range $e^{r^{\alpha}}$, arXiv: 1107.1632v1, (2011).
- [CCJJV] Cherix P., Cowling M., Jolissaint P., Julg P., Valette A., *Groups with the Haagerup property*, Progress in Maths 197, Birkhauser, 2001.
- [CGP] Coulhon T., Grigor'yan A., Pittet C., A geometric approach to on-diagonal heat kernel lower bounds on groups, Ann. Inst. Fourier (Grenoble) 51 (2001), no. 6, 17631827.
- [Ers1] Erschler A., On drift and entropy growth for random walks on groups, Ann. Probab. 31 (2003), no. 3, 11931204.
- [Ers2] Erschler A., On the asymptotics of the rate of departure to infinity, J. Math. Sci. (N. Y.) 121 (2004), no. 3, 24372440.
- [Ers3] Erschler A., On the degrees of growth of finitely generated groups, Funct. Anal. Appl. 39 (2005), no 4, 317-320. [Russian: Funktsional. Anal. i Prilhozen. 39 (2005), no 4, 86-89.]
- [Ers4] Erschler A., Piecewise automatic groups, Duke Math. J. 134 (2006), no. 3, 591613.
- [Ers5] Erschler A., Isoperimetric inequality for wreath product of Markov chains and selfintersections of random walks, Probab. Theory Relat. Fields 136(4), 560586 (2006).
- [Ers6] Erschler A., Poisson-Furstenberg boundaries, large-scale geometry and growth of groups, Proc. Int. Cong. Math., Vol II, 681-704, Hindustan Book Agency, New Delhi, 2010.
- [Fol] Folner E., On groups with full Banach mean value, Math. Scand. 3 (1955) 243-254.
- [Gri1] Grigorchuk R., Degrees of growth of finitely generated groups and the theory of invariant means, Math. USSR Izv. 25:2 (1985), 259-300.
- [Gri2] Grigorchuk R., Degrees of growth of p-groups and torsion free groups, Math. USSR-Sb. 54(1), 185-205 (1986). [Russian: Mat. Sb. (N.S.) 126(168) (1985), no.2, 194-214, 286.]
- [GZ] Grigorchuk R., Zuk A., On a torsion-free weakly branch group defined by a three state automaton, Internat. J. Algebra Comput., Vol. 12, No. 1 (2002), 1-24.
- [GS] Gupta N., Sidki S., On the Burnside problem for periodic groups, Math. Z. 182 (1983) 385-388.
- [Kai] Kaimanovich V., "Münchhausen trick" and amenability of self-similar groups, IJAC 15(5-6) (2005), 907-938.
- [KV] Kaimanovich V., Vershik A., Random walks on discrete groups: boundary and entropy, Ann. Probab. Volume 11, Number 3 (1983), 457-490.
- [KP] Kassabov M., Pak I., Groups of oscillating intermediate growth, arXiv: 1108.0262v1 (2011).
- [Kes] Kesten H., Full Banach mean values on countable groups, Math. Scand. 7 (1959), 146-156.
- [LP] Lee J., Peres Y., Harmonic maps on amenable groups and a diffusive lower bound for random walks, arXiv:0911.0274 (2009).
- [MP] Muchnik R., Pak I., On growth of Grigorchuk groups, Internat. J. Algebra Comput. 11 (2001), 1-17.
- [Nek] Nekrashevych V., Self-similar groups, volume 117 of Mathematical Surveys and Monographs. Amer. Math. Soc., Providence, RI, 2005.
- [PSC1] Pittet C., Saloff-Coste L., On the stability of the behavior of random walks on groups, J. Geom. Anal. 10 (2000), no. 4, 713-737.
- [PSC2] Pittet C., Saloff-Coste L., On random walks on wreath products, Ann. Probab. 30 (2002), no. 2, 948-977.
- [PSC3] Pittet C., Saloff-Coste L., Random walks on finite rank solvable groups, J. Eur. Math. Soc. 5 (2003), 313-342.
- [Rev] Revelle D., Rate of Escape of Random Walks on Wreath Products and Related Groups, Ann. Probab. 31 (2003), no. 4, 1917-1934.
- [Seg] Segal D., The finite images of finitely generated groups, Proc. London Math. Soc. (3) 82 (2001), no. 3, 597613.
- [Sha] Shannon C., A mathematical theory of communication, Bell System Tech. J. 27, (1948). 379423, 623656.
- [Sid] Sidki S., Finite automata of polynomial growth do not generate a free group, Geom. Dedicata 108 (2004), 193204.
- [Var] Varopoulos N., Théorie du potentiel sur les groupes nilpotents, C. R. Acad. Sci. Paris Sér. I Math. 301 (1985), 143-144.
- [Wil1] Wilson J., On exponential growth and uniformly exponential growth for groups, Invent. math. 155 (2004), 287-303.

[Wil2] Wilson J., Further groups that do not have uniformly exponential growth, Journal of Algebra 279 (2004), 292-301.

[Woe] Woess W., Random walks on infinite graphs and groups, Cambridge Tracts in Mathematics, 138, Cambridge University Press (2000).

[Zuk] Zuk A., Groupes engendrés par des automates, Séminaire N. Bourbaki 971, (2006).

 $E ext{-}mail\ address: jeremie.brieussel@gmail.com}$