

CRITICAL PERCOLATION OF VIRTUALLY FREE GROUPS AND OTHER TREE-LIKE GRAPHS

BY IVA KOZÁKOVÁ

Vanderbilt University

This article presents a method for finding the critical probability p_c for the Bernoulli bond percolation on graphs with the so-called tree-like structure. Such graphs can be decomposed into a tree of pieces, which have finitely many isomorphism classes. This class of graphs includes the Cayley graphs of amalgamated products, HNN extensions or general groups acting on trees. It also includes all transitive graphs with more than one end.

The idea of the method is to find a multi-type Galton-Watson branching process (with a parameter p), which has finite expected population size if and only if the expected percolation cluster size is finite. This provides sufficient information about p_c . In particular if the pairwise intersections of pieces are finite, then p_c is the smallest positive p , such that $\det(M - 1) = 0$, where M is the first-moment matrix of the branching process. If the pieces of the tree-like structure are finite, then p_c is an algebraic number, and we give an algorithm computing p_c as a root of some algebraic function.

We show that any Cayley graph of a virtually free group (that is, a group acting on a tree with finite vertex stabilizers) with respect to any finite generating set has a tree-like structure with finite pieces. In particular we show how to compute p_c of the Cayley graph of a free group with respect to any finite generating set.

Contents.

1	Introduction	2
2	Graphs with tree-like structure	5
	2.1 The definition	6
	2.2 Changing the generating set	7
3	Partitions of the border sets	10
4	Branching processes	12
	4.1 Preliminaries	12
	4.2 Coloring of the tree of pieces according to the percolation	14
	4.3 The complete branching process	15
	4.4 The branching process with a reduced number of types	16
	4.5 The case of finite border sets	19
5	Examples	22

MSC classes: 60K35, 60D05, 20P05

5.1	Free products of (transitive) graphs	22
5.2	$SL(2, \mathbb{Z})$	23
5.3	Grandparent tree	25
6	Fundamental groups of graphs of groups	25
6.1	Amalgamated products with standard generating sets	25
6.2	HNN extensions with standard generating sets	26
6.3	Graphs of groups	27
6.4	An example with infinite pieces	30
6.5	Groups acting on trees with finite vertex stabilizers	31
7	Transitive graphs with more than one end	33
7.1	Dunwoody's result	33
7.2	Application to free groups with non-standard generators	35
	Acknowledgments	38
	References	38
	Author's addresses	39

1. Introduction. We will use the notation $\mathcal{G} = (V, E)$ for a graph with the vertex set V and the edge set E . All graphs are assumed to be locally finite (vertices have finite degrees) and transitive (for any two vertices there exists an automorphism of \mathcal{G} mapping one to the other). We fix one vertex of the graph and call it the *origin*.

For every $p \in (0, 1)$, the *Bernoulli bond percolation* on \mathcal{G} is a product probability measure P_p on the space $\Omega = \{0, 1\}^E$, the subsets of the edge set E . The product measure is defined via $P_p(\omega(e) = 1) = p$ for all $e \in E$. The σ -algebra of P_p -measurable sets does not depend on p . We denote the σ -algebra by Σ and the expected value by E_p .

For any realization $\omega \in \Omega$, open edges form a random subgraph of \mathcal{G} . The *Percolation function* is defined to be the probability that the origin is contained in an infinite cluster. The behavior of the percolation model depends strongly on the value of probability p . There is a critical value p_c of the probability p such that for $0 \leq p < p_c$ all clusters are finite, and if $p_c < p \leq 1$ there is an infinite cluster P_p -almost surely (see Grimmett [4]).

Explicit values of p_c have been known only for some special cases. In particular, for lattices in \mathbb{R}^2 the value of p_c is obtained using dual graphs (for \mathbb{R}^d with $d \geq 3$ the values of p_c are not known). For the square lattice, Kesten ([5]) proved $p_c = 1/2$, for the triangular lattice $p_c = 2 \sin(\pi/18)$, and for the hexagonal lattice $p_c = 1 - 2 \sin(\pi/18)$ (see Grimmett [4]). Ziff and Scullard [11] found p_c for a larger class of lattices in \mathbb{R}^2 (they considered graphs that can be decomposed onto certain self-dual arrangements). The value of critical probability is also known for trees ($p_c = 1/\text{branching number}$) and

virtually cyclic groups, where $p_c = 1$. The critical probability can also be found in case of a free product of finite transitive graphs ([6]). As far as we know these are the only known graphs where p_c for the bond percolation has been computed exactly.

If we change the generating set of a group, the graph changes dramatically, and there have been no examples of groups (except virtually cyclic) where p_c was known for all generating sets. It is expected that many properties of percolation (behavior at the critical value) are invariant with respect to changing the generating set, so it is useful to know how p_c depends on the generating set of a group.

In this article we study the critical probability p_c for a class of graphs that admit the so-called *tree-like structure*. Roughly speaking, such a graph can be decomposed into a rooted tree of edge-disjoint pieces that intersect at the so-called border sets, which are cut sets of the graph. We always assume that there are finitely many isomorphism classes of pieces. This class of graphs includes, for example, all transitive graphs with more than one end, Cayley graphs of amalgamated products and HNN extensions (for example: $SL(2, \mathbb{Z}) = \mathbb{Z}_4 *_{\mathbb{Z}_2} \mathbb{Z}_6$). In the case of an amalgamated product the pieces correspond to the Cayley graphs of factor groups and the border sets consist of vertices from cosets of the amalgamated subgroup. The precise definition of the tree-like structure is contained in Section 2 and several examples, including the grandparent tree, can be found in Section 5.

Consider a realization of the percolation on a graph \mathcal{G} with a tree-like structure. For each piece P_i that is not the root, let B_i be the border set that is the intersection of P_i with its parent in the tree.

Since B_i is a cut set, we can define an equivalence relation on it, by saying that two vertices of B_i are equivalent if they are connected by an open path inside the union of all pieces that are not descendants of P_i . The data consisting of the equivalence relation and the distinguished equivalence class connected to the origin is called the *color* of the piece P_i . If no vertex of the border set is connected to the origin we say that this piece has white color. Since the p_c of the whole graph does not exceed the p_c of any subgraph, we can assume that the probability p is smaller than the minimum of all p_c 's of pieces. Then it is easy to see that the percolation process dies if and only if the tree of non-white pieces is finite. The fact that the pieces form a tree suggests using a branching process with individuals corresponding to the colored pieces, such that the distribution of children is induced by the percolation process. Unfortunately the color of a piece depends not only on the color of its parent (as required for branching process), but also on the colors of the siblings and their descendants.

Nevertheless, we define a different distribution on the colors of the children, and we obtain a branching process that has finite population size if and only if the expected size of the percolation cluster is finite. The next statement is the main result of the paper.

THEOREM 1.1. *Assume the graph \mathcal{G} has a tree-like structure.*

- (i) *For a percolation with parameter p there exist a branching process on the tree of pieces, such that the expected size of its population is finite if and only if the expected size of the percolation cluster at the origin is finite.*
- (ii) *If all the border sets are finite, then the branching process has finitely many types, and the first moment matrix is of finite size. In this case p_c is the smallest value of p such that the spectral radius of the first moment matrix is 1.*
- (ii) *If in addition the pieces are finite, then the entries of the first moment matrix are algebraic functions in p . Therefore p_c is algebraic. There exists an algorithm that, given the pieces and their border sets, computes a finite extension K of the field $\mathbb{Q}(p)$ and an algebraic function f in K such that p_c is the smallest positive root of f .*

The theorem is proved in Section 4.

We list here two corollaries of Theorem 1.1. The first one is already proved in [6] by the author using different method. The second corollary answers a question of M. Sapir (personal communication) about the symmetric group $SL_2(\mathbb{Z})$, which is an simple example of a group that is not free product. Both corollaries are proved in Section 5.

By the expected sub-critical cluster size $\chi_i(p)$ we mean $E_p(|C|)$, where C is the cluster containing the origin and $p < p_c$.

COROLLARY 1.2. *Let \mathcal{G} be a free product of (transitive) graphs. Denote by $\chi_i(p)$ the expected (sub-critical) cluster size in the i -th factor graph. The critical probability p_c of \mathcal{G} is the infimum of positive solutions of*

$$\sum_{j=1}^n \prod_{i=1, i \neq j}^n \chi_i(p) - (n-1) \prod_{i=1}^n \chi_i(p) = 0.$$

Note that in the case of free products (with respect to natural set of generators) the border sets consist of one vertex and so the branching process has just one type of individual (for more details see Section 5.1). It is in fact also a special case of Theorem 1.4 below.

We explicitly compute the critical probabilities of several Cayley graphs. In particular we prove the following.

COROLLARY 1.3. *The critical probability p_c of $SL_2(\mathbb{Z})$ given by presentation $\langle a, b | a^4, b^6, a^2b^{-3} \rangle$ is an algebraic number that is equal to $.4291140496\dots$*

The general case of Cayley graphs of amalgamated products and HNN extensions is covered by the following theorem. Consider a group G acting on a simplicial tree T . The standard generating set of the group G is any generating set consisting of elements in the vertex stabilizers and free letters. It follows from the Structure theorems of Bass-Serre theory (see for example [10] or [2]) that the group G is the fundamental group of a graph of groups (see Section 6.3).

THEOREM 1.4. *Let G be a group acting on a simplicial tree and let \mathcal{G} be its Cayley graph with respect to a standard generating set. Then \mathcal{G} has a tree-like structure whose pieces correspond to the Cayley graphs of the vertex stabilizers and border sets correspond to the edge stabilizers.*

In the last section of the paper, we prove that every transitive graph with more than one end has a tree-like structure with finite border sets. This can be applied to, say, Cayley graphs of free groups with arbitrary finite generating sets. Moreover, the pieces obtained from the general construction are finite in this case, and are explicitly described in Section 7.2.

THEOREM 1.5. *Let G be a virtually free group, that is, it acts on a simplicial tree with finite vertex stabilizers. Then its Cayley graph with respect to any finite generating set has a tree-like structure with finite border sets and finite pieces. Given a finite generating set, the pieces of the tree-like structure are algorithmically constructed.*

Therefore the p_c is an algebraic number and one can use the algorithm from Theorem 1.1 (iii) to compute p_c , given any finite generating set.

This theorem is proved in Section 6.5.

This gives the first example of a class of Cayley graphs closed under quasi-isometry where we can algorithmically find the value of p_c for every graph in the class (besides the graphs with 0 and 2 ends where $p_c = 1$).

2. Graphs with tree-like structure. In what follows all graphs are supposed to be transitive, locally finite, connected and infinite (e.g. Cayley graphs of finitely generated groups). By $V(\mathcal{G})$ (resp. $E(\mathcal{G})$) we denote the set of vertices (resp. edges) of a graph \mathcal{G} .

Here we present the formal definition of the tree-like structure and several basic properties. The reader may want to review the examples presented in Section 5 to obtain more intuitive picture.

2.1. *The definition.* Now we are going to define a tree-like structure of a graph.

Note that a rooted tree of elements define a partial order on these elements. In this partial an element is smaller then other if and only if it is its descendant. The root is a maximal element in such case.

DEFINITION 2.1. *A tree-like structure on a (transitive, locally finite, connected, infinite) graph \mathcal{G} is a triple (\mathbf{P}, J, γ) , where \mathbf{P} consists of pairs of non-empty subgraphs (P_i, B_i) of \mathcal{G} , $i \in I$ (P_i are called the pieces, B_i are called the border sets), J is a finite subset of I , γ is a model map from I to J , and the following conditions are satisfied.*

- (1) *For every $i \in I$, P_i is a subgraph of \mathcal{G} and $B_i \subseteq V(P_i)$.*
- (2) *For every $i \neq j$, $E(P_i) \cap E(P_j) = \emptyset$ and $\bigcup_{i \in I} E(P_i) = E(\mathcal{G})$.*
- (3) *There is a partial order on the pieces with maximal element P_0 , such that its graphical representation is a tree with a root P_0 . Moreover, if P_i is a child of P_j , then $P_i \cap P_j = B_i$.*

Denote by $U(P_i)$ the union of the pieces in the descendant subtree of P_i (P_i including), for all $i \in I$.

- (4) *For every $i \neq j$, if $P_i \not\subseteq U(P_j)$, then $P_i \cap U(P_j) \subseteq B_j$.*
- (5) *For every $i \in I$, there is an isomorphism between $U(P_i)$ and $U(P_{\gamma(i)})$ taking pieces to pieces and border sets to border sets, respecting the order on the pieces.*

We say that there is a finite number of isomorphism classes of pairs (P_i, B_i) 's. And for $j \in J$ the pieces P_j are called the model pieces.

If there is a piece P_i with no edges, then it consists only of vertices and we can remove such a piece (and add the vertices to the parent piece) and change the tree-like structure accordingly. Therefore we will assume that each P_i contains at least one edge. Clearly the tree-like structure can be degenerated in the sense that the whole graph is just one piece or the number of pieces is finite. In what follows we will always assume that it is non-degenerated, that is, the number of pieces is infinite. In all cases considered in this paper, the pieces have at most finitely many components (this simplifies the computations).

For every $i \in I$, denote by $\Lambda_i \subset I$ the set of indices of children of P_i (given by part (3) of Definition 2.1). Now we will present several basic properties of the tree-like structure.

LEMMA 2.2. *For every i , the set $U(P_i)$ is covered by P_i and the collection of $U(P_\lambda)$ for $\lambda \in \Lambda_i$. Moreover, pairwise intersections of $U(P_\lambda)$'s for $\lambda \in \Lambda_i$ consist only of vertices of P_i .*

PROOF. By property (2) each edge is included in exactly one piece and therefore if $P_k \subset U(P_{\lambda_1})$, then $P_k \not\subset U(P_{\lambda_2})$, provided $\lambda_2 \neq \lambda_1$. Therefore by property (4) the intersection $U(P_{\lambda_1}) \cap U(P_{\lambda_2})$ is included in $B_{\lambda_1} \cap B_{\lambda_2}$. Since each $B_\lambda \subset P_i$ by property (3), the claim follows. \square

LEMMA 2.3. *Every border set (except possibly B_0 , the border set of the root) is a vertex cut set of graph \mathcal{G} . In particular, let a subgraph $U(P_i)^c$ of \mathcal{G} be induced by the edges $E(\mathcal{G}) \setminus E(U(P_i))$. Then $U(P_i) \cup U(P_i)^c = \mathcal{G}$ and $U(P_i) \cap U(P_i)^c = B_i$.*

PROOF. If the complement $U(P_i)^c$ satisfies $U(P_i) \cap U(P_i)^c = B_i$, then B_i is indeed a cut set. By property (4) for any $P_k \subset U(P_i)^c$ we have $P_k \cap U(P_i) \subset B_i$ so the claim follows. \square

2.2. *Changing the generating set.* The following lemmas apply to the Cayley graphs of groups. In that case the group itself acts on its Cayley graph (by multiplication from the right) and so it is a subgroup of the automorphism group $\text{Aut}(\mathcal{G})$. If the isomorphisms of part (5) of Definition 2.1 are (almost) in the group G , we can make the generating set smaller or slightly bigger and still obtain a tree-like structure.

LEMMA 2.4. *Let S be a finite generating set of a group G . Suppose that there is a tree-like structure of the Cayley graph \mathcal{G} of G with respect to S , and that each of the isomorphisms from part (5) of Definition 2.1 can be extended to the whole \mathcal{G} , and that these extensions form a subgroup H of $\text{Aut}(\mathcal{G})$. If $G \cap H$ has finite index in H , then for any generating subset $S' \subset S$ the Cayley graph \mathcal{G}' of G with respect to S' has a tree-like structure with pieces containing the same vertices (and fewer edges).*

PROOF. If we restrict the isomorphisms in part (5) of Definition 2.1 from H to $G \cap H$, we will increase the number of isomorphism classes by finitely many because $G \cap H$ has finite index in H . Thus we can assume that $H \subset G$.

The graph \mathcal{G}' is obtained from \mathcal{G} by removing edges with labels in $S \setminus S'$. The pieces P'_i of the tree-like structure on \mathcal{G}' are obtained in the same way

(just by removing edges with labels in $S \setminus S'$). Let us verify the conditions of Definition 2.1 Every edge belongs to exactly one piece, thus property (2) follows. Conditions (1),(3),(4) depend only on vertices of pieces. Since vertices do not change, these conditions are satisfied.

It remains to show property (5). Since the isomorphisms between $U(P_i)$'s in \mathcal{G} are given by elements in G , they are also isomorphisms of $U(P'_i)$'s in \mathcal{G}' . Thus there are finitely many isomorphism classes between pieces of \mathcal{G}' . \square

LEMMA 2.5. *Let S, S' be finite generating sets of a group G . Suppose that there is a tree-like structure with pieces P_i of the Cayley graph \mathcal{G} of G with respect to S , and that the isomorphisms from part (5) of Definition 2.1 form a subgroup H of $\text{Aut}(\mathcal{G})$. Assume that $G \cap H$ has finite index in H and that the following condition is satisfied:*

- (*) *for all $x \in S' \setminus S$ and $g \in G$ there is a piece P_i containing both vertices labeled by g and gx .*

Then the Cayley graph \mathcal{G}' of G with respect to S' has a tree-like structure with pieces P'_i , where P'_i contains the same vertices as P_i .

PROOF. Again we can assume that $H \subset G$. If we prove the claim for $S'' = S \cup S'$, then for S' it follows from Lemma 2.4. Therefore we can assume that $S \subset S'$.

The graph \mathcal{G}' is obtained from \mathcal{G} by adding edges labeled by elements in $S' \setminus S$. In the tree-like structure, we add each new edge to the oldest (i.e. the maximal in the partial order) piece P_i containing both its endpoints. Such a piece exists by assumption (*). Suppose that there are two such oldest pieces P_i and P_j containing end points of an edge e . Then none of them is a descendant of the other and their intersection is in $B_i \cup B_j$ by condition (4). Therefore e should be added to their predecessor that is older than both of them. Therefore there is a unique such oldest piece for each edge. This implies property (2). Conditions (1),(3),(4) now depend only on vertices of pieces that have not been changed. Therefore these conditions hold as in the tree-like structure on \mathcal{G} .

It remains to prove property (5). Consider an isomorphism from property (5) of the graph \mathcal{G} . Let $h \in H \subset G$ such that $U(P_i) = hU(P_j)$, and h takes pieces to pieces and border sets to border sets (we can assume $i, j \neq 0$). We will show that the descendant trees of the modified pieces P'_i and P'_j in \mathcal{G}' can be mapped by the same isomorphism h , and it takes the modified pieces to pieces and border sets to border sets as well. The vertices of pieces have not been changed so a difficulty can arise only for edges. By contradiction, assume that $U(P'_i) \neq hU(P'_j)$ respective that there is a descendant P'_k of P'_i

and P'_l of P'_j such that $P_k = hP_l$ but $P'_k \neq hP'_l$. There exists an edge e in P'_k that is missing in hP'_l (or vice versa). If $e \notin hP'_l$, then there is a older (in the ordering) piece than P'_l containing both endpoints of $h^{-1}e$. Therefore these endpoints are in the border set B'_l . Consequently the endpoints of e are in B'_k , and so the edge must be included in some ancestor of P'_k , a contradiction. \square

Lemma 2.5 can be applied only if the pieces are large enough to make the condition (*) satisfied. The following lemma allows us to enlarge pieces. The new piece P'_i is a union of the children pieces of the piece P_i . If we started with finite pieces, then the constructed pieces are again finite. Moreover, the tree of pieces (partial order) remains unchanged.

LEMMA 2.6. *Assume graph \mathcal{G} admits a tree-like structure with pieces P_i and border sets B_i . Define new pieces P'_i and border set B'_i as follows*

$$\begin{aligned} P'_0 &= P_0 \cup \bigcup_{k \in \Lambda_0} P_k, & B'_0 &= B_0, \\ P'_i &= V(P_i) \cup \bigcup_{k \in \Lambda_i} P_k, & B'_i &= V(P_i), \quad \text{for } i \neq 0. \end{aligned}$$

Define the partial order on the pieces P'_i to be the same as on the original pieces P_i . Then the modified pieces P'_i and border sets B'_i form a new tree-like structure of \mathcal{G} .

PROOF. We need to verify conditions of Definition 2.1.

- (1)-(2) Clearly follow from the definition of P'_i and B'_i .
- (3) Let P'_j be a child of P'_i . Then $P'_i \cap P'_j$ contains vertices of P_j , that is B'_j . If $k \in \Lambda_j$, then by property (4) of the original tree-like structure, $U(P_k) \cap P'_i \subset B_k \subset V(P_j) = B'_j$. Therefore $P'_i \cap P'_j = B'_j$.
- (4) Assume P'_i is not a descendant of P'_j and $i \neq j$. Then P_i is not a descendant of P_j and no child of P_i is a descendant of P_j (it can be P_j itself). Therefore using property (4) of the original tree-like structure we see that $P'_i \cap U(P_j) \subset P_j$. Since the intersection contains only vertices (by (2)), we can write

$$P'_i \cap U(P'_j) = V((P_i \cup \bigcup_{k \in \Lambda_i} P_k) \cap U(P_j)) \subset V(P_j) = B'_j.$$

- (5) Assume that there exists an isomorphism from property (5) between $U(P_i)$ and $U(P_j)$ such that the pieces and border sets are respected. The modification of the piece P_i into P'_i uses the children of P_i that

are preserved by the isomorphism. Thus the modified pieces and their border sets are also preserved by this isomorphism between $U(P'_i)$ and $U(P'_j)$. Thus there are finitely many isomorphism classes. \square

3. Partitions of the border sets. Consider a graph \mathcal{G} with a tree-like structure as above and take a realization $\omega \in \Omega$ of a percolation process on \mathcal{G} . Some pairs of vertices of the border set B_0 can be connected by open paths in \mathcal{G} . In this way the realization determines a partition of the set B_0 : two vertices are in the same class of the partition if they are connected by an open path in \mathcal{G} . The percolation process induces a probability measure on the set of all partitions of B_0 . Similarly we obtain a partition of each B_j by looking at the open paths in the subgraph $U(P_j)$. We call this a *descendant partition* of the border set B_j . We say a partition of a set B is *induced by percolation* on H , if H is a subgraph of \mathcal{G} , B is a subset of vertices of H , and two vertices of B are in the same class of the partition if and only if they are connected by an open path in H .

In this section, we will use the decomposition into pieces to find the measure on the set of descendant partitions of border sets using recurrent relations.

Let $Z^{(i)}$ be the set of all partitions of the border set B_i that can be induced by the percolation on $U(P - i)$. Let $q_i : \Omega \rightarrow Z^{(i)}$ be a map assigning to each realization $\omega \in \Omega$ the partition on the border set B_i . We consider only partitions induced by percolation on $U(P - i)$ and thus the map q_i is surjective. The measure on $Z^{(i)}$ is a pullback of the percolation measure P_p by q_i . Thus $A \subset Z^{(i)}$ is measurable if and only if $q_i^{-1}(A) \in \Sigma$. For simplicity we use the same notation P_p for the probability measure on partitions and denote the σ -algebra of measurable sets of partitions by $\mathcal{Z}^{(i)}$.

For every finite subset F of B_i and every partition $z^{(i)}$ of B_i , denote by $z^{(i)}(F)$ the set of all partitions that coincide with $z^{(i)}$ on F . These sets are obviously in $\mathcal{Z}^{(i)}$.

If the subgraphs $U(P_i)$ and $U(P_j)$ are isometric (by an isomorphism from part (5) of the definition), then the σ -algebras of partitions $\mathcal{Z}^{(i)}$ and $\mathcal{Z}^{(j)}$ are isomorphic and the measures induced by the same percolation process are preserved.

LEMMA 3.1. *A descendant partition of B_i is determined by the state of the edges in P_i and by the descendant partitions of border sets $B_\lambda, \lambda \in \Lambda_i$ (which are children of P_i).*

PROOF. Indeed using property (2) of Definition 2.1 and Lemma 2.2 we see that every edge of an open path connecting two vertices of B_i in $U(P_i)$ is in exactly one of the following graphs: P_i or $U(P_\lambda)$, $\lambda \in \Lambda_i$. We can split the path into several segments, each of them being in some $U(P_\lambda)$ or P_i . A segment in $U(P_\lambda)$ starts at some point of B_λ and ends at some other point of B_λ . Therefore the existence of the open segment is determined by the partition on B_λ . Thus in order to decide whether there is an open path connecting certain vertices in B_i it is sufficient to know the states of edges in P_i and the partitions of B_λ , $\lambda \in \Lambda_i$. \square

For every measurable set $A \in \mathcal{Z}^{(i)}$ and every vector $\zeta \in \prod_{\lambda \in \Lambda_i} Z^{(\lambda)}$ of partitions on B_λ 's, $\lambda \in \Lambda_i$, we denote the conditional probability of A provided ζ by $P_p(A|\zeta)$. Denote by $\mu_p^{(i)}$ the product measure on $\prod_{\lambda \in \Lambda_i} Z^{(\lambda)}$ such that the measure on each $Z^{(\lambda)}$ is given by P_p .

Then we can write:

$$(1) \quad P_p(A) = \int_{\zeta \in \prod_{\lambda \in \Lambda_i} Z^{(\lambda)}} P_p(A|\zeta) d\mu_p^{(i)}(\zeta).$$

Note that since for pieces of the same isomorphism class we have isomorphic σ -algebras of the set of partitions and measures on them, we can consider only one such equation for each model border set (of index in the finite set J from the definition).

This defines an operator on the space of all measures on the direct product of $Z^{(i)}$'s. More precisely, let X_j be the space of the probability measures on $Z^{(j)}$ with the σ -algebra $\mathcal{Z}^{(j)}$. In fact $X_j \subset [0, 1]^{\mathcal{Z}^{(j)}}$. Denote $\prod_{j \in J} X_j$ by \mathbf{X} .

For $x = (x_1, \dots, x_{|J|}) \in \mathbf{X}$ denote by $\mu_x^{(j)}$ the product measure on the space $\prod_{\lambda \in \Lambda_j} Z^{(\lambda)}$ such that the measure on each $Z^{(\lambda)}$ is given by $x_{\gamma(\lambda)} \in X_{\gamma(\lambda)}$.

Define an operator $\Psi_p : \mathbf{X} \rightarrow \mathbf{X}$ (where $\Psi_p(x) = (\Psi_p(x)_1, \dots, \Psi_p(x)_{|J|})$) by

$$(2) \quad \Psi_p(x)_j(A) = \int_{\zeta \in \prod_{\lambda \in \Lambda_j} Z^{(\lambda)}} P_p(A|\zeta) d\mu_x^{(j)}(\zeta), \text{ for all } A \in \mathcal{Z}^{(j)}.$$

For any $p \in (0, 1)$ the measure on $Z^{(j)}$, $j \in J$ induced by the percolation on $U(P_j)$ is a fixed point of this operator Ψ_p , indeed compare (2) with (1).

If the pieces are finite, then the equations (1) form a finite system of polynomial equations in the unknown variables $P_p(z^{(j)})$. The probabilities of partitions are important for evaluating the first moment matrix of the branching process defined in Section 4.5.

We can endow each $Z^{(j)}$ with a topology generated by the cylindrical sets $z^{(j)}(F)$. This topological space is second countable and Hausdorff. Moreover, the space is compact since each sequence has an accumulation point. (Indeed let $F_i, i = 1, \dots$ be an increasing sequence of finite subsets of B_j , and $\bigcup F_i = B_j$. In order to find an accumulation point of a sequence x_k , we can consider a set of partitions that agree on F_i with infinitely many x_k 's, and let i go to infinity.)

The space \mathbf{X} of the probability measures on a compact metric space is a convex compact metric space in the weak* topology induced by continuous functions (follows from the Riesz representation theorem). Then a continuous operator on \mathbf{X} has a fixed point. The set of its fixed points is closed and since the operator Ψ_p is linear, the set is also convex. The operator Ψ_p acts linearly on the space of measures and has norm at most one because it preserves the subset of probability measures. Therefore the set of its fixed points is convex, compact.

4. Branching processes.

4.1. *Preliminaries.* Recall that a multi-type branching process is a Markov process that models a population in which each individual in generation n produces some random number of offspring of the various types in generation $n + 1$, according to a fixed probability distribution that depends only on the type of the individual.

Assume S is a set of types of individuals and that a type s individual produces children of different types according to a probability distribution p_s on \mathbb{N}_0^S , where $\mathbb{N}_0 = \{0, 1, 2, \dots\}$. Assume that all individuals produce offspring independently of each other and of the history of the process. Let vector $X_n \in \mathbb{N}_0^S$ represent the n -th generation, where each coordinate $X_n[s]$ represents the number of individuals of type s in the n -th generation. It is given by the recurrent relation

$$(3) \quad X_{n+1} = \sum_{s \in S} \sum_{i=1}^{X_n[s]} \xi_{ns}^i,$$

where ξ_{ns}^i are independent random variables with the distribution p_s . The sequence $\{X_n\}_0^\infty$ is called a *multi-type Galton-Watson branching process* with initial population size $X_0 \in \mathbb{N}_0^S$ and offspring distribution $p_s, s \in S$.

Let $e_s \in \mathbb{N}_0^S$ be a vector with 1 at the position $s \in S$ and zeros at other positions. Let $M_n(e_s, B)$ be the expected number of individuals of the n -th generation of types in $B \subset S$. For any fixed initial e_s , $M_n(e_s, \cdot)$ is a measure

on S given by

$$(4) \quad M_n(e_s, B) = E \left(\sum_{s \in B} X_n[s] | X_0 = e_s \right).$$

The initial population can be given by any vector a in \mathbb{N}_0^S (usually it is some e_s , an atomic measure). We will give some properties of the M_n 's using a notation from the book of Nummelin [9]. Assume the expected size of the first generation is bounded, we can condition on the states in the first generation and obtain:

$$(5) \quad M_n(a, B) = \int_{s \in S} M_{n-1}(s, B) dM_1(a, s).$$

Each $M_n : (S, P(S)) \rightarrow \mathbb{R}_+$ is a kernel acting as an operator \tilde{M}_n on the space of measures on S by

$$\tilde{M}_n(\mu)(B) = \int_{s \in S} M_n(s, B) d\mu(s).$$

The product of two kernels is defined as in expression (5), thus \tilde{M}_n is the n -th iterate kernel of \tilde{M}_1 .

A branching process is called *singular* if each individual has exactly one offspring almost surely. M is *irreducible* if and only if there exist a σ -finite measure ϕ on S such that for all ϕ -positive sets $B \subset S$ and $s \in S$, $M_n(e_s, B) > 0$ for some n .

If a non-singular branching process is irreducible, then the population becomes extinct or explodes exponentially. The irreducibility condition is very important here. The branching process we will construct can be reducible in general. This prevents us from making claims about the extinction of the process in general. If we restrict ourself to the finite set of types, we can make further claims, see Section 4.5.

If S is finite, then the operators M_n are matrices. The expected size of the first generation (the first moment matrix M) is given by

$$(6) \quad m_{rs} = E(X_1[s] | X_0 = e_r).$$

Two types r and s are said to be in the same class if an individual of type r is in the offspring of an individual of type s with positive probability and vice versa (i.e. for some n , the (r, s) entry of M^n , $m_{rs}^{(n)}$, is positive). Now the multi-type branching process is irreducible if all types are in the same class.

The process is *positively regular* if there exists n such that all elements of M^n are strictly positive. If the process is irreducible but non-positively

regular, then it is *periodic*. The period of a branching process is a number d such that the matrix M may be represented, after reordering the types of individuals, in the form

$$M = \begin{bmatrix} 0 & M(1,2) & 0 & \dots & 0 \\ 0 & 0 & M(2,3) & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & M(d-1,d) \\ M(d,1) & 0 & 0 & \dots & 0 \end{bmatrix},$$

where $M(i, i+1)$ denotes a non-zero matrix.

We will need the following result from the theory of branching processes.

LEMMA 4.1. *If an irreducible multi-type Galton-Watson process with finite number of types is non-singular, then the population becomes extinct with probability one, whenever the maximal eigenvalue of the first moment matrix M is at most one.*

For the proof see Mode [8], Theorem 7.1 on page 16 and Theorem 2.1 on page 54. Note that by The Perron-Frobenius theorem the maximal (in absolute value) eigenvalue of M will always be a non-negative real number.

4.2. *Coloring of the tree of pieces according to the percolation.* Next we will relate the percolation cluster size to the population of a multi-type branching process. The number of types will be as small as possible in order to simplify the computation at the cost that the population size will not match, only approximate the cluster size, in particular it will be finite if and only if the percolation cluster was also finite.

Assume that there is a distinct vertex o called the origin in the border set B_0 . Each realization of a percolation gives rise to a coloring of the tree of pieces in the following way.

Consider a piece P_i that is not the root. Consider a subgraph $U(P_i)^c$ of \mathcal{G} induced by the edges $E(\mathcal{G}) \setminus E(U(P_i))$. Note that the intersection of $U(P_i)^c$ and $U(P_i)$ is the border set B_i , and it is a cut set by Lemma 2.3. The subgraph $U(P_i)^c$ contains the origin. Vertices of the border set B_i may be connected by open paths in $U(P_i)^c$. This gives a new partition on B_i and one class of the partition may be connected to the origin. The color of the piece is the data consisting of the partition and the distinct origin-connected class (which may be empty). Denote all possible colors $y^{(i)}$ by $Y^{(i)}$, that is all colors that appear for some realization of percolation $\omega \in \Omega$. Say that a piece is white if it is of a color such that the class connected to the origin is

empty. Observe that in this model the colors of pieces depend not only on the parent but on the colors of its siblings and the whole subtree of their descendants. This does not give us a branching process directly, but this last difficulty is to be overcome.

Consider the maps $q'_i : \Omega \rightarrow Y^{(i)}$ assigning to each realization a color of the piece P_i . We can pullback the percolation probability measure P_p to $Y^{(i)}$. This determines a σ -algebra $\mathcal{Y}^{(i)}$ on $Y^{(i)}$. We can identify colors of the pieces P_i and $P_{\gamma(i)}$ (γ is the model map from part (5) of Definition 2.1) because the spaces $Y^{(i)}$ and $Y^{(\gamma(i))}$ are isomorphic.

LEMMA 4.2. *For every $i \in I$ and $v \in \Lambda_i$, $U(P_v)^c$ is covered by P_i , $U(P_i)^c$ and the collection of $U(P_\lambda)$ for $v \neq \lambda \in \Lambda_i$. So the color of P_v is determined by the state of the edges in P_i , the color of P_i and descendant partitions of border sets B_λ for $v \neq \lambda \in \Lambda_i$.*

PROOF. The argument is identical to the one used in Lemma 2.2, and Lemma 3.1. \square

Note that only a finite number of children will be non-white if $p < \min\{p_c(P_i)\}$, and all children of a white piece will be white as well.

4.3. The complete branching process. In the previous sections we assigned to each piece P_i (or its border set B_i) a descending partition and a color based on the realization of the percolation process, that is a pair $(z^{(i)}, y^{(i)}) \in \bigcup_{j \in J} (Z^{(j)} \times Y^{(j)})$. Denote $\bigcup_{j \in J} (Z^{(j)} \times Y^{(j)})$ by Θ .

Assume a piece P_j with border set B_j has a descendant partition $z^{(j)}$ and a color $y^{(j)}$. The descendant partitions and the colors of the pieces $P_\lambda, \lambda \in \Lambda_j$ (the offspring pieces of P_j) are random variables taking values in $Z^{(\lambda)} \times Y^{(\lambda)}$. Denote by (ζ, η) the vector of these random variables, with values in $\prod_{\lambda \in \Lambda_j} Z^{(\lambda)} \times \prod_{\lambda \in \Lambda_j} Y^{(\lambda)}$. Let Q be a random variable representing all the colors and descending partitions of pieces in $U(P_j)^c$.

LEMMA 4.3. *We claim that the distribution of the descendant partitions and colors of the offspring pieces depends only on the parent, in particular*

$$(7) \quad P_p((\zeta, \eta) \in A | (z^{(j)}, y^{(j)}) \& Q) = P_p((\zeta, \eta) \in A | (z^{(j)}, y^{(j)})),$$

for any measurable set $A \subset \prod_{\lambda \in \Lambda_j} Z^{(\lambda)} \times \prod_{\lambda \in \Lambda_j} Y^{(\lambda)}$.

PROOF. From Lemma 4.2 and Lemma 3.1 we have that (ζ, η) is independent of Q for a given $(z^{(j)}, y^{(j)})$. Therefore the claim follows.

In other words, the descendant partition and the color of the parent encodes every connectedness relation coming from $U(P_j)^c$. So given the descendant partition and the color of the piece P_i , the descendant partitions and the colors of pieces in $U(P_j)^c$ and of pieces in $U(P_j)$ are independent. \square

The purpose of this claim becomes clear after the following definition.

Let us define a measure $\mathcal{D}(z^{(i)}, y^{(i)})$ on $\prod_{\lambda \in \Lambda_i} Z^{(\lambda)} \times \prod_{\lambda \in \Lambda_i} Y^{(\lambda)}$ according to the probabilities in Lemma 4.3. We identify the space $Z^i \times Y_i$ with $Z^{\gamma^{(i)}} \times Y^{\gamma^{(i)}}$ and we count the number of repetitions of each $(z^{(j)}, y^{(j)})$, $j \in J$ among the offspring. In this way we define a map $\prod_{\lambda \in \Lambda_j} Z^{(\lambda)} \times \prod_{\lambda \in \Lambda_i} Y^{(\lambda)} \rightarrow \mathbb{N}_0^\Theta$. For any $i \in J$, we can pull back the distribution $\mathcal{D}(z^{(i)}, y^{(i)})$ to a distribution $\mathcal{D}'(z^{(i)}, y^{(i)})$ on \mathbb{N}_0^Θ .

DEFINITION 4.4. *The complete multi-type Galton–Watson branching process induced by the percolation with parameter p on a graph with a tree-like structure is given by the following conditions.*

The type of an individual is given by the color and the descendant partition. The set of types is $\Theta = \bigcup_{j \in J} (Z^{(j)} \times Y^{(j)})$.

There is one initial individual of type (z_0, y_0) , where z_0 is the diagonal descendant partition of B_0 (i.e. all classes are of size 1) and y_0 is the color with the diagonal partition of B_0 with only the origin in the distinct origin-connected class.

Every individual of some type $(z^{(j)}, y^{(j)})$ gives birth to $|\Lambda_j|$ individuals, with the distribution of types $\mathcal{D}'(z^{(j)}, y^{(j)})$.

We can represent this branching process by a tree. Clearly it will match the tree of pieces of the graph.

LEMMA 4.5. *The coloring of the tree of pieces according to the percolation has the same distribution as the above defined complete branching process.*

PROOF. Let us represent the coloring of the tree of pieces by a random process (X_n) , where $X_n \in \mathbb{N}_0^\Theta$. The colors and the descendant partitions are represented by elements of Θ and each coordinate of X_n gives a number of pieces with a specific color and a specific descending partition in the n -th generation of the tree of pieces.

By Lemma 4.3 the distribution of the offspring of an individual in the n -th generation is independent of the other individuals (in generation at most n) and depends only on its type $(z^{(j)}, y^{(j)})$. The distribution is $\mathcal{D}'(z^{(j)}, y^{(j)})$.

Therefore the process X_n coincides with the complete branching process above. \square

4.4. *The branching process with e reduced number of types.* Next we will reduce the number of types and obtain a different branching process. We will also impose an independency condition on the offspring of every individual, that is, the joint distribution of the offspring $\mathcal{D}(y^{(j)})$ will be a product measure. Nevertheless we will show that the expected population size of this reduced branching process contains enough information about p_c .

Using Lemma 4.2 we can make the following observation about the coloring of the tree of pieces. We look at the piece P_i and evaluate the conditional probability of a specific piece P_v having color $y^{(v)}$ (resp. $y^{(\gamma(v))}$) assuming the descendant partitions of the other border sets are given by the vector $\zeta \in \prod_{\lambda \in \Lambda_i} Z^{(\lambda)}$ (note that the descendant partition on P_v has no influence). Taking an expected value of these conditional probabilities over all possible descendant partitions gives us the probability $P_p(y^{(v)}|y^{(i)})$ of having a child piece of a given color $y^{(v)}$ from a piece of color $y^{(i)}$. These probabilities are the same for the piece P_i as for its model piece $P_{\gamma(i)}$, so we can write for $j \in J$

$$(8) \quad P_p(y^{(\gamma(v))} \in B|y^{(j)}) = \int_{\zeta \in \prod_{\lambda \in \Lambda_j} Z^{(\lambda)}} P_p(y^{(\gamma(v))} \in B|y^{(j)} \& \zeta) d\mu^{(j)}(\zeta),$$

where $v \in \Lambda_j$ and $B \in \mathcal{Y}^{(\gamma(v))}$, a measurable set of colors. These conditional probabilities may differ for different children v_1 and $v_2 \in \Lambda_j$ even if $\gamma(v_1) = \gamma(v_2)$ (that is if the set of possible colors coincide).

Denote by $Y'^{(j)}$ the subset of $Y^{(j)}$ such that $y^{(j)} \in Y'^{(j)}$ if and only if it is not white. Let $\mathbf{Y} = \bigcup_{j \in J} Y'^{(j)}$. Then \mathbf{Y} is the set of all non-white colors. Let the σ -algebra \mathcal{Y} on \mathbf{Y} be generated by the intersections of sets in $\mathcal{Y}^{(j)}$ with $Y'^{(j)}$.

DEFINITION 4.6. *The reduced multi-type Galton-Watson branching process induced by the percolation with parameter p on a graph with a tree-like structure is given by the following conditions.*

The types of individuals are colors in \mathbf{Y} .

There is one initial individual with color $y_0^{(0)}$ that is the diagonal partition of B_0 with only the origin in the distinct origin-connected class.

Every individual of some color $y^{(j)}$ gives birth to possibly $|\Lambda_j|$ individuals, each of which is born and has its color assigned independently of the others. The distribution of the color $y^{(\gamma(\lambda))}$ of a child indexed by $\lambda \in \Lambda_j$ follows the law in formula (8).

Note that white is no longer a legitimate color. A child indexed by $\lambda \in \Lambda_j$ is not born in the reduced branching process with the same probability as the probability, that the corresponding piece is white in the percolation, that is $\mathbb{P}_p(y^{(\gamma(v))} \text{ is white} | y^{(j)})$. This is not the only difference between the reduced branching process and the coloring of the tree of pieces according to the percolation. The joint distribution of the offspring of an individual is different (because of the independence), and only the first moment is the same. But the first moment is all we need.

REMARK 4.7. *In the definition of the tree-like structure we allowed some isomorphism classes of pieces to be finite. In particular very often the root is not isometric to any other piece. In such case, it is useful to start the branching process not at the root (with one element), but with a generation that already consists only of pieces that are in infinite isomorphism classes. This gives us the initial measure on the colors. Therefore we consider only a subset J' of J such that $j \in J'$ if and only if $|\gamma^{-1}(j)| = \infty$. Then $\mathbf{Y} = \bigcup_{j \in J'} Y^{(j)}$.*

THEOREM (1.1 (i)). *For a percolation with parameter p the reduced branching process on the tree of pieces has the property that the expected size of its population is finite if and only if the expected size of the percolation cluster at the origin is finite.*

PROOF. Note that, by the result of Aizenman and Barsky [1], the subcritical phase on transitive graphs is equivalently characterized by the finiteness of the expected cluster size (they proved this for \mathbb{Z}^d , the generalization to all transitive graphs was pointed out by Lyons and Peres [7]).

Clearly $p_c(\mathcal{G}) \leq p_c(P_i)$. If there is an infinite cluster at the origin for p , such that $p < p_c(P_i)$, then the cluster has to intersect infinitely many border sets almost surely. Therefore the number of border sets connected to the origin is finite if and only if the original cluster was also finite. In what follows we will always assume $p \leq p_c(P_i)$ (if P_i is finite, then we set $p_c(P_i) = 1$).

Given a realization of percolation we have introduced a coloring of the tree of pieces. The percolation cluster at the origin is infinite if and only if the non-white colored component of the tree of pieces is infinite (by the assumption that p is smaller than p_c of the pieces). The expected size of the colored component is a sum of the probabilities that a piece is non-white over all pieces of the tree.

The branching process from Definition 4.6 can be naturally illustrated by a tree isomorphic to the tree of pieces. The distribution of colors of a

specific individual in this branching process equals to the distribution of non-white colors of the related piece in the percolation. Therefore the expected population size of this branching process equals the expected number of non-white colored pieces in the percolation.

Therefore $p < p_c$ if and only if the related branching process has finite expected population size. \square

The branching process is not singular because for $p < 1$ (and some non-white color y) there is no color, which appears as an only child of y almost surely (i.e. with positive probability there are less or more children with different colors). Assume the initial measure is $\nu : \mathcal{Y} \rightarrow \mathbb{R}_+$ (it can be the atomic measure from Definition 4.6 or distribution of the first generation of pieces with infinite isomorphism classes from Remark 4.7). Then the expected population size of the branching process is $\sum_{n=0}^{\infty} \tilde{M}_n \nu(\mathbf{Y})$. The critical probability p_c is then a supremum of all p such that $\sum_{n=0}^{\infty} \tilde{M}_n \nu(\mathbf{Y}) < \infty$. In general, the convergence of $\sum_{n=0}^{\infty} \tilde{M}_n \nu(\mathbf{Y})$ may depend strongly on the initial measure ν . But we will show that if the border sets are finite, then it actually depends only on the spectral radius of M_1 .

4.5. *The case of finite border sets.* If the border sets are finite, then the space of partitions is finite and so is the space of colors \mathbf{Y} . Denote by $M = [m_{ab}]_{a,b \in \mathbf{Y}}$ the first moment matrix of the branching process, that is the matrix of expected number of offspring of each color $m_{ab} = \mathbb{E}_p(\#b|a)$, where

$$\begin{aligned} \mathbb{P}_p(y^{(\gamma(v))} | y^{(j)}) &= \int_{\zeta \in \prod_{\Lambda_j} Z^{(\lambda)}} \mathbb{P}_p(y^{(\gamma(v))} | y^{(j)} \& \zeta) d\mu^{(j)}(\zeta) \\ (9) \quad \mathbb{E}_p(\#y^{(k)} | y^{(j)}) &= \sum_{\lambda \in \Lambda_j, \gamma(\lambda)=k} \mathbb{P}_p(y^{(\gamma(\lambda))} | y^{(j)}). \end{aligned}$$

The expected number of individuals of the n -th generation is then given by the n -th power of M .

THEOREM (1.1 (ii)). *If all the border sets are finite, then the branching process has finitely many types, and the first moment matrix is of finite size. In this case p_c is the smallest value of p such that the spectral radius of the first moment matrix is 1.*

PROOF. Assume the graph \mathcal{G} has a tree-like structure such that the border sets B_i 's are finite. Using the already proved part (i) of Theorem 1.1 we

need to decide for which p the expected population size of the constructed branching process is finite.

This branching process is non-singular for $p < 1$ because if a piece has more than one child, then the offspring size is bigger than one with positive probability, and if every piece has exactly one child, then it has no offspring (only white) with positive probability. The expected population size is $\sum M^n$ applied to the initial measure. If the spectral radius is less than 1, then the sum $\sum M^n$ is always finite. If the spectral radius is at least 1, then there exists a possible initial measure, for which the expected population size is infinite (follows from The Perron-Frobenius theorem). If the process is irreducible, then it is independent of the choice of initial measure.

In case M is irreducible the result follows directly from Lemma 4.1. In the other case there are several classes of types and we denote by M_1, \dots, M_k the first moment matrices of each class. We can reorder the types in \mathbf{Y} such that M has diagonal blocks equal to M_i 's and all entries above these diagonal blocks are zero. Thus the spectral radius of M is the spectral radius of some M_s . There is a type (color from \mathbf{Y}) in the s -th class such that with positive probability there is a piece P_i with this color (because we considered only those colors that are realized by percolation). The process starting at this P_i will have infinite cluster size whenever the spectral radius of M_s is at least one. Therefore the expected size of the percolation cluster is finite whenever the maximal eigenvalue of M is less than one.

Note that $\det(M - 1) = 0$ if 1 is an eigenvalue of M and for $p = 0$ all eigenvalues of M are zero. The eigenvalues depend continuously on the matrix entries, which are continuous functions of p . Therefore p_c is the first positive value of p such that $\det(M - 1) = 0$. \square

THEOREM (1.1 (iii)). *If all the pieces are finite, then the entries of the first moment matrix of the reduced branching process are algebraic functions in p . Therefore p_c is algebraic.*

There exists an algorithm that, given the model pieces and their border sets, computes a finite extension K of the field $\mathbb{Q}(p)$ and a function f in K such that p_c is the smallest positive root of f .

PROOF. If the pieces are finite, then the probabilities of the descending partitions, $P_p(z^{(j)})$, can be found as solutions of the system of equations (1) introduced in the previous section. Let us index the partitions on B_j by natural numbers $1, \dots, |Z^{(j)}|$. Denote by $x_{j,i}$ the probability of the i -th possible partition on the border set of model piece P_j , that is $x_{j,i} = P_p(z_i^{(j)})$. Then we have the following system of equations in the unknown variables $x_{j,i}$.

$$(10) \quad x_{j,i} = \sum_k \left(\sum_{\Gamma \in L} \alpha(\Gamma) \right) \prod_{\lambda \in \Lambda_j} x_{\gamma(\lambda), k(\lambda)},$$

where

- k is a map that assigns partitions to the border sets B_λ 's. That is $k : \Lambda_j \rightarrow \mathbb{N}$, and for all $\lambda \in \Lambda_j$, $k(\lambda) \in \{1, 2, \dots, |Z^{(\lambda)}|\}$. We sum over all such possible functions k .
- For a given k , L is the set of subgraphs of P_j such that $\Gamma \in L$ if and only if the following holds: provided the partitions on the children pieces are $z_{k(\lambda)}^{(\lambda)}$ and the open edges of P_j are given by Γ , the partition on B_j (induced by $U(P_j)$) equals $z_i^{(j)}$. Note that by Lemma 3.1 the partition on B_j is uniquely determined by the provided informations.
- $\alpha(\Gamma)$ is the probability of Γ in the percolation, that is $\alpha(\Gamma) = p^{E(\Gamma)}(1-p)^{E(P_i)-E(\Gamma)}$.

The number of equations is equal to the number of variables. The degree in p of each equation (10) is equal to the number of edges in a piece and the degree in the unknown variables $x_{j,i}$'s is $|\Lambda_j|$, the number of offspring of the piece P_j . Denote by K the algebraic extension of $\mathbb{Q}(p)$ containing roots of this system of equations.

Similarly as above we can rewrite the formulas (9) as follows

$$(11) \quad P(y^{(\gamma(v))} | y^{(j)}) = \sum_k \left(\sum_{\Gamma \in L'} \alpha(\Gamma) \right) \prod_{\lambda \in \Lambda_j} x_{\gamma(\lambda), k(\lambda)},$$

where

- k and $\alpha(\Gamma)$ are defined as above, and
- L' depends on k and it is the set of subgraphs of P_j such that $\Gamma \in L'$ if and only if the color on B_v is $y^{(\gamma(v))}$ provided the color of the parent is $y^{(j)}$, the descendant partitions on the children pieces are $z_{k(\lambda)}^{(\lambda)}$ and the open edges of P_j are given by Γ (by Lemma 4.2 these uniquely determine the color on B_v).

Consequently the entries of the first moment matrix M , given by formula (9), are polynomial functions in p and $x_{j,i}$'s. The critical probability p_c is the first positive value of p such that $\det(M - 1) = 0$. The function $\det(M - 1)$ is an element of the field K , and p_c is an algebraic number in this case. \square

5. Examples.

5.1. *Free products of (transitive) graphs.* The free product \mathcal{G} of transitive graphs G_1, \dots, G_n is an infinite connected graph constructed as a union of copies of G_i 's such that each vertex of \mathcal{G} belongs to exactly one copy of each G_i , and every simple closed path in \mathcal{G} is included in one copy of some G_i . In particular, if $G_1 = \langle S_1 \rangle$ and $G_2 = \langle S_2 \rangle$, a Cayley graph of $G_1 * G_2$ with respect to $S_1 \cup S_2$ is a free product of Cayley graphs of G_1 and G_2 with respect to S_1 and S_2 respectively.

We present here a different method to obtain a result shown previously by the author in [6].

COROLLARY (1.2). *Let \mathcal{G} be a free product of transitive graphs $\mathcal{G}_1, \dots, \mathcal{G}_n$. Denote by $\chi_i(p)$ the expected sub-critical cluster size in the i -th factor graph G_i . The critical probability p_c of \mathcal{G} is the infimum of positive solutions of $\sum_{j=1}^n \prod_{i=1, i \neq j}^n \chi_i(p) - (n-1) \prod_{i=1}^n \chi_i(p) = 0$.*

PROOF. The tree-like structure is very natural in this case. The copies of factor graphs $\mathcal{G}_1, \dots, \mathcal{G}_n$ are pieces (thus $J = \{1, \dots, n\}$) and each vertex is in fact a border set.

The partitions of border sets are trivial (the spaces of partitions have size 1) in this case. For the coloring we need only two colors $Y^{(i)} = \{y_1^{(i)}, y_2^{(i)}\}$. Assuming that $y_2^{(i)}$ is white, then $\mathbf{Y} = \{y_1^{(1)}, \dots, y_1^{(n)}\}$.

Denote by b_i the border vertex of piece P_i and by $\tau_p^{(i)}(a \leftrightarrow b)$ the probability that a and b are connected in \mathcal{G}_i . Denote by $\chi_i(p)$ the expected cluster size in \mathcal{G}_i . Now we are ready to compute the matrix $M = (m_{ij})$.

$$\begin{aligned}
 \mathbb{P}_p(y_1^{(\gamma(\lambda))} | y_1^{(j)}) &= \tau_p^{(j)}(b_j \leftrightarrow b_\lambda) \\
 \mathbb{E}_p(\#y_1^{(k)} | y_1^{(j)}) &= \chi_j(p) - 1 \\
 (12) \quad m_{ij} &= \chi_i(p) - 1, \text{ for } i \neq j \\
 m_{ii} &= 0
 \end{aligned}$$

In order to find for which p the spectral radius of M equals 1 we solve the equation $\det(M - 1) = 0$ by Theorem 1.1 (iii). The determinant is computed as follows (we subtract the first column from all the others and then expand the determinant along the first row):

$$\det(M - 1) = \det \begin{bmatrix} -1 & \chi_1(p) - 1 & \dots & \chi_1(p) - 1 \\ \chi_2(p) - 1 & -1 & \dots & \chi_2(p) - 1 \\ \vdots & \vdots & \ddots & \vdots \\ \chi_n(p) - 1 & \chi_n(p) - 1 & \dots & -1 \end{bmatrix}$$

$$\begin{aligned}
\det(M - 1) &= \det \begin{bmatrix} -1 & \chi_1(p) & \chi_1(p) & \cdots & \chi_1(p) \\ \chi_2(p) - 1 & -\chi_2(p) & 0 & \cdots & 0 \\ \chi_3(p) - 1 & 0 & -\chi_3(p) & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \chi_n(p) - 1 & 0 & 0 & \cdots & -\chi_n(p) \end{bmatrix} \\
&= - \prod_{i=2}^n (-\chi_i(p)) + \sum_{j=2}^n (\chi_j(p) - 1) \prod_{i=1, i \neq j}^n (-\chi_i(p)) \\
(13) \quad &= (-1)^n \left(\sum_{j=1}^n \prod_{i=1, i \neq j}^n \chi_i(p) - (n-1) \prod_{i=1}^n \chi_i(p) \right)
\end{aligned}$$

Therefore p_c is the infimum of positive solutions of

$$\sum_{j=1}^n \prod_{i=1, i \neq j}^n \chi_i(p) - (n-1) \prod_{i=1}^n \chi_i(p) = 0.$$

□


5.2. $SL(2, \mathbb{Z})$. The easiest example of a Cayley graph with a tree-like structure that is not a free product, is $SL(2, \mathbb{Z})$. It is the amalgamated product $\mathbb{Z}_4 *_{\mathbb{Z}_2} \mathbb{Z}_6$ with standard generating set $\{a, b\}$ so that $\mathbb{Z}_4 = \langle a \rangle$ and $\mathbb{Z}_6 = \langle b \rangle$. We can illustrate the general method here because the required computations (for obtaining p_c) are relatively simple, or at least doable.

COROLLARY (1.3). *The critical probability p_c of $SL_2(\mathbb{Z})$ given by presentation $\langle a, b | a^4, b^6, a^2b^{-3} \rangle$ is an algebraic number equal to .4291140496...*

PROOF. The Cayley graph has the tree-like structure with pieces of two isomorphism classes (squares P_1 and hexagons P_2) corresponding to the factor groups, see Figure 1. The hexagonal pieces consist of six-tuples of vertices connected by bold lines in Figure 1 and squares contain dashed edges. Each vertex is contained in exactly one square and one hexagon and each neighboring square and hexagon share two vertices - the border sets.

First we need to find the distribution of the descendant partitions of the border sets. There are two possible descendant partitions on each border set, B_1 and B_2 . Denote them by $Z^{(1)} = \{z_1^{(1)}, z_2^{(1)}\}$ and $Z^{(2)} = \{z_1^{(2)}, z_2^{(2)}\}$, where $z_1^{(i)}$ means that the two vertices of the border set B_i are connected and $z_2^{(i)}$ means that they are not connected. Now we can find the probability of these partitions using the formula (1), which leads to a quadratic equation.

In Section 4 we introduced the coloring of a tree. For a border set of size two there are three non-white colors: either both vertices are connected

FIG 1. The Cayley graph of $SL(2, \mathbb{Z})$ with one border set circled

to the origin or exactly one of them is connected to the origin. The two different situations when exactly one vertex of the border set is connected to the origin are symmetric, i.e. there is an obvious isomorphism of $U(P_i)$ to itself such that each vertex of B_i is mapped onto the other one. Therefore these two colors have the same distribution of offspring. This simplification is special for the Cayley graph of $SL(2, \mathbb{Z})$, because this new symmetry is not one of the isomorphisms in condition (5) of the definition of the tree-like structure. Since the distribution of offspring does not depend on which of the two vertices is connected to the origin, we identify these two colors as one.

The set of colors becomes $Y^{(i)} = \{y_r^{(i)}, y_b^{(i)}, y_w^{(i)}\}$ where r (red) means that both vertices of the border set are connected to the origin, b (blue) means that exactly one vertex is connected to the origin (thus the border set is disconnected) and w stands for white - no vertex connected to the origin. We evaluate the probabilities of getting a child of a given color, according to formula (11) and the first moment matrix M takes the following form.

$$(14) \quad M = \begin{bmatrix} 0 & 0 & P_p(y_r^{(\lambda)} | y_r^{(1)}) & P_p(y_b^{(\lambda)} | y_r^{(1)}) \\ 0 & 0 & P_p(y_r^{(\lambda)} | y_b^{(1)}) & P_p(y_b^{(\lambda)} | y_b^{(1)}) \\ 2P_p(y_r^{(\lambda)} | y_r^{(2)}) & 2P_p(y_b^{(\lambda)} | y_r^{(2)}) & 0 & 0 \\ 2P_p(y_r^{(\lambda)} | y_b^{(2)}) & 2P_p(y_b^{(\lambda)} | y_b^{(2)}) & 0 & 0 \end{bmatrix}.$$

If we solve the equation $\det(M - 1) = 0$ we obtain $p_c = .4291140496 \dots$

□

5.3. *Grandparent tree.* An interesting example of a transitive graph with infinitely many ends that is not a Cayley graph is the grandparent tree. It is obtained from the three-regular tree, by adding edges connecting each vertex to its grandparent, that is, we pick one end and at each vertex we add an edge connecting it to a vertex at distance 2 in the direction of the distinguished end.

In the suggested decomposition below, we obtain pieces of size four with border sets of size two. Note that in the Figure 2 the graph is oriented so that the distinguished end is to the top. In the right picture we see that the border set of a piece consist of the two upper vertices and each of its two children intersects it by a pair of vertices - the middle one and the left (resp. right) one.




FIG 2. Part of the grandparent tree and its piece with cycled border sets.

Note that the pieces that lie on the path from the root to the distinguished end have different orientation - the border set is a different pair of vertices inside the piece (exercise to the reader).

A careful analysis of the branching process can be carried out as in the case of $SL(2, \mathbb{Z})$. The p_c obtained is $.158656326\dots$

6. Fundamental groups of graphs of groups. In this section we will generalize the example of $SL(2, \mathbb{Z})$ to arbitrary graphs of groups by showing that the Cayley graphs of fundamental group of graphs of groups have the tree-like structure. To make the transition simpler, we first consider amalgamated products and HNN extensions before proceeding to arbitrary graphs of groups.

6.1. *Amalgamated products with standard generating sets.* Recall that an amalgamated product $G_1 *_H G_2$ is obtained from two groups G_1 and G_2

provided monomorphisms $i_k : H \rightarrow G_k$, $k = 1, 2$. The group $G_1 *_H G_2$ is a quotient of the free product $G_1 * G_2 / N$, where N is the smallest normal subgroup containing elements $i_1(h)i_2^{-1}(h)$, $h \in H$.

Consider the right Cayley graph \mathcal{G} of an amalgamated product $G_1 *_H G_2$ with respect to the generating set $S_1 \cup S_2$, where $G_1 = \langle S_1 \rangle$ and $G_2 = \langle S_2 \rangle$. The structure of the Cayley graph \mathcal{G} is the following. First consider the copies of the Cayley graph \mathcal{G}_1 of G_1 and the Cayley graph \mathcal{G}_2 of G_2 containing the origin. These subgraphs intersect by the vertices of H , which we consider as a border set. Each coset of H inside G_1 is another border set connecting \mathcal{G}_1 with another copy of \mathcal{G}_2 ; each coset of H inside G_2 is a border set connecting it with another copy of \mathcal{G}_1 . The copies of \mathcal{G}_1 and \mathcal{G}_2 form a tree, it is a G-tree (see [2] Section 8.6) which is usually called the Bass-Serre tree of the amalgamated product.

Let us pick one of the pieces, say \mathcal{G}_1 , and call it a root P_0 . It contains $[G_1 : H]$ border sets B_λ , $\lambda \in \Lambda_1$. All of these P_λ , $\lambda \in \Lambda_1$, are in the same isomorphism class, say I_2 , because there exists an isomorphism that takes P_λ to $P_{\lambda'}$ with $\lambda, \lambda' \in \Lambda_1$, and carries descendants to descendants. This isomorphism is just a left multiplication by an element of $G_1 *_H G_2$. Children of pieces in the isomorphism class I_2 are in the same isomorphism class again, denote it by I_1 . So the isomorphism class of a piece depends only on the parity of the generation. More precisely there are three isomorphism classes: the root, the set of pieces of odd generations and the set of pieces of even generations. The root differs from an even generation piece by the number of children, but it is only one such exceptional piece. No color assigned to this piece is in \mathbf{Y} by Remark 4.7, so we do not need to consider this piece in the remaining computations.

The parity of a generation implies that the branching process defined in Section 4 is periodic and so the matrix M has always two anti-diagonal blocks as in the case of $SL_2(Z)$, see (14).

6.2. HNN extensions with standard generating sets. Recall that an HNN extension G is constructed from a base group G_1 having a presentation $G_1 = \langle S \mid R \rangle$, and from an isomorphism α between two subgroups H and K of G_1 . Let t be a new symbol not in S (free letter), and define $G = G_1 *_\alpha = \langle S, t \mid R, tht^{-1} = \alpha(h), \forall h \in H \rangle$.

Consider the Cayley graph \mathcal{G} of this HNN extension G with respect to the generating set $S \cup \{t\}$. The Cayley graph \mathcal{G}_1 of G_1 is a part of the Cayley graph \mathcal{G} . Each coset of H (resp. K) is attached to another copy of \mathcal{G}_1 , the attachment is done by edges labeled by t , these edges correspond to the isomorphism α . Therefore the piece \mathcal{G}_1 is connected to $[G_1 : H] + [G_1 : K]$

other pieces. This way we obtain the Bass-Serre tree of the HNN extension.

Let us denote by P_0 a subgraph of \mathcal{G} containing the graph \mathcal{G}_1 and all t -edges incident to at least one vertex in \mathcal{G}_1 (and we also add its other endpoint). In \mathcal{G} , the that contains the copy of \mathcal{G}_1 at v and all t -edges incident to at least one vertex in this copy that are not in P_0 . The border set B_1 consists of vertices in H in the copy of \mathcal{G}_1 . Repeating this for all t -edges emerging from P_0 we obtain $[G_1 : K]$ pieces in the first generation. Similarly by following edges labeled by t^{-1} emerging from P_0 we obtain $[G_1 : H]$ other pieces in the first generation. Then we do the same for next generations. This procedure, gives the tree-like structure of the graph. Each piece P_i contains a copy of \mathcal{G}_1 and some t -edges connected to it (in fact it contains all such t -edges except the $|K|$ edges that connect the border set B_i and the parent). There are three isomorphism classes of pieces: the root, pieces whose border sets are copies of K and pieces whose border sets are copies of H . In this case the process is not periodic since a piece of each class may be incident to some other piece of the same class.

6.3. *Graphs of groups.* Let us recall the definition of the fundamental group of a graph of groups [2].

DEFINITION 6.1. A graph of groups \mathfrak{G} consists of

- (i) a connected graph X with vertex set $V(X)$ and edge set $E(X)$,
- (ii) for each vertex v of X a group G_v , and for each edge e of X a group G_e , and
- (iii) for each edge $e = (v_1, v_2)$ monomorphisms $\tau : G_e \rightarrow G_{v_1}$ and $\sigma : G_e \rightarrow G_{v_2}$.

Denote by E be the free group with basis $\{t_e; e \in E(X)\}$. Let $F(\mathfrak{G})$ be the group $(E * \ast_{v \in V(X)} G_v) / N$, where N is the normal closure of the subset $\{t_e^{-1} \tau(a) t_e \sigma(a)^{-1} : e \in E(X), a \in G_e\}$.

Let T be a maximal tree in X . We define the fundamental group $\pi(\mathfrak{G}, X, T)$ to be $F(\mathfrak{G}) / M$, where M is the normal closure of $\{t_e, e \in E(T)\}$.

Note that the groups $\pi(\mathfrak{G}, X, T)$ are independent of T up to isomorphism.

We will consider the Cayley graph \mathcal{G} of the fundamental group $\pi(\mathfrak{G}, X, T)$ with the following generators: $(\bigcup_{v \in V(X)} S_v) \cup \{t_e : e \in E(X) \setminus E(T)\}$, where $G_v = \langle S_v \rangle$ for $v \in V(X)$. This set of generators depends on T . We call this set of generators *standard*. In order to obtain locally finite graph we will consider only finite sets of generators, in particular this restricts us to finite graphs X of finitely generated groups.

Observe that if X consists of one edge between two distinct vertices, then its fundamental group is an amalgamated product of the vertex groups. If the two vertices coincide, then the fundamental group is a HNN extension. The tree-like structure we described for these specific cases will be now generalized to the Cayley graph of any (finite) graph of groups.

THEOREM (1.4). *The Cayley graph of the fundamental group of a graph of groups has a tree-like structure.*

PROOF. We will define the pieces and verify the conditions (1)-(5) in Definition 2.1.

First observe that the Cayley graph \mathcal{G} is covered by translates of the Cayley graphs \mathcal{G}_v of the vertex groups G_v , $v \in V(X)$ by left multiplications by elements of G , and edges labeled by t_e , $e \in E(X) \setminus E(T)$. For each $e = (v_1, v_2) \in E(X) \setminus E(T)$ denote by R_e the set of edges labeled by t_e such that they start at a vertex of \mathcal{G}_{v_1} and end at a vertex of the translate of \mathcal{G}_{v_2} by t_e . Then we denote by $g\mathcal{G}_v$ the translate of the subgraph \mathcal{G}_v by g , the representative of a coset in $\pi(\mathfrak{G}, X, T)/G_v$. Denote by ghR_e the translate of the set R_e by gh , where g is a representative of a coset in $\pi(\mathfrak{G}, X, T)/G_{v_1}$ and h is a representative of a coset in $G_{v_1}/\tau(G_e)$. These sets will be used in the construction of pieces of the tree-like structure.

Consider the quotient map $\pi(\mathfrak{G}, X, T) \rightarrow \pi_1(X)$ taking all G_v to identity. The fundamental group $\pi_1(X)$ of a finite graph X is free of rank $|E(X) \setminus E(T)|$. Every translate of R_e is a preimage of one edge in the Cayley graph of $\pi_1(X)$, thus it is an edge cut set of \mathcal{G} . Moreover, if $e = (v_1, v_2) \in E(T)$, then any translate $g\tau(G_e)$ by $g \in \pi(\mathfrak{G}, X, T) \rightarrow \pi_1(X)$ is a vertex cut set.

We say that $g_1\mathcal{G}_{v_1}$ and $g_2\mathcal{G}_{v_2}$ are neighbors if there is an edge $e = (v_1, v_2) \in E(T)$ (or $e = (v_2, v_1)$) and $g_1\mathcal{G}_{v_1} \cap g_2\mathcal{G}_{v_2} \neq \emptyset$. We say that $g_1\mathcal{G}_{v_1}$ and $g_2\mathcal{G}_{v_2}$ are delayed neighbors if there is an edge $e = (v_1, v_2) \in E(X) \setminus E(T)$ ($e = (v_2, v_1)$ resp.) and there is translate hR_e of a set of edges labeled by t_e connecting a vertex of $g_1\mathcal{G}_{v_1}$ to a vertex of $g_2\mathcal{G}_{v_2}$, we say the neighbors are delayed by hR_e .

Consider a graph with vertices $g\mathcal{G}_v$ and edges between neighbors and delayed neighbors. This graph is a tree. Assume not and let there be a cycle. If some edge of the cycle arises from delayed neighbors, then this R_e is not a cut set. If all edges in the cycle are neighbors, they induce a loop in T , which is also a contradiction. We choose a root of this tree to be \mathcal{G}_{v_0} for some $v_0 \in V(X)$.

Now we are ready to define pieces and border sets. We take the tree from above and define the pieces: For the root \mathcal{G}_{v_0} we define a piece $P_{v_0, o}$ to be a union of \mathcal{G}_{v_0} and sets hR_e for all its delayed neighbors. For any other $g\mathcal{G}_v$

we define a piece $P_{v,g}$ to be a union of \mathcal{G}_{v_0} and sets hR_e for all its delayed neighbors that are not parent of $g\mathcal{G}_v$ in the above tree order. Define the border sets of a piece $P_{v,g}$ to be its intersection with the parent piece. For the root define the border set to contain only the origin.

Let us now verify the conditions in Definition 2.1:

- (1) The Cayley graphs \mathcal{G}_v are connected and the pieces remain connected after adding the incident edges t_e . The intersection of P_{v_1,g_1} with its parent P_{v_2,g_2} is a translate of $\tau(G_e)$ ($\sigma(G_e)$ resp.) inside P_{v_1,g_1} provided $e = (v_1, v_2)$ ($e = (v_2, v_1)$ resp.). Thus the border set is indeed subset of the vertex set of the piece.
- (2) We need to show that every edge $e \in E(\mathcal{G})$ is in exactly one P_i . Clearly this holds for edges labeled by $s \in S_v$. Every edge labeled by t_e has endpoints in two different pieces, which by the construction become parent and child, and thus it is included only in the parent piece.
- (3) The tree was defined above, such that the border sets satisfy required condition (3).
- (4) For all $j \neq 0$ the border set B_j is a cut set. Using the notation $U(P_i)^c$ for the subgraph induced on edges in $E(G) \setminus E(U(P_i))$, we obtain $U(P_i) \cap U(P_i)^c = B_i$. So claim (4) follows.
- (5) Now we want to show that there are only finitely many isomorphism classes of pieces. The root is a unique piece of its class and we exclude it from the following consideration. Assume P_i was obtained from translate of \mathcal{G}_{v_1} and its parent from \mathcal{G}_{v_2} that are neighbors due to an edge $e = (v_1, v_2) \in E(X)$. This edge together with its direction characterizes the isomorphism class of P_i . Indeed if P_j is characterized by the same edge $e = (v_1, v_2) \in E(X)$, then clearly there is an element $g \in \pi(\mathcal{G}, X, T)$ acting on the Cayley graph as an isomorphism f such that $f(P_i) = P_j$ and $f(B_i) = B_j$. Therefore $f(U(P_i)) = U(P_j)$ as well. Moreover, since the pieces (and border sets) are identified in each subtree by the same procedure, this isomorphism respects the structure of pieces in the subtree.

□

COROLLARY 6.2. *For the Cayley graph of a fundamental group of a finite graph of finite groups, with the natural generating sets, the p_c is an algebraic number.*

PROOF. It follows from Theorem 1.1 (iii) and Theorem 1.4. □

REMARK 6.3. *A fundamental group of a graph of groups with trivial*

edge groups is a free product of its vertex groups and copies of \mathbb{Z} , so the first moment matrix M can be expressed as in the example 5.1.

6.4. *An example with infinite pieces.* Consider an amalgamated product $G_1 *_{\mathbb{Z}_2} G_2$ with $G_1 = \mathbb{Z}_2 \times \mathbb{Z}$ and $G_2 = \mathbb{Z}_4$. Its Cayley graph with respect to natural generators can be decomposed similarly as above, with border sets of size two. Therefore the set of partition $Z^{(i)}$ has two elements, and the set of colors \mathbf{Y} has four elements as for the Cayley graph of $Sl_2(\mathbb{Z})$. The first moment matrix M has again two anti-diagonal blocks.


Let A (B resp.) be the probability that the two vertices of the border set in G_1 (G_2 resp.) are connected (in the descendant subtree). Then A and B satisfy:

$$\begin{aligned} A &= 2p - p^2 + (1 - p)^2(2C - C^2), \\ B &= (2p - p^2)^2 A + (2p^2 - p^4)(1 - A), \\ C &= p^2(1 - (1 - p)^2(1 - B)) + (1 - p)^2(1 - B)C, \\ C &= \frac{p^2(1 - (1 - p)^2(1 - B))}{1 - p^2(1 - p)^2(1 - B)}. \end{aligned}$$

In order to express A , we split the piece $\mathbb{Z}_2 \times \mathbb{Z}$ into three parts, the first one containing only two edges between vertices of B_i . Then we split the rest at the B_i and obtain two other (identical) parts (corresponding to $\mathbb{Z}_2 \times \mathbb{N}$, see for example the last block in Figure 3). The probability that vertices of B_i are connected in one of these two latter parts is equal to C .

These formulas lead to a cubic equation, and exactly one of the solutions is real and in $[0, 1]$. Now we are ready to express the entries of first moment matrix M . There are two anti-diagonal blocks M_{12} and M_{21} , again the one corresponding to \mathbb{Z}_4 is already known from previous example. For the other factor group $\mathbb{Z}_2 \times \mathbb{Z}$ we denote by T_n the transition matrix from the parent to the n -th child (order them by distance from the origin). In order to obtain the transition matrix, we split the graph of $\mathbb{Z}_2 \times \mathbb{Z}$ into several blocks, see Figure 3. For each block we express the transition probabilities and thus T_n is composition of these probabilities, in particular it is a product of matrices corresponding to each block.

The first block consists of vertices to the left from the parent and including the parent. The probability of the connection in the left part is C and the probability that the two vertices in B_0 are connected is $2p - p^2$. This gives the first matrix in formula (15). The second block is used to express matrix T i.e. the transition to the first child B_1 from the parent B_0 . The third one is repeated $(n - 1)$ -times in order to reach the n -th child. It differs from


 FIG 3. Blocks of the Cayley graph of $\mathbb{Z}_2 \times \mathbb{Z}$

T by the factor of B coming from the descendant partition on the sibling border set (this connection is drawn by the dash-dotted line). The last factor in the product corresponds to the rightmost part of the picture and equals the probability of connection on the right side of the child in question (it is again equal to C).

$$\begin{aligned}
 (15) \quad T_n &= \begin{bmatrix} P_p(r_n|r) & P_p(b_n|r) \\ P_p(r_n|b) & P_p(b_n|b) \end{bmatrix} \\
 &= \begin{bmatrix} 1 & 0 \\ 1 - (1 - C)(1 - p)^2 & (1 - C)(1 - p)^2 \end{bmatrix} T \\
 &\quad \cdot \left(\begin{bmatrix} 1 & 0 \\ B & 1 - B \end{bmatrix} T \right)^{n-1} \begin{bmatrix} 1 & 0 \\ C & (1 - C) \end{bmatrix}, \\
 T &= \begin{bmatrix} P_p(r|r) & P_p(b|r) \\ P_p(r|b) & P_p(b|b) \end{bmatrix} = \begin{bmatrix} p^2 + 2p(1 - p)(2p - p^2) & 2p(1 - p)^3 \\ p(2p - p^2) & p(1 - p)^2 \end{bmatrix}, \\
 (16) \quad M_{21} &= \begin{bmatrix} E_p(r|r) & E_p(b|r) \\ E_p(r|b) & E_p(b|b) \end{bmatrix} = 2 \sum_{n=1}^{\infty} T_n \\
 &= 2 \begin{bmatrix} 1 & 0 \\ C & (1 - C) \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 2p - p^2 & (1 - p)^2 \end{bmatrix} T \\
 &\quad \cdot \left(1 - \begin{bmatrix} 1 & 0 \\ B & 1 - B \end{bmatrix} T \right)^{-1} \begin{bmatrix} 1 & 0 \\ C & (1 - C) \end{bmatrix}
 \end{aligned}$$

Again we can solve the equation $\det(M - 1) = 0$ and we obtain $p_c = .2951 \dots$

6.5. *Groups acting on trees with finite vertex stabilizers.* The goal of this section is to prove Theorem 1.5. To this end, we combine the result about standard generating sets (Theorem 1.4) with the general results from Section 2.

THEOREM (1.5). *Let G be a virtually free group, that is, it acts on a*

simplicial tree T with finite vertex stabilizers. Then its Cayley graph with respect to any finite generating set has a tree-like structure with finite pieces. Given a finite generating set of G , the pieces of the tree-like structure are algorithmically constructed.

Therefore the p_c is an algebraic number and one can use the algorithm from Theorem 1.1 (iii) to compute p_c , given any finite generating set.

PROOF. The group G in this case is a fundamental group of a finite graph of finite groups. Recall that the standard generating set, as defined in Section 6.3, consists of generators of the vertex groups and free letters corresponding to the edges outside the spanning tree of the factor graph T/G . Assume S_1 is maximal generating set such that it contains all elements of the vertex groups and free letters. In Section 6.3 we constructed a tree-like structure for the Cayley graph, with pieces corresponding to vertex groups. This tree of pieces P_i is a starting point of our generalization to an arbitrary generating set.

Denote by m the number of vertex groups in T/G . Elements of an arbitrary (finite) generating set S can be represented by reduced words in S_1 of bounded length (depending on S). Denote the maximal length by N .

The next step is to apply Lemma 2.6 several times, so that the piece P'_i contains all its descendants up to n -th generation. Every application of that lemma enlarge the pieces by one generation. We will find n large enough to guarantee that there exists a piece containing both endpoints of each edge labeled by an element in S .

Let P'_i be a piece in the n -times “enlarged” tree-like structure. Then $V(P'_i) = \bigcup_{k \in \Lambda_i(n)} V(P_k)$, where $\Lambda_i(0) = \{i\}$ and $\Lambda_i(n+1) = \{i\} \cup \bigcup_{s \in \Lambda_i(n)} \Lambda_s$. Since the original pieces were finite, Λ_j 's are finite and so are the modified pieces P'_i . An intuitive picture of how such enlarged pieces look like, comes from the free groups, see Corollary 7.6.

Recall that the pieces P_i correspond to transitions of vertex groups gG_v , and they are connected in the tree of pieces if they are so-called neighbors (corresponding to edge in spanning tree) or delayed neighbors (corresponding to other edges in the graph of groups), see Section 6.3. Now we can use a graph distance between pieces. Let $s \in S_1$ and x be a vertex in some piece $g_1G_{v_1}$. How far is a piece containing the vertex xs ?

Assume $s \in G_{v_2}$, so it is in some vertex stabilizer. There is a path between v_1 and v_2 in the spanning tree of the graph T/G , visiting vertices $v_1 = u_1, u_2, \dots, u_k = v_2$. The size of the spanning tree is equal to the the number of vertex groups minus 1, that is $m - 1$. For every i there is $g_i \in G$ such that $x \in g_iG_{u_i}$. Therefore there is a path in the tree of pieces through vertices

$g_1G_{u_1}, g_2G_{u_2}, \dots, g_kG_{u_k}$. Moreover, $xs \in g_kG_{v_2}$. Thus the distance between pieces containing x and xs is at most $m - 1$.

Assume s is a free letter. Then it corresponds to an edge e in the graph of groups outside the spanning tree, starting at vertex v_2 and terminating at vertex v_3 . As before there is a path from $g_1G_{v_1}$ to a piece $g_kG_{v_2}$, and $xs \in g_{k+1}G_{v_3}$ a delayed neighbor of $g_kG_{v_2}$. Thus the distance between pieces containing x and xs is at most m , in this case. Similarly for s^{-1} .

Let $x \in G$ with word length (in S_1) at most N . The distance between pieces containing g and gx is at most mN , for any $g \in G$. If the distance between two pieces is at most mN , then their closest common ancestor differs from them by less than mN generations. In particular, if $\text{dist}(P_i, P_j) \leq mN$, then there is P_k , such that P'_k contains vertices of P_i and P_j provided $n > mN$.

Now we can apply Lemma 2.5 to conclude that the Cayley graph with respect to $S \cup S_1$ has a tree-like structure with finite pieces. Consequently by Lemma 2.4 the Cayley graph with respect to S has also a tree-like structure with finite pieces. \square

7. Transitive graphs with more than one end. In what follows we will generalize the above decomposition into the tree-like structure, which was natural for amalgamated products and HNN extensions, to the case of transitive graphs with more than one end.

Recall that the number of ends of a graph is the supremum of the number of connected components of any of its subgraphs that was obtained by removing a finite set of vertices. An infinite transitive graph can have one, two or infinitely many ends.

7.1. *Dunwoody's result.* First let us recall the notation and a result obtained by Dunwoody in [3].

Let \mathcal{G} be an infinite connected graph with more than one end. Let c be a subset of vertices of \mathcal{G} , denote by $c^* = V(\mathcal{G}) \setminus c$ and by ∂_{Ec} the set of edges having one endpoint in c and the other in c^* . Denote by $\partial_V c$ the set of vertices in c having a neighbor in c^* . A set of vertices c such that ∂_{Ec} is finite is called a cut. A cut is said to be non-trivial if both c and c^* are infinite.

LEMMA 7.1 ((Dunwoody)). *Let \mathcal{G} be a graph with more than one end and let $H \subset \text{Aut}(\mathcal{G})$. There exists a non-trivial cut d such that for any $g \in H$ one of the inclusions $d \subset gd$, $d \subset gd^*$, $d^* \subset gd$, $d^* \subset gd^*$ holds.*

Let d be a cut satisfying the above lemma and for any other cut b we

denote

$$\begin{aligned}
T_b &= \{gd \mid g \in \text{Aut}(\mathcal{G}), gd \subsetneq b, \text{ and there is no } h \in \text{Aut}(\mathcal{G}) \\
&\quad \text{such that } gd \subsetneq hd \subsetneq b \text{ or } gd \subsetneq hd^* \subsetneq b\}, \\
T_b^* &= \{gd^* \mid g \in \text{Aut}(\mathcal{G}), gd^* \subsetneq b, \text{ and there is no } h \in \text{Aut}(\mathcal{G}) \\
&\quad \text{such that } gd^* \subsetneq hd \subsetneq b \text{ or } gd^* \subsetneq hd^* \subsetneq b\}, \\
Q_1 &= d \cap \bigcap_{c \in T_d \cup T_d^*} (c^* \cup \partial_V c), \\
(17) \quad Q_2 &= d^* \cap \bigcap_{c \in T_{d^*} \cup T_{d^*}^*} (c^* \cup \partial_V c).
\end{aligned}$$

If there is $g \in \text{Aut}(\mathcal{G})$ such that $gd = d^*$ the above definition gives us $T_d = T_d^*$. In order not to consider each cut twice, we set $T_d^* = \emptyset$ in this case.

Now we will decompose the graph into pieces isomorphic to Q_1 and Q_2 .

LEMMA 7.2. *Let $c_1, c_2 \in T_d \cup T_d^*$ and $c_1 \neq c_2$. Then the distance from $c_1 \setminus Q_1$ to $c_2 \setminus Q_1$ in \mathcal{G} is at least 2. Moreover, $d = Q_1 \dot{\cup} \bigcup_{c \in T_d \cup T_d^*} (c \setminus \partial_V c)$. Similarly for d^* .*

PROOF. From the property that there is no $h \in \text{Aut}(\mathcal{G})$ such that $c_i \subsetneq hd \subsetneq d$ or $c_i \subsetneq hd^* \subsetneq d$ for $i = 1, 2$ it follows that $c_1 \not\subset c_2$ and $c_1^* \not\subset c_2^*$. Since $d^* \subset c_i^*$, we have $c_1^* \not\subset c_2$. Therefore by Dunwoody's result in Lemma 7.1 $c_1 \subset c_2^*$ and $c_2 \subset c_1^*$. If $\partial_V c_i \subset Q_1$ the claim would follow.

Assume $c \in T_d \cup T_d^*$ such that $\partial_V c \not\subset Q_1$. It means that there is $c' \in T_d \cup T_d^*$ such that $(c' \cup \partial_V c') \cap d$ does not contain $\partial_V c$, and therefore $(c' \setminus \partial_V c') \cap \partial_V c \neq \emptyset$. But from above we have that $c' \subset c^*$, what is a contradiction. \square

THEOREM 7.3. *Every transitive graph with more than one end admits a tree-like structure with finite border sets.*

PROOF. The pieces we consider are Q_1, Q_2 and their translates. Assume d is a cut satisfying the above Dunwoody's Lemma 7.1 and that the origin is in $\partial_V d^*$. Let the root piece P_0 be a subgraph of \mathcal{G} induced on the vertices $Q_2 \cup \partial_V d^*$.

Now with each translate of d (d^* resp.) we have a translate of Q_1 (Q_2 resp.). Consider the cuts in $T_{d^*} \cup T_{d^*}^*$ and the translates of Q_1 and Q_2 corresponding to them. The pieces of the first generation are subgraphs induced by the vertices of these translates of Q_1 and Q_2 (we include all the edges that are not already in P_0). Let the first generation also contain a piece induced by edges in the original Q_1 .

The whole tree of pieces is constructed inductively. We always consider translates of Q_1 and Q_2 corresponding to the cuts in $T_b \cup T_b^*$, where b are the cuts used in the previous generation. The piece is a subgraph induced on the vertices of the appropriate translate of Q_1 and Q_2 , and we exclude all edges which are already in some piece of previous generation. The border set B_i is defined as the intersection of P_i with its parent ($B_0 := \{o\}$).

Now we can start verifying the properties of Definition 2.1.

- (1) The pieces and border sets such that B_i is a subset of vertex set of P_i .
- (2) Every edge is in exactly one piece - in the first appearing in the construction and containing both endpoints of the edge.
- (3) The partial order of the elements follows from the construction, as well as the property that border set is the intersection of the piece with its parent.
- (4) If we remove a border set B_i the graph falls apart. In particular $U(P_i)$ is the subgraph induced by vertices of c' and by Lemma 7.2 it intersects the $U(P_i)^c$ only by vertices in $\partial_V c$, which are in B_i .
- (5) There are two isomorphism classes of pieces (excluding the root), based on whether P_i arises from transition Q_1 or Q_2 .

□

7.2. Application to free groups with non-standard generators. The simplest example of a transitive graph with infinitely many ends is a regular tree. The Cayley graph of free group with respect to free generators is a regular tree and the percolation on it is well understood. The result of this section gives us a simple way to find p_c for any finite generating set.

Assume $F_n = \langle x_1, \dots, x_n \rangle$. Let \mathcal{G} be its Cayley graph with respect to the standard (free) generators. Denote by H the subgroup of $\text{Aut}(\mathcal{G})$ generated by left translations by elements in F_n and isomorphisms arising from permutations of the generators.

The following lemma shows that the the cut d satisfying the condition in Dunwoody's Lemma 7.1 can be found explicitly.

LEMMA 7.4. *Consider the free group F_n with any finite set of generators. Denote by d the set of vertices labeled by words starting with letter x_1 (one element of any free generating set). Then d is such that for any $g \in H$ one of the inclusions $d \subset gd$, $d \subset gd^*$, $d^* \subset gd$, $d^* \subset gd^*$ holds.*

PROOF. For any finite generating set, d and d^* are infinite with finite boundaries. Indeed for any m there are only finitely many pairs $(a, b); a \in d, b \in d^*$ with the word distance less than m . Without loss of generality we

assume that the generating set is symmetric on the permutation of generators, so the whole group H acts on the Cayley graph by isomorphisms.

Now for any $h \in H$, if the origin 1 is contained in hd , then there is a word x such that all vertices in hd^* are labeled by reduced words starting with x and $x \in d^*$. If the origin is in hd^* , then hd has a similar property. Assume that for a given $h \in H$ such a word is w and the origin is in hd (or in hd^* respectively). Then if w starts with letter x_1 , then $d^* \subset hd^*$ (or $d^* \subset hd$ respectively), otherwise $d \subset hd$ (or $d \subset hd^*$ respectively). \square

The following theorem is a special case of Theorem 1.5. We include here a different, more hands-on proof. We present here the pieces of the tree-like structure of free groups explicitly. They are constructed using a cut set satisfying Dunwoody's Lemma 7.1. We could as well use the Lemma 2.6 (starting with pieces that are single edges of Cayley graph of free group with respect to free generators), the pieces obtained would be exactly the same.

THEOREM 7.5. *The Cayley graph of free group F_n (with n free generators) with respect to any finite generating set has a tree-like structure with finite pieces. Therefore its p_c is an algebraic number and one can use the algorithm from Theorem 1.1 (iii) to compute p_c , given any finite generating set.*

PROOF. Recall that we want to show that the Cayley graph of F_n with respect to any finite set of generators has a tree-like structure with finite pieces.

Any finite set of generators is contained in some ball S_k , i.e. the set of all words of length at most k (in the standard word metric). Therefore, using Lemma 2.4, it is enough to find the tree-like structure with finite pieces for the generating set S_k . The isomorphisms will be from the group H , which admits the group F_n as a finite index subgroup.

Consider the cut d from Lemma 7.4. We observed that always either hd or hd^* is a set of all vertices labeled by a reduced word starting with some x . We denote such a cut D_x and its complement (containing the origin) by D_x^* . For example for d we write D_{x_1} .

Applying the construction in the proof of Lemma 7.3 to the cut d we obtain the following tree-like structure. The pieces are isomorphic to Q_1 and Q_2 . Since d and d^* are isomorphic, it is enough to describe just one of

them, say Q_2 (general formula (17)).

$$T_{d^*} \cup T_{d^*}^* = \{D_x : |x| = 1, x \neq x_1\}$$

$$Q_2 = D_{x_1}^* \cap \bigcap_{|x|=1, x \neq x_1} (D_x^* \cup \partial_V D_x)$$

If a vertex is labeled by a reduced word longer than k , then it is in $D_x \setminus \partial_V D_x$ for some $|x| = 1$, thus not in Q_2 . Therefore $Q_2 \subset S_k$, hence it is finite.

Note that since $|T_{d^*} \cup T_{d^*}^*| = 2n - 1$ the branching number of the tree of pieces is $2n - 1$. □

The tree-like structure is explicitly given by the procedure of Lemma 7.3. For the Cayley graph of F_2 with respect to S_2 it results in pieces as in Figure 4. In the picture we see that the whole Q_2 splits into three parts. For the tree-like structure we can take pieces corresponding to these thirds of Q_2 . In particular the middle piece P_i (with labeled vertices in the picture) has border set $B_i = \{o, a\}$, and its children pieces (three of them) share with P_i the middle vertex a and one of the remaining three vertices (ab, a^2, ab^{-1}). The value of the critical probability is .139... in this case.




FIG 4. Piece in Cayley graph of F_2 with respect to S_2

If the generating set is a general ball S_k we can identify the following tree-like structure.

COROLLARY 7.6. *Consider the free group $F_2 = \langle a, b \rangle$ and its Cayley graph with respect to the generating set S containing all words of length at*

most k (in the standard word metric). There is a tree-like structure with border sets of size $\frac{3^{k-1}+1}{2}$, pieces of size $\frac{3^k+1}{2}$, and the tree of pieces has branching number 3.

PROOF. The root piece P_0 is a subgraph induced by vertices of S_{k-1} with border set containing the origin. The four pieces of first generation are subsets of S_k such that they contain the origin and vertices labeled by words starting with a specific letter. The children of the piece corresponding to the letter x are induced by subsets of xS_k containing x and vertices labeled by words starting with xy , where $x \neq y \in \{a, a^{-1}, b, b^{-1}\}$. Clearly each such piece has exactly three children. Since $|S_k| = 4 * (3^k - 1)/(3 - 1) + 1$ we can see $|P_i| = (|S_k| - 1)/4 + 1 = (3^k + 1)/2$ for all $i \neq 0$. Again we included every edge in exactly one piece, that is in the oldest one containing both its endpoints. The border set is defined as the intersection of a piece with its parent. It is not difficult to see that B_i coincides with vertices of a piece in Cayley graph with respect to S_{k-1} , and thus $|B_i| = (3^{k-1} + 1)/2$. \square

Acknowledgments. The author would like to thank Mark Sapir, Tatiana Smirnova-Nagnibeda and Russell Lyons for fruitful conversations and useful advice. I am also grateful to Vadim Kaimanovich for bringing Dunwoody's result to my attention.

REFERENCES

- [1] Michael Aizenman and David J. Barsky, *Sharpness of the phase transition in percolation models*, Comm. Math. Phys. **108** (1987), no. 3, 489–526. MR874906
- [2] Daniel E. Cohen, *Combinatorial group theory: a topological approach*, London Mathematical Society Student Texts, vol. 14, Cambridge University Press, Cambridge, 1989. MR1020297
- [3] Martin J. Dunwoody, *Cutting up graphs*, Combinatorica **2** (1982), no. 1, 15–23.
- [4] Geoffrey Grimmett, *Percolation*, second ed., Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 321, Springer-Verlag, Berlin, 1999. MR1707339
- [5] H. Kesten, *The critical probability of bond percolation on the square lattice equals 1/2*, Communications in Mathematical Physics **74** (1980), 41–59.
- [6] Iva Kozakova, *Critical percolation on the free products*, International Journal of Algebra and Computation **18** (2008), no.4.
- [7] Russel Lyons and Yuval Peres, *Probability on trees and networks*, Cambridge University Press (To appear), <http://mypage.iu.edu/~rdlyons/prbtree/prbtree.html>.
- [8] Charles J. Mode, *Multitype branching processes. Theory and applications*, Modern Analytic and Computational Methods in Science and Mathematics, No. 34, American Elsevier Publishing Co., Inc., New York, 1971. MR0279901
- [9] Esa Nummelin, *General irreducible Markov chains and nonnegative operator*, Cambridge Tracts in Mathematics, No. 83, Cambridge University Press, Cambridge, 1984. MR776608

- [10] Jean-Pierre Serre, *Trees*, Springer-Verlag, Berlin, 1980, Translated from the French by John Stillwell. MR607504
- [11] Robert M. Ziff and Christian R. Scullard, *Exact bond percolation thresholds in two dimensions*, arXiv:cond-mat/0610813 v1 (2006).

MATHEMATICS DEPARTMENT, VANDERBILT UNIVERSITY
1326 STEVENSON CENTER, NASHVILLE, TN 37240
E-MAIL: iva.kozakova@vanderbilt.edu