ON NORMAL APPROXIMATIONS TO U-STATISTICS

By Vidmantas Bentkus, Bing-Yi Jing, and Wang Zhou

Vilnius Institute of Mathematics and Informatics, Hong Kong University

of Science and Technology, and National University of Singapore

Let X_1, \ldots, X_n be i.i.d. random observations. Let $\mathbb{S} = \mathbb{L} + \mathbb{T}$ be a U-statistics of order $k \geq 2$, where \mathbb{L} is a linear statistic having asymptotic normal distribution, and \mathbb{T} is a stochastically smaller statistic. We show that the rate of convergence to normality for \mathbb{S} can be simply expressed as the rate of convergence to normality for the linear part \mathbb{L} plus a correction term, $(\operatorname{var} \mathbb{T}) \ln^2(\operatorname{var} \mathbb{T})$, under the condition $\mathbb{E} \mathbb{T}^2 < \infty$. An optimal bound without this log factor is obtained under a lower moment assumption $\mathbb{E} |\mathbb{T}|^{\alpha} < \infty$ for $\alpha < 2$. Some other related results are also obtained in the paper. Our results extend, refine and yield a number of related known results in the literature.

1. Introduction. There has been a vast literature related to normal approximations and to the rates of convergence to normality for U-statistics of order $k \geq 2$. Undoubtedly, the case for k = 2 has been most studied and best understood. So we will start our discussion with this case as well. Let $X, X_1, X_2, ..., X_n, n \geq 2$, be independent and identically distributed (i.i.d.) random variables (r.v.'s). Define a U-statistic of order 2 by

(1.1)
$$U_n = \frac{2}{n(n-1)} \sum_{1 \le i \le j \le n} h(X_i, X_j),$$

where the kernel h(x, y) is a real-valued Borel measurable function, symmetric in its arguments with $Eh(X_1, X_2) = \theta$. By the Hoeffding decomposition,

^{*}Bing-Yi Jing's research was partially supported by Hong Kong RGC grants $\rm HKUST6011/07P$ and $\rm HKUST6015/08P$. Wang Zhou's research was partially supported by grant R-155-000-083-112 at National University of Singapore

AMS 2000 subject classifications: Primary 62E20

Keywords and phrases: U-statistics, Berry-Esseen bound, rate of convergence, central limit theorem, normal approximations, self-normalized, studentized U-statistics.

we have

$$\frac{\sqrt{n}(U_n - \theta)}{2\sigma_g} = \frac{1}{\sqrt{n}\sigma_g} \sum_{i=1}^n g(X_i) + \frac{1}{\sqrt{n}(n-1)\sigma_g} \sum_{1 \le i < j \le n} \eta(X_i, X_j),$$

$$(1.2) \qquad \stackrel{\text{def}}{=} \mathbb{L} + \mathbb{T},$$

where \mathbb{L} and \mathbb{T} are the linear and quadratic terms, respectively, and

$$g(x) = E[h(x, X_2)] - \theta, \qquad \sigma_g^2 = Var[g(X_1)],$$

 $\eta(x, y) = h(x, y) - g(x) - g(y) + \theta.$

Throughout the paper, it is assumed that $\sigma_g^2 \in (0, \infty)$.

For any generic r.v. Y, denote its distribution function (d.f.) by $F_Y(x) = \mathbb{P}(Y \leq x)$. Then,

$$F_{\mathbb{L}+\mathbb{T}}(x) = \mathbb{P}\left(\frac{\sqrt{n}(U_n - \theta)}{2\sigma_g} \le x\right).$$

It is well-known that $F_{\mathbb{L}+\mathbb{T}}(x)$ converges to the standard normal d.f., denoted by $\Phi(x)$, provided $Eh^2(X_1,X_2)<\infty$; see Hoeffding (1948). In fact, this moment condition can be reduced to $Eg^2(X_1)<\infty$ and $E|h(X_1,X_2)|^{4/3}<\infty$; see Remark 4.2.4 of Koroljuk and Borovskich [(1994), page 131]. There also has been much work on the convergence rates to normality for U-statistics of order 2. However, the sharpest Berry-Esseen bound of order $O(n^{-1/2})$ comes from Koroljuk and Borovskich (1985) and Friedrich (1989), who established the following ideal bound

(1.3)
$$||F_{\mathbb{L}+\mathbb{T}} - \Phi|| \ll \frac{1}{\sqrt{n}} \left(\frac{E|g_1|^3}{\sigma_g^3} + \frac{E|\eta_{12}|^{5/3}}{\sigma_g^{5/3}} \right),$$

where the following notation has been used:

- $g_i = g(X_i)$ and $\eta_{ij} = \eta(X_i, X_j)$.
- For any function $f: \mathbb{R} \to \mathbb{R}$, define $||f|| = \sup_{x \in \mathbb{R}} |f(x)|$.
- By $a \ll b$ for $a, b \ge 0$ we mean $a \le cb$ for some positive constant c not depending on the underlying distribution function.

Indeed, Bentkus *et al.* (1994) showed that the moment conditions in (1.3): $E|g_1|^3 < \infty$ and $E|\eta_{12}|^{5/3} < \infty$, are the weakest possible in the Berry-Esseen bounds of order $O(n^{-1/2})$ for *U*-statistics of order 2.

Our main purpose of this paper is to extend the optimal results on Berry-Esseen bounds for U-statistics of order 2 to those of higher orders. In stark contrast with studies on U-statistics of order 2, there is a very limited literature on optimal or near-optimal error bounds for U-statistics of higher orders. In this paper, we intend to fill in the gap. The work is of both theoretical and practical value since many symmetric statistics may be approximated arbitrarily closely by U-statistic of sufficiently high order under appropriate conditions. As an application, we will derive a near optimal error bound for studentized U-statistics of order 2 in Section 3.3.

At first sight, it appears to be an easy task to extend the optimal Berry-Esseen bounds from second order *U*-statistics to higher order ones. (At least we naively thought so in the beginning.) However, a close inspection soon leads one to believe that this is far from trivial. We note that the usual approach of dealing with *U*-statistics is to first use the *Hoeffding decomposition* to turn the statistic of interest into a sum of sequentially smaller and uncorrelated terms, and then to use *truncation techniques* to each of these terms. Broadly speaking, there are two main difficulties with this approach, which we must overcome. Specifically, we have the following goals in mind.

(A) A direct and transparent methodology is needed.

The truncation techniques are manageable for second order U-statistics. However, as the order gets larger, it becomes more and more intangible and unworkable. In order to treat U-statistics of higher order than 2, in this paper, we may have to abandon the idea of truncation, and instead choose a more direct and more transparent approach.

(B) A simple and unified form of error bounds is needed.

For second order U-statistics, the Hoeffding decomposition produces a linear statistic \mathbb{L} plus a degenerate U-statistic \mathbb{T} in (1.2). As a result, the two terms involving $E|g_1|^3$ and $E|\eta_{12}|^{5/3}$ in (1.3) are from the linear term \mathbb{L} and the degenerate U-statistic term \mathbb{T} , respectively. More generally, applying the Hoeffding decomposition to U-statistics of order k > 2 leads to a sum of k terms. Consequently, the resulting optimal error bound would contain k terms, which would become more complicated as k gets larger. In this paper, we strive to provide a simple and general error bounds for U-statistics of general orders.

Let us now look at (B) in more detail. Again, we will examine the case of U-statistics of order 2 first. A closer look at the proof of (1.3), as done in Koroljuk and Borovskich (1985) and Friedrich (1989), shows that a more refined Berry-Esseen bound can be given as follows:

$$(1.4) ||F_{\mathbb{L}+\mathbb{T}} - \Phi|| \ll \frac{1}{\sqrt{n}} \left(\frac{E|g_1|^3}{\sigma_g^3} + \frac{E|\eta_{12}|^{5/3}}{\sigma_q^{5/3}} + \frac{E|g_1g_2\eta_{12}|}{\sigma_g^3} \right).$$

Using some truncation argument, it can be shown that (1.4) is equivalent to

$$(1.5) ||F_{\mathbb{L}+\mathbb{T}} - \Phi|| \ll \frac{1}{\sqrt{n}} \frac{E|g_1|^3}{\sigma_a^3} + \frac{1}{n} \frac{E|\eta_{12}|^2}{\sigma_a^2} + \frac{1}{\sqrt{n}} \frac{E|g_1g_2\eta_{12}|}{\sigma_a^3}.$$

Note that the last term in (1.5) shows the interaction effect between \mathbb{L}^2 and \mathbb{T} in (1.2). This suggests that if we take the correlation between \mathbb{L} and \mathbb{T} into account, we might be able to improve the error bound. Indeed, if we let

$$N_2(x) = \Phi(x) + \frac{1}{\sqrt{n}} \kappa_2 \Phi'''(x), \text{ where } \kappa_2 = E g_1 g_2 \eta_{12} / \sigma_g^3,$$

be the *adjusted* normal approximation, Alberink and Bentkus (2001a, 2001b) showed that the following optimal error bound holds:

(1.6)
$$||F_{\mathbb{L}+\mathbb{T}} - N_2|| \ll \frac{1}{\sqrt{n}} \frac{E|g_1|^3}{\sigma_g^3} + \frac{1}{n} \frac{E|\eta_{12}|^2}{\sigma_g^2};$$

see Theorem 1 of Alberink and Bentkus (2001a, 2001b). We reiterate that the error bound given in (1.5) is optimal, and implies the optimal bound (1.3) as a special case by a simple truncation argument.

Finally, we could rewrite (1.5) in a very simple form

(1.7)
$$||F_{\mathbb{L}+\mathbb{T}} - N_2|| \ll ||F_{\mathbb{L}} - \Phi|| + \operatorname{var} \mathbb{T}.$$

In other words, the error bounds in the adjusted normal approximations for U-statistics of order 2 is simply the error bounds for the dominant linear part \mathbb{L} plus the variance of the remaining "error" term \mathbb{T} .

It is somewhat surprising that the optimal error bound for U-statistics of order 2 can be written in such a simple and compact form given by (1.7). The appearance of the variance term var \mathbb{T} in (1.7) is appealing since the variance is the most natural and easy-to-interpret measure to describe the

effects of the error term \mathbb{T} . Furthermore, from a computational point of view, it is easier to calculate the variance of \mathbb{T} than some other moments of \mathbb{T} .

The purpose of this paper is to derive optimal or near-optimal error bound in the normal approximations for U-statistics of order $k \geq 2$, similar to those given in (1.7). We emphasize that the truncation techniques are heavily used in Alberink and Bentkus (2001a, 2001b) for second order U-statistics, and will not work for U-statistics of higher orders.

The paper is arranged as follows. In Section 2, some definitions and notation will be given. In Section 3, we establish an optimal or near optimal error bounds for U-statistic of order $k \geq 2$. In Section 4, we give some approximations to expectations of smooth functions of U-statistics and some of their applications. Possible extensions to other statistics are discussed in Section 5. We provide the details of the proofs in Section 6.

2. Definitions and notation. Let X, X_1, \ldots, X_n be a sequence of i.i.d. r.v.'s. Suppose that a symmetric statistic $\mathbb{S} = \mathbb{S}(X_1, \ldots, X_n)$ can be decomposed into

$$(2.1) S = \mathbb{L} + \mathbb{T},$$

where \mathbb{L} and \mathbb{T} are symmetric *U*-statistics, \mathbb{L} of order 1 and \mathbb{T} of order k, respectively. This means that \mathbb{L} is of the form

$$(2.2) \mathbb{L} = \sum_{i=1}^{n} L_i$$

and \mathbb{T} is of the form, by way of Hoeffding decomposition,

(2.3)
$$\mathbb{T} = \sum_{i=1}^{n} T_i + \sum_{1 \leq i < j \leq n}^{n} T_{ij} + \dots + \sum_{1 \leq i_1 < \dots < i_k \leq n}^{n} T_{i_1 \dots i_k} \\
= \mathbb{T}_1 + \dots + \mathbb{T}_k,$$

where $L_i = l(X_i)$ and $T_{i_1...i_k} = t_k(X_{i_1},...,X_{i_k})$ are Borel measurable functions and t_k is invariant under permutation of variables. Without loss of generality, it is assumed that

(2.4)
$$\mathbb{E} \mathbb{L} = \mathbb{E} \mathbb{T} = 0, \text{ and } \text{var } \mathbb{L} = 1,$$

which can always be achieved after re-centering and re-scaling the relevant statistics. Typically, \mathbb{L} is the dominant term and \mathbb{T} can then be called the *error* term.

Now let us define, for $\alpha \in [1, 2]$,

$$\beta = n \mathbb{E} |L_1|^3, \qquad \gamma = \text{var } \mathbb{T}, \qquad \gamma^{(\alpha)} = \sum_{p=1}^k \binom{n}{p} \mathbb{E} |T_{1...p}|^{\alpha}.$$

Note that

$$\mathbb{E} |\mathbb{L}|^{3} \ll n \mathbb{E} |L_{1}|^{3} = \beta,$$

$$\mathbb{E} |\mathbb{T}|^{2} = \sum_{i=1}^{k} \mathbb{E} \mathbb{T}_{i}^{2} = \sum_{i=1}^{k} \binom{n}{i} \mathbb{E} |T_{1...i}|^{2} = \gamma,$$

$$(2.5) \qquad \mathbb{E} |\mathbb{T}|^{\alpha} \leq k^{\alpha-1} \sum_{i=1}^{k} \mathbb{E} |\mathbb{T}_{i}|^{\alpha} \ll \sum_{i=1}^{k} \binom{n}{i} \mathbb{E} |T_{1...i}|^{\alpha} = \gamma^{(\alpha)},$$

where the second inequality in (2.5) follows by first applying an inequality (2.18) from Gine, Latala and Zinn (2000):

$$\mathbb{E} |\mathbb{T}_i|^{\alpha} \le C(\alpha, i) \mathbb{E} |\sum_{1 \le j_1 < \dots < j_i \le n} T_{j_1 \dots j_i}^2|^{\frac{\alpha}{2}},$$

where $C(\alpha, i)$ is a constant depending only on α and i, and then applying the following simple inequality (noting $\alpha/2 \leq 1$):

$$\Big| \sum_{1 \le j_1 < \dots < j_i \le n} T_{j_1 \dots j_i}^2 \Big|^{\frac{\alpha}{2}} \le \sum_{1 \le j_1 < \dots < j_i \le n} |T_{j_1 \dots j_i}|^{\alpha}.$$

Therefore, β , γ and $\gamma^{(\alpha)}$ are closely related to $E|\mathbb{L}|^3$, \mathbb{ET}^2 and $\mathbb{E}|\mathbb{T}|^{\alpha}$, respectively. Furthermore, $\gamma = \gamma^{(2)}$.

Finally, the following convention will be adopted throughout the paper: conditions appearing in a statement are implied implicitly in that statement. For instance, in (3.4), the conditions $\beta < \infty$ and $\gamma < \infty$ are assumed even though (3.4) still holds true without these moment condition.

3. Error bounds for U-statistics. Asymptotic normality of the linear term \mathbb{L} and its rates of convergence are well-known. For instance, the latter is described by the Berry-Esseen bound:

If $\mathbb{T} \to 0$ in probability, we have $||F_{\mathbb{S}} - \Phi|| \to 0$ as $n \to \infty$. Following the discussion in the Introduction, we are interested in the error bound for $||F_{\mathbb{S}} - \Phi||$ in the form of

$$(3.2) ||F_{\mathbb{S}} - N_p|| \ll ||F_{\mathbb{L}} - \Phi|| + f(\mathbb{E}|\mathbb{T}|^{\alpha}) \ll \beta + f(\gamma^{(\alpha)}),$$

for some function f, N_p for some $p \ge 0$ (to be defined below), and $\alpha \le 2$. In particular, when $\alpha = 2$, (3.2) becomes

$$||F_{\mathbb{S}} - N_p|| \ll ||F_{\mathbb{L}} - \Phi|| + f(\operatorname{var} \mathbb{T}) \ll \beta + f(\gamma).$$

In other words, we wish to investigate the question: if a linear statistic \mathbb{L} is perturbed by some error term \mathbb{T} , how do we correct the usual Berry-Berry bound for normal approximations, using only the variance of \mathbb{T} ?

The adjusted normal approximation N_p (which was mentioned in (3.2)) can be defined as follows. Define $N_0(x) = \Phi(x)$ and

(3.3)
$$N_p(x) = \Phi(x) + G_1(x) + \dots + G_p(x), \quad p = 1, 2, \dots$$

where, for $p = 1, \ldots, k$,

$$\kappa_p = \operatorname{cov}(\mathbb{L}^p, \mathbb{T}) = \mathbb{E} \mathbb{L}^p \mathbb{T} = \binom{n}{p} \mathbb{E} L_1 \dots L_p \mathbb{T},$$

$$G_p(x) = (-1)^{p+1} \kappa_p \Phi^{(p+1)}(x).$$

Two questions immediately arise from the above definition:

1. Under what conditions can we guarantee the existence of κ_p for $p \geq 1$? By the Holder's inequality, we have

$$\mathbb{E} |L_1 \dots L_i \mathbb{T}| \leq \left(\mathbb{E} |L_1|^3 \dots \mathbb{E} |L_i|^3 \right)^{1/3} \left(\mathbb{E} |\mathbb{T}|^{3/2} \right)^{2/3}.$$

Therefore, κ_p exists if $\mathbb{E} |\mathbb{T}|^{3/2} < \infty$ as we assume that $\mathbb{E} |L_i|^3 < \infty$ for $1 \leq i \leq n$. If $\mathbb{E} |T|^{3/2} = \infty$, we will see later that we only use $N_0(x) = \Phi(x)$.

2. How does one interpret N_p ?

The adjusted normal approximation N_p plays a central role in this paper. For U-statistics of order 2, first appeared in Alberink and Bentkus (2001a), we have $\kappa_1 = 0$ and $\kappa_2 = -n^{-1/2} \mathbb{E} g_1 g_2 \eta_{12} / \sigma_g^3$, hence,

$$N_2(x) = \Phi(x) - \frac{1}{\sqrt{n}} \frac{\mathbb{E} g_1 g_2 \eta_{12}}{\sigma_g^3} \Phi'''(x)$$

Note that N_2 is different from its corresponding Edgeworth expansion:

$$F_{\mathbb{L}+\mathbb{T}}(x) = \Phi(x) - \frac{1}{\sqrt{n}} \left(\frac{\mathbb{E} g_1 g_2 \eta_{12}}{2\sigma_g^3} + \frac{\mathbb{E} g_1^3}{6\sigma_g^3} \right) \Phi'''(x) + o(n^{-1/2}),$$

which holds under the following optimal conditions: (i) d.f. of $g(X_1)$ is nonlattice; (ii) $\mathbb{E} |g_1|^3 < \infty$ and $\mathbb{E} |\eta_{12}|^{5/3} < \infty$; see Jing and Wang (2003). On the other hand, N_p 's and Edgeworth expansions are similar in the sense that both try to improve the normal approximations by including some higher order correction terms. We point out that Edgeworth expansions do require some smoothness conditions while Berry-Esseen bounds do not.

Of course, the adjusted normal approximations N_p are introduced not to compete with Edgeworth expansions. Rather, it is a convenient tool in obtaining a neat and unified error bound in the normal approximation for U-statistics of general order.

In the following, we will give error bounds under two different sets of conditions: $\mathbb{E} \mathbb{T}^2 < \infty$, and $\mathbb{E} |\mathbb{T}|^{\alpha} < \infty$ for $\alpha \in [1,2)$, respectively. Since the Hoeffding decomposition of \mathbb{T} requires $\mathbb{E} |\mathbb{T}| < \infty$, we will not consider the case $\mathbb{E} |\mathbb{T}|^{\alpha} < \infty$ for $\alpha \in (0,1)$.

3.1. Error bounds under $\mathbb{ET}^2 < \infty$. Here is our key result.

Theorem 1. We have

$$(3.4) ||F_{\mathbb{S}} - N_k|| \ll \beta + \gamma \ln^2(\gamma)$$

where the last term $\gamma \ln^2(\gamma)$ is taken to be zero when $\gamma = 0$.

For second order *U*-statistics, Bentkus and Alberink (2001a, 2001b) showed

$$||F_{\mathbb{S}} - N_k|| \ll \beta + \gamma,$$

which is sharper than that in Theorem 1 when k=2. We conjecture that the logarithmic factor in Theorem 1 can be removed for U-statistics of order k>2. Actually, we have derived an error bound without log factors, which for the U-statistics of order 2 implies the result of Bentkus and Alberink (2001a, 2001b). Unfortunately, for $k \geq 3$, the error bound involves certain

conditional variances, and the proof of this bound is indeed very complicated, compared to the proof of Theorem 1. We do not provide this bound here since in applications this bound has only minor advantages, compared to the simple and indeed convenient bound of Theorem 1.

Applying Theorem 1, we can derive the following optimal error bound for $||F_{\mathbb{S}} - \Phi||$ for *U*-statistics of order *k*. (A similar theorem to (3.5) is also derived by Chen and Shao (2007), using Stein's method.)

Theorem 2. We have

Furthermore, the exponent 1/2 in (3.5) is the best possible for U-statistics of order $k \geq 2$.

Comparing Theorems 1 and 2, we see that, the adjusted normal approximation $N_k(x)$, which takes into account of the correlations between \mathbb{L}^k and \mathbb{T} , does have improvement over the standard normal distribution $\Phi(x)$.

3.2. Error bounds under $\mathbb{E} |\mathbb{T}|^{\alpha} < \infty$ for $\alpha \in [1, 2)$. Theorem 1 contains a log factor, which will be inherited when applying the result. We will now provide an optimal error bound without the log factor under a lower moment assumption $\mathbb{E} |\mathbb{T}|^{\alpha} < \infty$ for $\alpha \in [1, 2)$.

To do this, we define $N_0 = \Phi$ and

$$N_p = \Phi + G_1 + \dots + G_p,$$

where $p = p(\alpha, k)$ is the largest integer such that $p < (\alpha - 1)/(2 - \alpha)$ and $p \le k$. For example, if $\alpha \le 3/2$, then $N_p = \Phi$. If $3/2 < \alpha \le 5/3$ and $k \ge 1$, then $N_p = \Phi + G_1$. If $5/3 < \alpha \le 7/4$ and $k \ge 2$, then $N_p = \Phi + G_1 + G_2$.

Theorem 3. For $\alpha \in [1,2)$, let $p = p(\alpha,k)$ be defined above, we have

$$||F_{\mathbb{S}} - N_p|| \ll \beta + \gamma^{(\alpha)}.$$

Theorem 3 would be of most interest when α is close to 2 in applications. As an example, we will apply Theorem 3 below to obtain an near optimal Berry-Esseen bound for Studentized U-statistics of order 2.

3.3. An application to Studentized U-statistics of order 2. The optimal error bound for standardized U-statistics of order 2 was given in (1.3). We conjecture that the same optimal error bound applies to its corresponding Studentized U-statistics defined by

$$\widehat{\mathbb{S}} = \frac{\sqrt{n}(U_n - \theta)}{2\widehat{\sigma}_q},$$

where $\hat{\sigma}_g^2$ is the jackknife variance estimator:

$$\widehat{\sigma}_g^2 = \frac{n-1}{(n-2)^2} \sum_{i=1}^n \left(\frac{1}{n-1} \sum_{j=1, j \neq i}^n h_{ij} - U_n \right)^2.$$

However, we have so far only managed to prove a near-optimal result, as given in Theorem 4 below. Although this result is the best available bound in the literature, it falls a little short of the optimal bound (i.e., $\epsilon = 0$).

Theorem 4. For any $\epsilon > 0$, we have

$$||F_{\widehat{\mathbb{S}}} - \Phi|| \ll \frac{1}{\sqrt{n}} \left(\frac{E|g_1|^3}{\sigma_g^3} + \frac{E|\eta_{12}|^{5/3+\epsilon}}{\sigma_g^{5/3+\epsilon}} \right). \quad \Box$$

The proof of Theorem 4 is very involved, and hence will not be given here. Its proof can be obtained from the authors. The proof basically involves approximating $\widehat{\mathbb{S}}$ by a U-statistic of sufficiently high order and then applying nontrivial truncation techniques. Currently we are trying to eliminate ϵ in Theorem 4, and hope to be able to report on this in the near future.

4. Error bounds for expectations of smooth functions. Here, we give some approximations to expectations of smooth functions of U-statistics. These results (e.g., Theorems 5 and 6) are of interest in their own right. There are also other useful applications, e.g., in approximating characteristic functions. For instance, Corollaries 1 and 2 have been used in the proofs of Theorems 1 and 3, respectively; see (6.6) and (6.22).

Throughout this section, let $f: \mathbb{R} \to \mathbb{R}$ be a sufficiently smooth function.

Theorem 5. We have

$$\| \iint_{\mathbb{R}} f(u) d \mathbb{P} \{ \mathbb{S} \le u \} - \iint_{\mathbb{R}} f(u) dN_k(u) \| \ll C(\beta + \gamma),$$

where
$$C \stackrel{def}{=} ||f''|| + \dots + ||f^{(k+4)}||$$
.

Choosing $f(u) = \exp\{itu\}$ in Theorem 5, we obtain a useful inequality for the characteristic functions.

Corollary 1. We have

$$\left| \mathbb{E} \exp\{it\mathbb{S}\} - \int_{\mathbb{R}} \exp\{itu\} dN_k(u) \right| \ll (|t| + |t|^{k+4})(\beta + \gamma). \quad \Box$$

Theorem 5 and Corollary 1 can be extended to the case of the lower moment assumption $\mathbb{E} |\mathbb{T}|^{\alpha} < \infty$ with $\alpha < 2$. However, the technical details are more involved in Theorem 6 than those in Theorem 5.

Theorem 6. For $\alpha \in [1, 2)$, we have

$$\| \iint_{\mathbb{R}} f(u) d \mathbb{P} \{ \mathbb{S} \le u \} - \iint_{\mathbb{R}} f(u) dN_p(u) \| \ll C \left(\beta + \gamma^{(\alpha)} \right),$$

where $p = p(\alpha, k)$ is defined in Theorem 3, and C = ||f'|| + ||f'''|| for $\alpha = 1$, and $C = ||f'|| + \cdots + ||f^{(k+4)}||$ for $\alpha \in (1, 2)$.

Choosing $f(u) = \exp\{itu\}$ in Theorem 6, we get

Corollary 2. For $\alpha \in [1, 2)$, we have

$$\left| \mathbb{E} \exp\{it\mathbb{S}\} - \int_{\mathbb{R}} \exp\{itu\} dN_p(u) \right| \ll C(t)(\beta + \gamma^{(\alpha)})$$

where $p = p(\alpha, k)$ is defined in Theorem 3, and $C(t) = |t| + |t|^3$ for $\alpha = 1$, and $C(t) = |t| + |t|^{k+4}$ for $\alpha \in (1, 2)$.

5. Extensions. One may consider extending the work on the error bounds from U-statistics to more general class of statistics. Consider a non-linear statistic $\mathbb{W} = \mathbb{W}(X_1, ..., X_n)$, based on a sequence of r.v.'s $X_1, ..., X_n$. Examples include symmetric statistics, U-statistics with non-i.i.d. observations, L-statistics, studentized statistics, exchangeable statistics, finite population statistics, etc. Error bounds for symmetric statistics and nonsymmetric statistics were considered by van Zwet (1984) and Chen and Shao

(2007). Alberink (2000) studied error bounds for U-statistics with non-i.i.d. observations.

Suppose that we can linearize W into

$$\mathbb{W} = \mathbb{L} + \Delta$$
,

where \mathbb{L} is a linear statistic and Δ is an error term. One way to linearize \mathbb{W} is by the Hoeffding decomposition. We have seen how it works for U-statistics in this paper. For symmetric statistics, we refer to van Zwet (1984). The Hoeffding decompositions for independent observations and non-symmetric statistics was discussed by Karlin and Rinott (1982). For orthogonal decomposition of finite population statistics, see Bloznelis and Götze (2001). Hoeffding decompositions for exchangeable statistics was considered by Peccati (2004).

If \mathbb{L} is asymptotically normal and $\Delta \to 0$ in probability, \mathbb{W} is asymptotically normal. Using truncation and the Chebyshev inequality, we can easily get the following error bound:

$$||F_{\mathbb{W}} - \Phi|| \le ||F_{\mathbb{L}} - \Phi|| + (1+p)[(||\Phi'||)/p]^{p/(1+p)}(E|\Delta|^p)^{1/(1+p)}.$$

See (1.3) of Chen and Shao (2007). By taking p = 2, we get

(5.1)
$$||F_{\mathbb{W}} - \Phi|| \le ||F_{\mathbb{L}} - \Phi|| + 2(\operatorname{var} \Delta)^{1/3}.$$

For symmetric statistics, the next example shows that the exponent 1/3 in (5.1) is the best possible already. The example is similar to an example of Chen and Shao (2007).

Example 1. Take X, X_1, \ldots, X_n to be i.i.d. N(0,1) r.v.'s. Define

(5.2)
$$\mathbb{L} = n^{-1/2}(X_1 + \dots + X_n), \quad \Delta = -\varepsilon(|\mathbb{L}|^{-a} - \mathbb{E}|\mathbb{L}|^{-a}),$$

where 0 < a < 1/2 and $\varepsilon > 0$. Then, $\mathbb{W} = \mathbb{L} + \Delta$ is a symmetric statistic with finite variance since $\mathbb{E} \Delta^2 \le 2\varepsilon^2 \mathbb{E} |\mathbb{L}|^{-2a} < \infty$. Now if we have

$$(5.3) ||F_{\mathbb{W}} - \Phi|| \leq c||F_{\mathbb{L}} - \Phi|| + c(\operatorname{var}\Delta)^{\vartheta}$$

or

(5.4)
$$||F_{\mathbb{W}} - \Phi|| \le c \,\mathbb{E} \,|X|^3 / \sqrt{n} + c(\operatorname{var} \Delta)^{\vartheta},$$

then the exponent ϑ must satisfy $\vartheta \leq 1/3$.

Proof. Clearly, $\mathbb{L} \sim N(0,1)$. So $||F_{\mathbb{L}} - \Phi|| = \sup_x |\mathbb{P}\{\mathbb{L} \leq x\} - \Phi(x)| = 0$. Thus it suffices to show that the weaker inequality (5.4) implies $\vartheta \leq 1/3$. Also note that var $\Delta = c_1 \varepsilon^2$, where $c_1 = \text{var}(|\mathbb{L}|^{-a})$ is some positive constant depending only on a. It follows that $||F_{\mathbb{W}} - \Phi||$ and $\operatorname{var} \Delta$ in this example do not depend on n. Letting $n \to \infty$ in (5.4) yields

(5.5)
$$||F_{\mathbb{W}} - \Phi|| \le c(\operatorname{var} \Delta)^{\vartheta} = cc_1^{\vartheta} \varepsilon^{2\vartheta}.$$

On the other hand, for sufficiently small $\varepsilon > 0$, we have

$$||F_{\mathbb{W}} - \Phi|| = \sup_{x} |\mathbb{P}\{\mathbb{W} \le x\} - \Phi(x)|$$

$$\geq \mathbb{P}\{\mathbb{W} \le \varepsilon \mathbb{E} |\mathbb{L}|^{-a}\} - \mathbb{P}\{\mathbb{L} \le \varepsilon \mathbb{E} |\mathbb{L}|^{-a}\}$$

$$= \mathbb{P}\{\mathbb{L}|\mathbb{L}|^{a} \le \varepsilon\} - \mathbb{P}\{\mathbb{L} \le \varepsilon \mathbb{E} |\mathbb{L}|^{-a}\}$$

$$= \mathbb{P}\{0 \le \mathbb{L}^{a+1} \le \varepsilon\} - \mathbb{P}\{0 \le \mathbb{L} \le \varepsilon \mathbb{E} |\mathbb{L}|^{-a}\}$$

$$\geq c_{2}\varepsilon^{1/(a+1)} - c_{3}\varepsilon,$$
(5.6)

where c_2 and c_3 are positive constants depending only on a. The inequalities (5.5) and (5.6) imply $cc_1^{\vartheta}\varepsilon^{2\vartheta} \geq c_2\varepsilon^{1/(a+1)} - c_3\varepsilon$, i.e.,

$$(5.7) cc_1^{\vartheta} \varepsilon^{2\vartheta - 1/(a+1)} \ge c_2 - c_3 \varepsilon^{a/(a+1)}.$$

If $2\vartheta - 1/(a+1) > 0$, then letting $\varepsilon \downarrow 0$ in (5.7) would imply that $0 \ge c_2 > 0$, a contradiction. Hence we must have $2\vartheta - 1/(a+1) \le 0$ for all 0 < a < 1/2. Letting $a \uparrow 1/2$, we have $2\vartheta - 2/3 \le 0$, i.e., $\vartheta \le 1/3$.

6. Proofs of Theorems 1–6. We first prove several useful lemmas.

Lemma 1. We have

- (a) $1 \leq \sqrt{n} \beta$.
- (b) $n \mathbb{E} |L_1|^q \le \beta^{q-2} \text{ for } 2 \le q \le 3.$
- (c) $|\kappa_s| \leq \sqrt{\gamma_s} \leq \sqrt{\gamma}$ for s = 1, ..., k. (d) $|\kappa_s| \leq \beta^{\delta s} + \gamma_s^{(\alpha)} \leq \beta^{\delta s} + \gamma^{(\alpha)}$, where $3/2 < \alpha < 2$, $\delta = (2-\alpha)/(\alpha-1)$, and s is an integer such that $1 \le s < \delta^{-1} \wedge k$.

Proof. We will use Hölder's inequality: $\mathbb{E}|XY| \leq |(\mathbb{E}|X|^a)^{1/a}(\mathbb{E}|Y|^b)^{1/b}$ with a > 1 and 1/a + 1/b = 1. Recall that $\mathbb{EL}^2 = n \mathbb{EL}^2 = 1$.

- (a) $1 = (n n^{-1})^{3/2} = (n \mathbb{E} L_1^2)^{3/2} = n^{3/2} (\mathbb{E} L_1^2)^{3/2} \le n^{3/2} \mathbb{E} |L_1|^3 = \sqrt{n} \beta.$
- (b) $n \mathbb{E} |L_1|^q = n \mathbb{E} |L_1|^{2(3-q)} |L_1|^{3(q-2)} \le n (\mathbb{E} L_1^2)^{3-q} (\mathbb{E} |L_1|^3)^{q-2} = \beta^{q-2}.$
- (c) $|\kappa_s| \leq \binom{n}{s} \mathbb{E} |L_1 \dots L_s T_{1...s}| \leq \binom{n}{s} \left(\mathbb{E} L_1^2 \dots \mathbb{E} L_s^2 \mathbb{E} T_{1...s}^2 \right)^{1/2}$ = $\left(\binom{n}{s} / n^s \right)^{1/2} \gamma_s^{1/2} \leq \gamma_s^{1/2} \leq \gamma^{1/2}$.
- (d) We use Hölder's inequality with exponents $a = \alpha$ and $b = \alpha/(\alpha 1)$ and another basic inequality $x^{1/a}y^{1/b} \leq x + y$ for $x, y \geq 0$. Also note that $n \mathbb{E} |L_1|^q \leq \beta^{q-2}$ from (b), and $q-2=\delta$. We have

$$|\kappa_{s}| \leq {n \choose s} \mathbb{E} |L_{1} \dots L_{s} T_{1 \dots s}| \leq {n \choose s} (\mathbb{E} |L_{1}|^{q})^{s/q} (\mathbb{E} |T_{1 \dots s}|^{\alpha})^{1/\alpha}$$

$$\leq {n \choose s} (\mathbb{E} |L_{1}|^{q})^{s} + {n \choose s} \mathbb{E} |T_{1 \dots s}|^{\alpha} \leq (n \mathbb{E} |L_{1}|^{q})^{s} + \gamma_{s}^{(\alpha)}$$

$$\leq \beta^{(q-2)s} + \gamma_{s}^{(\alpha)} = \beta^{\delta s} + \gamma_{s}^{(\alpha)}. \quad \Box$$

Lemma 2. For sufficiently smooth functions $g: \mathbb{R} \to \mathbb{R}$, one has

Proof. The result follows from multiplying the obvious inequalities

$$||g(x+h) - g(x)|| \le 2||g||$$
 and $||g(x+h) - g(x)|| \le ||g'|||h|$,

and taking a square root, and then applying $\sqrt{2ab} \leq a + b$. \square

6.1. **Proof of Theorem 1.** Denote $\Delta \stackrel{\text{def}}{=} ||F_{\mathbb{S}} - N_k||$. We consider two separate cases:

Case I: $\max\{\beta, \gamma_1, ..., \gamma_k\} > c_k$;

Case II: $\max\{\beta, \gamma_1, ..., \gamma_k\} \leq c_k$,

where $c_k > 0$ is a constant depending only on k to be determined later.

Case I is relatively easy to prove. For Case II, we take a classical approach. That is, using the well-known Esseen's smoothing inequality, we reduce the problem to the estimation of some characteristic functions.

Proof for Case I: We shall prove that $\Delta \ll \beta + \gamma$, which clearly implies the desired $\Delta \ll \beta + \gamma \ln^2(\gamma)$. From Lemma 1, we have $\kappa_p^2 \leq \gamma_p$. Using this inequality, estimating $\mathbb{P} \{ \mathbb{S} \leq x \} \leq 1$ and $|\Phi^{(p)}(x)| \ll 1$, we derive

(6.2)
$$\Delta \ll 1 + |\kappa_1| + \dots + |\kappa_k| \ll 1 + \sqrt{\gamma_1} + \dots + \sqrt{\gamma_k} \ll 1 + \sqrt{\gamma}$$
.

We consider the alternative cases $\gamma \geq 1$ and $\gamma < 1$ separately. If $\gamma \geq 1$ then using (6.2) we have $\Delta \ll \sqrt{\gamma} \leq \gamma \leq \beta + \gamma$. If $\gamma < 1$ then (6.2) implies $\Delta \ll 1$. Using the condition: $\max\{\beta, \gamma_1, ..., \gamma_k\} > c_k$, we have

$$\Delta \ll 1 \le c_k^{-1} \max\{\beta, \gamma_1, ..., \gamma_k\} \le c_k^{-1} (\beta + \gamma_1 + \dots + \gamma_k) \ll \beta + \gamma.$$

Proof for Case II: We can assume that n is sufficiently large so that

$$n \geq 1/c_k$$
.

Indeed, for $n \leq 1/c_k$ the bound of Theorem 1 holds trivially. To see this, we note that, by (c) of Lemma 1, we have $\kappa_p^2 \leq \gamma_p$. Since $\max\{\beta, \gamma_1, ..., \gamma_k\} \leq c_k$ by assumption, we then have $\kappa_p^2 \leq \gamma_p \leq c_k \leq n^{-1} \leq 1$. Hence, it follows that

$$|N_k(x)| \ll 1 + |\kappa_1| + \dots + |\kappa_k| \le 1 + k \ll 1.$$

Therefore $\Delta \ll 1$. Using $1 \leq \sqrt{n}\beta$ (see Lemma 1), for small $n \leq 1/c_k$ we have $1 \ll \beta$. Inequalities $\Delta \ll 1$ and $1 \ll \beta$ imply $\Delta \ll \beta \ll \beta + \gamma$, which is somewhat better than the bound of Theorem 1.

By Lemma 1, $\kappa_p^2 \leq \gamma_p$, and condition: $\max\{\beta, \gamma_1, ..., \gamma_k\} \leq c_k$ implies that $\kappa_p^2 \ll 1$ for $1 \leq p \leq k$. Thus the function $N_k(x)$ has a bounded derivative,

(6.3)
$$|N'_k(x)| \ll 1 + |\kappa_1| + \dots + |\kappa_k| \ll 1.$$

Due to (6.3), to estimate Δ we can apply Esseen's smoothing inequality (see, for example, Feller (1971), Chapter XVI, Lemma 3.2): for any a > 0 we have

(6.4)
$$\Delta \ll \frac{1}{a} + \int_{-a}^{a} |t|^{-1} |f(t) - g(t)| dt,$$

where

(6.5)
$$f(t) = \mathbb{E} \exp\{it\mathbb{S}\}, \qquad g(t) = \int_{\mathbb{R}} \exp\{itx\} dN_k(x).$$

Note that $g(t) = (1 + \kappa_1(it)^3 + \dots + \kappa_k(it)^{k+1}) \exp\{-t^2/2\}$. We choose

$$a = \frac{\sqrt{c_k}}{\beta + \gamma}.$$

Since $\max\{\beta, \gamma_1, ..., \gamma_k\} \leq c_k$, we have $\beta \leq c_k$ and $\gamma \leq kc_k$, resulting in

$$a > \sqrt{c_k}/(c_k + \dots + c_k) = 1/((1+k)\sqrt{c_k}) > 1$$

if c_k is chosen small enough, e.g., $c_k \leq (1+k)^{-2}$.

Split the integral in (6.4) as $\int_{-a}^{a} = \int_{|t| < C} + \int_{C < |t| < a}$, where $C = C_k$ is a sufficiently large positive constant depending only on k to be chosen later.

To estimate $\int_{|t| < C}$ we use Corollary 1 of Section 4:

(6.6)
$$|f(t) - g(t)| \ll (|t| + |t|^{k+4})(\beta + \gamma).$$

It follows that $\int_{|t|<C} \ll \beta + \gamma$, a bound which is somewhat better than the desired bound with $\gamma \ln^2(3+1/\gamma)$ in place of γ .

It remains to consider the integral $\int_{C \leq |t| \leq a}$. Introduce the characteristic functions $\vartheta = \vartheta(t)$ and $\varrho = \varrho(t)$,

(6.7)
$$\vartheta = \mathbb{E} \exp\{itL_1\}, \quad \varrho = \exp\{-t^2/(2n)\}.$$

By Lemma 3 below, we have, for $4k \le m \le n/4$,

(6.8)
$$|f(t) - g(t)| \ll |t|^{k+4} (|\vartheta|^m + \varrho^m)(\beta + \gamma) + \gamma t^2 m/n$$
, for $|t| \ge 1$.

We shall use the well-known simple bound

(6.9)
$$|\vartheta| \le \exp\{-t^2/(4n)\}, \text{ for } |t| \le 1/\beta.$$

In particular, the inequality (6.9) holds for C < |t| < a.

We choose the integer number $4k \le m \le n$ as

$$m = \left[4(k+5)nt^{-2}\ln|t| \right], \quad C < |t| < a,$$

where [x] is the integer part of $x \in \mathbb{R}$. The number m = m(t) depends on t. If $C = C_k$ is sufficiently large, then m is a well defined integer such that $4k \le m \le n/4$, for sufficiently large n. Now (6.8) and (6.9) imply

$$(6.10) |f(t) - g(t)| \ll |t|^{-1}(\beta + \gamma) + \gamma \ln|t|, C < |t| < a.$$

Integrating (6.10) over C < |t| < a we derive $\int_{C \le |t| \le a} \ll \beta + \gamma + \gamma \ln^2 a$. To conclude the proof of the theorem, we note that $1 \le a \le 3 + 1/\gamma$ due to our choices of constants, and therefore $\gamma + \gamma \ln^2 a \ll \gamma \ln^2(\gamma)$.

We now prove the following lemma, which is used in the proof above. The lemma gives an expansion of the characteristic functions for $|t| \ge 1$.

Lemma 3. Assume that $\beta \leq 1$ and $\gamma \leq 1$. Then for $n \geq 4k$ and $4k \leq m \leq n/4$ we have

$$|f(t) - g(t)| \ll |t|^{k+4} (|\vartheta|^m + \varrho^m)(\beta + \gamma) + \gamma t^2 m/n, \quad \text{for } |t| \ge 1.$$

Proof. The proof follows from Propositions 1–3 below. When applying Propositions 1 and 2 one has to replace m by 2m.

Let us introduce some notation first. Let $\Omega_m = \{1, ..., m\}$ and let A be a subset of $\Omega_n = \{1, ..., n\}$, and use |A| to denote the number of elements in A. For convenience, we write $T_{i_1,...,i_k}$ or $T_{i_1...i_k}$ instead of $T_{\{i_1,...,i_k\}}$. Then we can write $\mathbb{T} = \mathbb{T}_1 + \cdots + \mathbb{T}_k$, where, for $1 \leq p \leq k$,

$$\mathbb{T}_p = \sum_{|A|=p} T_A$$
, with $T_A = t_p(X_j, j \in A)$.

We now split \mathbb{T} into two parts $\mathbb{T}^{(m)}$, $\mathbb{T}^{(0)}$ so that

$$\mathbb{T} = \mathbb{T}^{(m)} + \mathbb{T}^{(0)}, \quad \mathbb{T}^{(m)} = \mathbb{T}_1^{(m)} + \dots + \mathbb{T}_k^{(m)}, \quad \mathbb{T}^{(0)} = \mathbb{T}_1^{(0)} + \dots + \mathbb{T}_k^{(0)},$$

where

$$\mathbb{T}_s^{(m)} = \sum_{|A|=s, A \cap \Omega_m \neq \varnothing} T_A$$
, and $\mathbb{T}_s^{(0)} = \sum_{|A|=s, A \cap \Omega_m = \varnothing} T_A$.

We are now ready to prove Propositions 1-3.

Proposition 1. Let $1 \le m \le n$. The characteristic function

$$f = f(t) = \mathbb{E} \exp\{it\mathbb{S}\} \equiv \mathbb{E} \exp\{it(\mathbb{L} + \mathbb{T}^{(0)} + \mathbb{T}^{(m)})\}$$

satisfies $f = f_0 + f_1 + R$ with

(6.11)
$$f_0 = \mathbb{E} \exp\{it(\mathbb{L} + \mathbb{T}^{(0)})\}, \quad f_1 = (it) \mathbb{E} (\exp\{it(\mathbb{L} + \mathbb{T}^{(0)})\}\mathbb{T}^{(m)}),$$

and a remainder term R such that $|R| \ll \gamma t^2 m/n$.

Proof. We use $\tau, \tau_0, \tau_1, \tau_2, \ldots$ to denote i.i.d. r.v.'s which are uniformly distributed on [0, 1], and further assume that they are independent of all

other r.v.'s. Given a smooth function g, we will frequently use the Taylor expansion of the following form

$$g(x+h) = \sum_{s=0}^{k} g^{(s)}(x)h^{s}/s! + \mathbb{E}(1-\tau)^{k}g^{(k+1)}(x+\tau h)h^{k+1}/k!.$$

For instance, we can expand f in powers of $\mathbb{T}^{(m)}$, and obtain $f = f_0 + f_1 + R$ with

$$R = (it)^{2} \mathbb{E} (1 - \tau) \exp\{it(\mathbb{L} + \mathbb{T}^{(0)} + \tau \mathbb{T}^{(m)})\} (\mathbb{T}^{(m)})^{2}.$$

Thus,

$$|R| \le t^2 \mathbb{E} (\mathbb{T}^{(m)})^2 \ll t^2 \sum_{p=1}^k m n^{p-1} \mathbb{E} T_{1...p}^2 = t^2 \frac{m}{n} \sum_{p=1}^k n^p \mathbb{E} T_{1...p}^2 \ll \frac{m}{n} t^2 \gamma.$$

Proposition 2. Let $2k \leq m \leq n$. The function $f_0 + f_1$ with f_0 and f_1 defined by (6.11) satisfies $f_0 + f_1 = f_2 + f_3 + R_1$ with $f_2 = \mathbb{E} \exp\{it\mathbb{L}\}$, $f_3 = \mathbb{E}(it) \exp\{it\mathbb{L}\}\mathbb{T}$, and the remainder term R_1 satisfies $|R_1| \ll \gamma t^2 |\vartheta|^{m/2}$.

Proof. Since $\mathbb{T} = \mathbb{T}^{(0)} + \mathbb{T}^{(m)}$, it suffices to check that

(6.12)
$$f_0 \sim f_2 + (it) \mathbb{E} \exp\{it\mathbb{L}\} \mathbb{T}^{(0)}, \quad f_1 \sim (it) \mathbb{E} \exp\{it\mathbb{L}\} \mathbb{T}^{(m)}$$

with remainder terms bounded as R_1 .

Let us prove the first relation in (6.12). Note that

$$f_0 = \vartheta^m \mathbb{E} \exp\{it(\mathbb{L}^{(0)} + \mathbb{T}^{(0)})\}, \text{ where } \mathbb{L}^{(0)} = L_{m+1} + \dots + L_n$$

is the part of \mathbb{L} independent of X_1, \ldots, X_m . Now we can expand in powers of $\mathbb{T}^{(0)}$. We estimate the remainder term similar to the proof of Proposition 1. To estimate the variance of $\mathbb{T}^{(0)}$ we use the obvious inequality var $\mathbb{T}^{(0)} \leq \text{var } \mathbb{T}$.

Let us prove the second relation in (6.12). We consider only the case where m is an even integer. It suffices to check that

(6.13)
$$\mathbb{E} \exp\{it(\mathbb{L} + \mathbb{T}^{(0)})\}\mathbb{T}_p^{(m)} \sim \mathbb{E} \exp\{it\mathbb{L}\}\mathbb{T}_p^{(m)}, \text{ for } p = 1, \dots, k.$$

Let us show that

(6.14)
$$\mathbb{E} \exp\{it(\mathbb{L} + \mathbb{T}^{(0)})\}\mathbb{T}_{n}^{(m)} = \mathbb{E} \exp\{it(\mathbb{L} + \mathbb{T}^{(0)})\}\mathbb{T}_{n}^{*}$$

with $\mathbb{T}_p^* = \sum_{s=1}^p \sum_{(s)} \alpha_s T_A$ and $\alpha_s = \binom{m}{s} / \binom{m/2}{s}$, where $\sum_{(s)}$ is taken over all subsets $A \subset \Omega_n \setminus \Omega_{m/2}$ such that $|A \cap (\Omega_m \setminus \Omega_{m/2})| = s$.

To prove (6.14), let us start with a representation of $\mathbb{E} \exp\{it(\mathbb{L}+\mathbb{T}^{(0)})\}\mathbb{T}_p^{(m)}$. Sorting subsets A according to the cardinality of the intersection $A \cap \{1, \ldots, m\}$ and using the symmetry and i.i.d. assumptions, we get

$$\mathbb{E} \exp\{it(\mathbb{L} + \mathbb{T}^{(0)})\}\mathbb{T}_{p}^{(m)}$$

$$= \sum_{A: |A|=p, A\cap\Omega_{m}\neq\emptyset} \mathbb{E} T_{A} \exp\{it(\mathbb{L} + \mathbb{T}^{(0)})\}$$

$$= \sum_{s=1}^{p} \sum_{A: |A|=p, |A\cap\Omega_{m}|=s} \mathbb{E} T_{A} \exp\{it(\mathbb{L} + \mathbb{T}^{(0)})\}$$

$$(6.15) = \sum_{s=1}^{p} {m \choose s} \sum_{B: |B|=p-s, |B\cap\Omega_{m}|=\emptyset} \mathbb{E} T_{A_{s}\cup B} \exp\{it(\mathbb{L} + \mathbb{T}^{(0)})\},$$

where $A_1, \ldots, A_s \subset \{1, \ldots, m\}$ are arbitrary fixed subsets of cardinality s. For example, $A_s = \{1, \ldots, s\}$. A consideration similar to (6.15), starting with $\mathbb{E} \exp\{it(\mathbb{L} + \mathbb{T}^{(0)})\}$ \mathbb{T}_p^* instead of $\mathbb{E} \exp\{it(\mathbb{L} + \mathbb{T}^{(0)})\}$ $\mathbb{T}_p^{(m)}$, shows that $\mathbb{E} \exp\{it(\mathbb{L} + \mathbb{T}^{(0)})\}$ \mathbb{T}_p^* equals to the right hand side of (6.15), which proves the identity (6.14).

The statistic \mathbb{T}_p^* is independent of $X_1, \ldots, X_{m/2}$. Therefore (6.14) implies

(6.16)
$$\mathbb{E} \exp\{it(\mathbb{L} + \mathbb{T}^{(0)})\}\mathbb{T}_p^{(m)} = \vartheta^{m/2} \mathbb{E} \exp\{it(\mathbb{L}^{(m/2)} + \mathbb{T}^{(0)})\}\mathbb{T}_p^*.$$

Now we can expand in powers of $\mathbb{T}^{(0)}$. This leads to

$$\mathbb{E} \exp\{it(\mathbb{L} + \mathbb{T}^{(0)})\} \, \mathbb{T}_p^{(m)} \sim \vartheta^{m/2} \, \mathbb{E} \, \exp\{it\mathbb{L}^{(m/2)}\} \, \mathbb{T}_p^* \equiv \, \mathbb{E} \, \exp\{it\mathbb{L}\} \, \mathbb{T}^{(m)}$$

up to an error bounded by $t^2|\vartheta|^{m/2} \mathbb{E}|\mathbb{T}^{(0)}\mathbb{T}_p^*|$ (this can be checked as we did for (6.15)). Using the Hölder inequality, we have

$$\mathbb{E} |\mathbb{T}^{(0)} \mathbb{T}_p^*| \le (\operatorname{var} \mathbb{T}_p \operatorname{var} \mathbb{T}_p^*)^{1/2} \ll \operatorname{var} \mathbb{T} = \gamma,$$

since var $\mathbb{T}_p^* \ll \operatorname{var} \mathbb{T}_p \leq \operatorname{var} \mathbb{T}$. Combining the bounds, we get (6.13). \square

Proposition 3. Assume that $|t| \geq 1$. Let $\beta \leq 1$ and $\gamma \leq 1$. Let $n \geq 4k$ and $1 \leq m \leq n/4$. Then the function $f_4 = \mathbb{E} \exp\{it\mathbb{L}\} + (it)\mathbb{E} \exp\{it\mathbb{L}\}\mathbb{T}$ satisfies $f_4 = g + R_2$ with $g(t) = (1 + \kappa_1(it)^3 + \dots + \kappa_{k-1}(it)^{k+1})\varrho^n$ and the remainder term R_2 such that $|R_2| \ll (\beta + \gamma)|t|^{k+4}(\varrho^m + |\vartheta|^m)$.

Proof. Using the i.i.d. and symmetry assumptions, we can write $f_4 = \vartheta^n + it \sum_{p=1}^k \vartheta^{n-p} \binom{n}{p} D_p$, where

$$D_{p} = \mathbb{E} T_{1...p} \exp\{itL_{1}\} \dots \exp\{itL_{p}\}$$

= $(it)^{p} \mathbb{E} L_{1} \dots L_{p} T_{1...p} \exp\{it(\tau_{1}L_{1} + \dots + \tau_{p}L_{p})\}.$

In view of the inequalities $|\exp\{itu\} - 1| \ll \sqrt{|u|}$ and $\sqrt{u_1 + \cdots + u_p} \ll \sqrt{u_1 + \cdots + \sqrt{u_p}}$ for $u_s \ge 0$, and $\sqrt{|t|} \le |t|$ for $|t| \ge 1$, and the assumptions $n \mathbb{E} |L_1|^3 = \beta$ and $\mathbb{E} L_1^2 = 1/n$, we have

$$\left| \binom{n}{p} D_p - (it)^p \kappa_p \right| \leq |t|^p \binom{n}{p} \mathbb{E} |L_1 \dots L_p T_{1\dots p}| |\exp\{it(\sum_{j=1}^p \tau_j L_j)\} - 1|
\ll \binom{n}{p} |t|^{p+1/2} \mathbb{E} |L_1 \dots L_p T_{1\dots p}| (\sum_{j=1}^p \sqrt{|L_j|})
= p \binom{n}{p} |t|^{p+1/2} \mathbb{E} |L_1|^{3/2} |L_2 \dots L_p T_{1\dots p}|
\ll |t|^{k+1} \sqrt{\beta} n^{-p/2} \binom{n}{p} (\mathbb{E} T_{1\dots p}^2)^{1/2}
\ll |t|^{k+1} \sqrt{\beta} \sqrt{\gamma}
\leq |t|^{k+1} (\beta + \gamma).$$

This in turn leads to $f_4 = f_5 + R_3$ with $f_5 = \vartheta^n + \sum_{p=1}^k (it)^{p+1} \kappa_p \vartheta^{n-p}$ and the remainder term R_3 such that $|R_3| \ll |t|^{k+2} |\vartheta|^m (\beta + \gamma)$.

Recall $1 \leq \sqrt{n}\beta$. By a simple Taylor expansion, we have $\beta \leq 1$, we have $|\vartheta^p - 1| \ll t^2 \mathbb{E} L_1^2 = t^2/n \leq t^2\beta^2 \leq t^2\beta$. Hence $|\vartheta^{n-p} - \vartheta^n| \ll t^2\beta |\vartheta|^m$. Therefore, using $\kappa_p \leq \sqrt{\gamma} \leq 1$, we can write $f_5 = f_6 + R_4$ with

$$f_6 = \vartheta^n \left(1 + \sum_{p=1}^k (it)^{p+1} \kappa_p \right),$$

and a remainder term R_4 such that $|R_4| \ll |t|^{k+3} |\vartheta|^m (\beta + \gamma)$. A very standard calculation shows that

(6.17)
$$|\vartheta^n - \varrho^n| \ll |t|^3 (|\vartheta|^m + \varrho^m)\beta.$$

Thus, using again $\kappa_p \leq \sqrt{\gamma} \leq 1$, it follows that $f_6 = g(t) + R_5$ with $|R_5| \ll |t|^{k+4} (|\vartheta|^m + \varrho^m)\beta$. Collecting all the inequalities, we complete the proof. \square

6.2. **Proof of Theorem 3.** The proof is similar to that of Theorem 1. Recall that $\gamma_s^{(\alpha)} = \binom{n}{s} \mathbb{E} |T_{1...s}|^{\alpha}$. Without loss of generality we assume that

(6.18)
$$\beta \leq c_k$$
, and $\gamma_s^{(\alpha)} \leq c_k$, for $s = 1, \dots, k$;

and

$$(6.19) n \geq 1/c_k,$$

where c_k is a positive sufficiently small constant. A reduction leading to (6.18) and (6.19) is based on applications of $|\kappa_s| \leq \beta^{\delta s} + \gamma_s^{(\alpha)} \leq \beta^{\delta s} + \gamma^{(\alpha)}$ (see Lemma 1) instead of $\kappa_p^2 \leq \gamma_p$ used in the proof of Theorem 1. Using (6.18), (6.19) and (d) of Lemma 1, it is easy to check that the function N_p has a bounded derivative. Thus we can apply Esseen's smoothing inequality: for any a > 0,

(6.20)
$$||F_{\mathbb{S}} - N_p|| \ll \frac{1}{a} + \int_{-a}^{a} |t|^{-1} |f(t) - g^{(\alpha)}(t)| dt,$$

with f defined by (6.5) and

(6.21)
$$g^{(\alpha)}(t) = \int_{\mathbb{R}} \exp\{itx\} dN_p(x).$$

We choose $a = \sqrt{c_k}/(\beta + \gamma^{(\alpha)})$. Split the integral in (6.20) as $\int_{-a}^a = \int_{|t| < C} + \int_{C < |t| < a}$, where $C = C_k$ is a sufficiently large positive constant depending only on k and α to be chosen later.

To estimate $\int_{|t|< C}$ we use Corollary 2 of Section 4. Estimating $|t| \leq C$, the corollary implies

$$(6.22) |f(t) - g^{(\alpha)}(t)| \ll |t|^{\alpha} (\beta + \gamma^{(\alpha)}).$$

It follows that $\int_{|t|< C} \ll \beta + \gamma^{(\alpha)}$. Note that the presence of the factor $|t|^{\alpha}$ guaranties the convergence of the integral in a neighborhood of t = 0.

It remains to consider the integral $\int_{C \le |t| \le a}$. By Lemma 4 below, we have

$$|f(t) - g^{(\alpha)}(t)| \ll |t|^{k+4} (|\vartheta|^m + \varrho^m)(\beta + \gamma^{(\alpha)}) + \gamma^{(\alpha)}|t|^{\alpha} m/n,$$

for $|t| \ge 1$ and $4k \le m \le n/4$. We integrate this bound as in the proof of Theorem 1. A small difference arises since now instead of $\gamma |t|^2 m/n$ we have the

summand $\gamma^{(\alpha)}|t|^{\alpha}m/n$. The choice of $m \sim nt^{-2} \ln |t|$ leads to $\gamma^{(\alpha)}|t|^{\alpha}m/n \sim \gamma^{(\alpha)}|t|^{\alpha-2} \ln |t|$, which is an integrable function with respect to the measure dt/|t| at $|t| = \infty$, since we assume that $\alpha < 2$. As a consequence, we now no longer have any log factors. \square

We now prove the following lemma, which is used in the proof above. This lemma extends Lemma 3 to the case of lower moment assumption.

Lemma 4. Let $1 \le \alpha \le 2$. Assume $\beta \le 1$ and $\gamma \le 1$. Then for $n \ge 4k$ and $4k \le m \le n/4$ we have

$$|f(t)-g^{(\alpha)}(t)| \ll |t|^{k+4}(|\vartheta|^m+\varrho^m)(\beta+\gamma^{(\alpha)})+\gamma^{(\alpha)}|t|^\alpha m/n, \quad \textit{for } |t| \geq 1,$$

where ϑ , ϱ , f and $g^{(\alpha)}$ are given by (6.7), (6.5) and (6.21), respectively.

Proof. We consider the cases $\alpha = 1$ and $1 < \alpha \le 2$ separately. First consider the case $\alpha = 1$. Using $\exp\{itx\} = 1 + O(|tx|^{\alpha})$, the expansion of Proposition 1 is replaced by $f = f_0 + R$ with a remainder R such that $|R| \ll \gamma^{(\alpha)} |t|^{\alpha} m/n$. The function f_0 can be represented as

$$f_0 \equiv \mathbb{E} \exp\{it(\mathbb{L} + \mathbb{T}^{(0)})\} = \vartheta^m \mathbb{E} \exp\{it(\mathbb{L}^{(0)} + \mathbb{T}^{(0)})\}.$$

Thus by an expansion in powers of $\mathbb{T}^{(0)}$ we can replace it by $\mathbb{E} \exp\{it\mathbb{L}\} = \vartheta^n$. We conclude the proof replacing ϑ^n by $\varrho^n \equiv g \equiv g^{(\alpha)}$, see (6.17).

Next consider the case $1 < \alpha \le 2$. Using $\exp\{itx\} = 1 + itx + O(|tx|^{\alpha})$ and repeating arguments used in the proof of Proposition 1, we have $f = f_0 + f_1 + R$ with a remainder term R such that $|R| \ll \gamma^{(\alpha)}|t|^{\alpha}m/n$. The functions f_0, f_1 are the same as in Proposition 1. A repetition of the proof of Proposition 2 leads to $f_0 + f_1 = f_2 + f_3 + R_1$ with $|R_1| \ll t^2 \gamma^{(\alpha)} |\vartheta|^{m/2}$. A small difference in the proof is that now we have to use Taylor expansions and Hölder's inequalities adjusted to our lower moment assumption.

To conclude the proof we have to show that $f_2 + f_3 = g^{(\alpha)} + R_2$ with a remainder term R_2 such that $|R_2| \ll (\beta + \gamma^{(\alpha)})t^{k+4}(\varrho^m + |\vartheta|^m)$. The proof is similar to that of Proposition 3 provided that some adjustments related to the lower moment assumption are made. The flavor of the adjustments is like that used in the proof of Theorem 6. We omit formal exposition of indeed lengthy and technical details.

6.3. **Proof of Theorem 2.** We prove the first half now. If $\gamma \geq 1$, then we estimate $||F_{\mathbb{S}} - \Phi|| \leq 2$, and it follows that $||F_{\mathbb{S}} - \Phi|| \ll \beta + \gamma$. In the case $\gamma \leq 1$, we use the bound of Theorem 1. Since $|G_s| \ll |\kappa_s|$ and $\kappa_s^2 \leq \gamma$ from Lemma 1, we have

$$||F_{\mathbb{S}} - \Phi|| \le ||F_{\mathbb{S}} - N_k|| + \sum_{s=1}^k |G_s| \ll \beta + \gamma \ln^2(\gamma) + \sqrt{\gamma} \ll \beta + \sqrt{\gamma}.$$

Next, we will show the second half of the theorem with an example. Let X, X_1, \ldots, X_n be i.i.d. N(0, 1) r.v.'s., and define $\mathbb{S} = \mathbb{L} + \mathbb{T}$, where

$$\mathbb{L} = \frac{1}{\sqrt{n}} \sum_{i=1}^{n} X_i, \qquad \mathbb{T} = \frac{2\varepsilon}{(n-1)\sqrt{n}} \sum_{1 \le i \le j \le n} X_i X_j,$$

where $\varepsilon > 0$. Now we can show that, if we have

then we must have $\vartheta \leq 1/2$.

We prove the above claim by contradiction. Namely, assume that the contrary holds, i.e., that (6.23) holds with some $\vartheta > 1/2$. Clearly, $\beta = \mathbb{E} |X_1|^3/\sqrt{n} \ll n^{-1/2}$, $\gamma = \text{var } \mathbb{T} = O(n^{-1})$, $\kappa_1 = 0$, $\kappa_2 = \varepsilon n^{-1/2}$, $G_1(x) = 0$, and $G_1(x) = \varepsilon n^{-1/2} \Phi'''(x)$. In view of Theorem 1 and the assumption (6.23), we have

$$|G_2(x)| = ||G_2|| \le ||F_S - \Phi|| + ||F_S - \Phi - G_2|| \ll \beta + \gamma^{\vartheta} + \gamma \ln^2(\gamma)$$

Multiplying \sqrt{n} on both sides, we get

$$|\varepsilon\Phi'''(x)| \ll 1 + \sqrt{n}\gamma^{\vartheta} + \sqrt{n}\gamma \ln^2(\gamma).$$

Letting $n \to \infty$, and in view of $\gamma = O(n^{-1})$ and $\vartheta > 1/2$, we get $\varepsilon |\Phi''(x)| \ll 1$, which contradicts the assumption that $\varepsilon > 0$ is an arbitrary positive number. Thus we must have $\vartheta \leq 1/2$.

6.4. **Proof of Theorem 5.** Let η be a r.v. from N(0,1). Note that the left hand side of the inequality in Theorem 5 can be written as

$$\mathbb{E} f(\mathbb{S}) = \int_{\mathbb{R}} f(u) d\mathbb{P} \{ \mathbb{S} \le u \},$$

$$\mathbb{E} f(\eta) + \kappa_1 \mathbb{E} f''(\eta) + \dots + \kappa_k \mathbb{E} f^{(k+1)}(\eta) = \int_{\mathbb{R}} f(u) dN_k(u).$$

Expanding $f(\mathbb{S}) = f(\mathbb{L} + \mathbb{T})$ in powers of \mathbb{T} , we get

$$\mathbb{E} f(\mathbb{S}) = \mathbb{E} f(\mathbb{L}) + \mathbb{E} f'(\mathbb{L})\mathbb{T} + \vartheta, \quad \vartheta \stackrel{\text{def}}{=} \mathbb{E} (1 - \tau) f''(\mathbb{L} + \tau \mathbb{T})\mathbb{T}^2,$$

where τ is uniformly distributed on [0, 1], and independent of all other r.v.'s. To prove the theorem it suffices to check that

(6.24)
$$\|\mathbb{E} f(\mathbb{L}) - \mathbb{E} f(\eta)\| \ll \|f'''\|\beta,$$

(6.25)
$$\|\mathbb{E}\mathbb{T}f'(\mathbb{L}) - I\| \ll C(\beta + \gamma),$$

where

(6.27)
$$I = \kappa_1 \mathbb{E} f''(\eta) + \dots + \kappa_k \mathbb{E} f^{(k+1)}(\eta) \stackrel{\text{def}}{=} I_1 + \dots + I_k.$$

The estimate (6.24) is well known since $\mathbb{L} = L_1 + \cdots + L_n$ is a sum of i.i.d. r.v.'s, see for example Bentkus *et al.* (2000).

The bound (6.26) is obvious. Indeed, we have $\|\theta\| \le \|f''\| \mathbb{E} \mathbb{T}^2 = \|f''\| \gamma$. It remains to prove (6.25). Using $\mathbb{T} = \mathbb{T}_1 + \cdots + \mathbb{T}_k$, the linear structure of \mathbb{T}_s , and the i.i.d. assumption, we can write

(6.28)
$$\mathbb{E} \,\mathbb{T} \,f'(\mathbb{L}) = J_1 + \dots + J_k, \quad J_s \stackrel{\text{def}}{=} \binom{n}{s} \,\mathbb{E} \,T_{1\dots s} f'(\mathbb{L}).$$

In view of (6.27) and (6.28), the proof of (6.25) reduces to checking that

Let us split $\mathbb{L} = W + R$, where $W = L_1 + \cdots + L_s$ and $R = L_{s+1} + \cdots + L_n$. Writing $K_s = \kappa_s \mathbb{E} f^{(s+1)}(R)$, instead of (6.29) it suffices to prove that

$$(6.30) ||I_s - K_s|| \ll C(\beta + \gamma),$$

and

$$(6.31) $||K_s - J_s|| \ll C(\beta + \gamma).$$$

Proof of (6.30). Note that $I_s - K_s = \kappa_s \left(\mathbb{E} f^{(s+1)}(\eta) - \mathbb{E} f^{(s+1)}(R) \right)$. Since $\mathbb{L} = R + W$, to prove (6.30) it suffices to show that

(6.32)
$$|\kappa_s| \|\mathbb{E} f^{(s+1)}(\eta) - \mathbb{E} f^{(s+1)}(\mathbb{L})\| \ll C(\beta + \gamma),$$

and

$$(6.33) \qquad |\kappa_s| \parallel \mathbb{E} f^{(s+1)}(R+W) - \mathbb{E} f^{(s+1)}(R) \parallel \ll C(\beta+\gamma).$$

Let us prove (6.32). We consider the cases $\beta \geq 1$ and $\beta < 1$ separately. Using $\kappa_s^2 \leq \gamma$ (see Lemma 1), in the case $\beta \geq 1$ we have

$$|\kappa_s| \|\mathbb{E} f^{(s+1)}(\eta) - \mathbb{E} f^{(s+1)}(\mathbb{L})\| \le 2 \|f^{(s+1)}\| \sqrt{\gamma} \ll C \sqrt{\gamma\beta} \le C (\gamma + \beta).$$

In the case $\beta \leq 1$ we use the bound (6.24) replacing f by $f^{(s+1)}$. Since $\kappa_s^2 \leq \gamma$, we get

$$|\kappa_s| \|\mathbb{E} f^{(s+1)}(\eta) - \mathbb{E} f^{(s+1)}(\mathbb{L})\| \ll \|f^{(s+4)}\| \sqrt{\gamma} \beta \le C \sqrt{\gamma\beta} \le C (\gamma + \beta).$$

Let us prove (6.33). We apply (6.1) replacing g by $f^{(s+1)}$, the variable x by R, and h by W respectively. Since $\kappa_s^2 \leq \gamma$, we have

(6.34)
$$|\kappa_s| \| \mathbb{E} f^{(s+1)}(R+W) - \mathbb{E} f^{(s+1)}(R) \|$$

$$\ll (\|f^{(s+1)}\| + \|f^{(s+2)}\|) \gamma^{1/2} \mathbb{E} |W|^{1/2}.$$

Using $|W|^{1/2} \leq |L_1|^{1/2} + \cdots + |L_s|^{1/2}$ and $\mathbb{E} L_m^2 = 1/n$, we derive $\mathbb{E} |W|^{1/2} \ll n^{-1/4} \leq \beta^{1/2}$. In view of (6.34), an application of $\sqrt{\gamma\beta} \leq \gamma + \beta$ yields (6.33). Proof of (6.31). Using short Taylor expansions, we can represent J_s as

(6.35)
$$J_s = \binom{n}{s} \mathbb{E} L_1 \dots L_s T_{1\dots s} f^{(s+1)} (V + R)$$

with $V = \tau_1 L_1 + \cdots + \tau_s L_s$. Representation (6.35) can be proved in s steps. Let us consider details related only to the first step. The degeneracy property of the kernels implies that

(6.36)
$$\mathbb{E} T_{1...s} f'(L_2 + \dots + L_n) = 0$$

since the conditional expectation $\mathbb{E}(T_{1...s} | X_2, ..., X_{s+1}) = 0$, and $L_2 + \cdots + L_n$ is independent of X_1 . Using (6.36), expanding in powers of L_1 we get

$$J_s = \binom{n}{s} \mathbb{E} L_1 T_{1...s} f''(\tau_1 L_1 + L_2 + \dots + L_n).$$

Proceeding in a similar way with L_2, \ldots, L_s in place of L_1 , we arrive at (6.35)

Using the definition of κ_s , and the expression (6.35) for J_s , we can write $K_s - J_s = \binom{n}{s} \mathbb{E} L_1 \dots L_s T_{1\dots s} \left(f^{(s+1)}(R) - f^{(s+1)}(V+R) \right)$. By an application of (6.1) with $g = f^{(s+1)}$, x = R and h = V, it follows that

(6.37)
$$||K_s - J_s|| \leq C\binom{n}{s} \mathbb{E} |L_1 \dots L_s T_{1\dots s}| |V|^{1/2}.$$

Using the inequality $|V|^{1/2} \leq |L_1|^{1/2} + \cdots + |L_s|^{1/2}$, the Hölder inequality, and the assumption $\mathbb{E}|L_1|^3 = \beta/n$ and $\mathbb{E}L_1^2 = 1/n$, we get

$$\mathbb{E} |L_1 \dots L_s T_{1...s}| |V|^{1/2} \leq s \mathbb{E} |L_1|^{3/2} |L_2 \dots L_s T_{1...s}|$$

$$\ll (\mathbb{E} |L_1|^3 L_2^2 \dots L_s^2 \mathbb{E} T_{1...s}^2)^{1/2}$$

$$= n^{-s/2} \left(\beta \mathbb{E} T_{1...s}^2\right)^{1/2}.$$

$$(6.38)$$

Using $\mathbb{E} T_{1...s}^2 = \mathbb{E} \mathbb{T}_s^2/\binom{n}{s} \leq \gamma/\binom{n}{s}$, relations (6.37) and (6.38) yield $||K_s - J_s|| \ll C\sqrt{\alpha}\sqrt{\beta\gamma}$, $\alpha = \binom{n}{s}/n^s$. Noting that $\alpha \leq 1$ and $\sqrt{\beta\gamma} \leq \beta + \gamma$, we derive $||K_s - J_s|| \ll C(\beta + \gamma)$, which concludes the proof of (6.31).

6.5. **Proof of Theorem 6.** Let η be a r.v. from N(0,1). Note that the left hand side of the inequality in Theorem 6 can be written as

$$\mathbb{E} f(\mathbb{S}) = \int_{\mathbb{R}} f(u) d\mathbb{P} \{ \mathbb{S} \le u \},$$

$$\mathbb{E} f(\eta) + \kappa_1 \mathbb{E} f''(\eta) + \dots + \kappa_p \mathbb{E} f^{(p+1)}(\eta) = \int_{\mathbb{R}} f(u) dN_p(u).$$

We consider the cases $\alpha = 1$ and $1 < \alpha < 2$, separately.

First consider the case $\alpha=1.$ In this case $I^{(\alpha)}=0.$ Similarly to (6.1) one can check that

(6.39)
$$f(\mathbb{S}) = f(\mathbb{L}) + \vartheta$$
, with $\|\vartheta\| \ll \|f\|^{1-\alpha} \|f'\|^{\alpha} \|\mathbb{T}\|^{\alpha}$.

It is easy to show that

$$(6.40) \quad \mathbb{E} |\mathbb{T}_p|^{\alpha} \ll \gamma_p^{(\alpha)} = \binom{n}{p} \ \mathbb{E} |T_{1\dots p}|^{\alpha}, \quad \mathbb{E} |\mathbb{T}|^{\alpha} \ll \gamma^{(\alpha)} = \gamma_1^{(\alpha)} + \dots + \gamma_k^{(\alpha)}.$$

Then the proof follows from (6.39), (6.24), and (6.40).

Now consider the case $1 < \alpha < 2$. Similar to (6.1), we have

$$f(\mathbb{S}) = f(\mathbb{L}) + f'(\mathbb{L})\mathbb{T} + \vartheta, \text{ with } \|\vartheta\| \ll (\|f'\| + \|f''\|)|\mathbb{T}|^{\alpha}.$$

From (6.40), we get $\mathbb{E} |\mathbb{T}|^{\alpha} \ll \gamma^{(\alpha)}$. Also from (6.24) it follows easily that $\|\mathbb{E} f(\mathbb{L}) - \mathbb{E} f(\eta)\| \ll \|f'''\| \beta$. It remains to show that

(6.41)
$$\|\mathbb{E}\,\mathbb{T}\,f'(\mathbb{L}) - I^{(\alpha)}\| \ll C(\beta + \gamma^{(\alpha)}),$$

where $I^{(\alpha)} = \kappa_1 \mathbb{E} f''(\eta) + \cdots + \kappa_p \mathbb{E} f^{(p+1)}(\eta)$. We have

(6.42)
$$\mathbb{E} \mathbb{T} f'(\mathbb{L}) = J_1 + \dots + J_k, \quad J_s \stackrel{\text{def}}{=} \binom{n}{s} \mathbb{E} T_{1\dots s} f'(\mathbb{L}).$$

In view of (6.41) and (6.42), the proof of the theorem reduces to checking

$$(6.43) ||I_s - J_s|| \ll C(\beta + \gamma^{(\alpha)}), \text{for } 1 \le s \le p,$$

with $I_s = \kappa_s \mathbb{E} f^{(s+1)}(\eta)$, and

(6.44)
$$||J_s|| \ll C(\beta + \gamma^{(\alpha)}), \text{ for } p < s \le k,$$

where p is an integer satisfying the condition of the theorem.

Estimation of J_s for s > p. We have to prove (6.44). Consider the difference operator δ_m such that $\delta_m f(x) = f(x + L_m) - f(x)$. Arguments similar to those used to derive (6.35), allow to write

$$(6.45) J_s = \binom{n}{s} \mathbb{E} T_{1...s} \delta_1 \dots \delta_s f'(R)$$

with $R = L_{s+1} + \cdots + L_n$. Similar to (6.1), we have

$$\|\delta_1 \dots \delta_s f'(R)\| \ll (\|f'\| + \|f^{(s+1)}\| |L_1|^{\varepsilon} \dots |L_s|^{\varepsilon}, \quad 0 < \varepsilon < 1.$$

From this and the Hölder inequality with exponents $q = \alpha/(\alpha - 1)$ and α , so that $1/q + 1/\alpha = 1$, we have

$$||J_s|| \ll C\binom{n}{s} \mathbb{E} |L_1|^{\varepsilon} \dots |L_s|^{\varepsilon} |T_{1\dots s}|$$

$$\ll C\binom{n}{s} (\mathbb{E} |L_1|^{\varepsilon q} \dots |L_s|^{\varepsilon q})^{1/q} (\mathbb{E} |T_{1\dots s}|^{\alpha})^{1/\alpha}$$

$$= C\left(\binom{n}{s} (\mathbb{E} |L_1|^{\varepsilon q})^s\right)^{1/q} (\gamma^{(\alpha)})^{1/\alpha}$$

$$\ll C\binom{n}{s} (\mathbb{E} |L_1|^{\varepsilon q})^s + C\gamma^{(\alpha)}.$$

To complete the proof of (6.44) we need to prove $\binom{n}{s} \left(\mathbb{E} |L_1|^{\varepsilon q} \right)^s \ll \beta$. Choose $\varepsilon = (2+1/s)/q$. The condition s > p guaranties that $\varepsilon \leq 1$. Using $n \mathbb{E} |L_1|^{2+1/s} \leq \beta^{1/s}$ (see (b) of Lemma 1), we have $\binom{n}{s} \left(\mathbb{E} |L_1|^{\varepsilon q} \right)^s \leq (n \mathbb{E} |L_1|^{2+1/s})^s \leq \beta$, which concludes the proof for s > p.

Estimation of J_s for $1 \le s \le p$. We have to prove (6.43). There is something to prove only if $3/2 < \alpha < 2$. Indeed, in the case $1 < \alpha \le 3/2$, the integer p from the condition of the theorem satisfies $p \le 0$, and we have no J_s to estimate.

Let us split $\mathbb{L} = W + R$, where $W = L_1 + \cdots + L_s$ and $R = L_{s+1} + \cdots + L_n$. Writing $K_s = \kappa_s \mathbb{E} f^{(s+1)}(R)$ instead of (6.43) it suffices to prove that $||I_s - K_s||$ and $||K_s - J_s||$ are bounded from above like $||I_s - J_s||$ in (6.43). Estimation of $||I_s - K_s||$. Note that $I_s - K_s = \kappa_s \mathbb{E} (f^{(s+1)}(\eta) - f^{(s+1)}(R))$. Since $\mathbb{L} = R + W$, it suffices to show that

$$(6.46) |\kappa_s| \left\| f^{(s+1)}(\eta) - f^{(s+1)}(\mathbb{L}) \right\| \ll C(\beta + \gamma^{(\alpha)}),$$

and

$$(6.47) \quad |\kappa_s| \left\| \mathbb{E} f^{(s+1)}(R+W) - \mathbb{E} f^{(s+1)}(R) \right\| \ll C(\beta + \gamma^{(\alpha)}).$$

Let us prove (6.46). We consider the cases $\beta \geq 1$ and $\beta \leq 1$ separately. Using $|\kappa_s| \ll \beta^{\delta s} + \gamma^{(\alpha)}$ and $\delta s \leq 1$ (see (d) of Lemma 1), in the case $\beta \geq 1$ we have

$$|\kappa_s| \left\| \mathbb{E} f^{(s+1)}(\eta) - \mathbb{E} f^{(s+1)}(\mathbb{L}) \right\| \le 2 \left\| f^{(s+1)} \right\| (\beta^{\delta s} + \gamma^{(\alpha)}) \ll C \left(\beta + \gamma^{(\alpha)}\right).$$

In the case $\beta \leq 1$ we combine (d) of Lemma 1 with the bound (6.24) replacing f by $f^{(s+1)}$. We get

$$|\kappa_s| \left\| \mathbb{E} f^{(s+1)}(\eta) - \mathbb{E} f^{(s+1)}(\mathbb{L}) \right\| \ll \|f^{(s+4)}\| (\beta^{\delta s} + \gamma^{(\alpha)})\beta \le C (\beta + \gamma^{(\alpha)}).$$

Let us now prove (6.47). Again we consider the cases $\beta \geq 1$ and $\beta \leq 1$ separately. We omit details in the case $\beta \geq 1$ since they are very similar to those in the proof of (6.46). In the case $\beta \leq 1$ we use (d) of Lemma 1 and apply $||f^{(s+1)}(R+W) - \mathbb{E} f^{(s+1)}(R)|| \leq C||W||$ to get

$$(6.48) |\kappa_s| \|\mathbb{E} f^{(s+1)}(R+W) - \mathbb{E} f^{(s+1)}(R)\| \ll C (\beta^{\delta s} + \gamma^{(\alpha)}) \mathbb{E} |W|.$$

Using $|W| \le |L_1| + \dots + |L_s|$ and $\mathbb{E} L_m^2 = 1/n$, we derive $\mathbb{E} |W| \ll n^{-1/2} \le \beta$. In view of our assumption $\beta \le 1$, the inequality (6.48) yields (6.47).

Estimation of $||K_s - J_s||$. By (6.35) we have

$$J_s = \binom{n}{s} \mathbb{E} L_1 \dots L_s T_{1\dots s} f^{(s+1)} (V + R)$$

with $V = \tau_1 L_1 + \cdots + \tau_s L_s$. This representation, combined with the definition κ_s , allows to write

$$K_s - J_s = \binom{n}{s} \mathbb{E} L_1 \dots L_s T_{1\dots s} \left(f^{(s+1)}(R) - f^{(s+1)}(V+R) \right).$$

Again we consider the cases $\beta \geq 1$ and $\beta \leq 1$ separately. We omit details in the case $\beta \geq 1$ since they are very similar to those in the proof of (6.46). In the case $\beta \leq 1$ we apply

$$\|\mathbb{E} f^{(s+1)}(R) - f^{(s+1)}(R+V)\| \le C\|V\|^{\varepsilon}, \quad 0 \le \varepsilon \le 1.$$

From the inequality $|V|^{\varepsilon} \ll |L_1|^{\varepsilon} + \cdots + |L_s|^{\varepsilon}$, the i.i.d. assumption, the Hölder inequality with exponents $q = \alpha/(\alpha - 1)$ and α , and also $\binom{n}{s} \leq n^s$, we have

$$||K_{s} - J_{s}|| \ll C\binom{n}{s} \mathbb{E} |L_{1} \dots L_{s} T_{1\dots s}| |V|^{\varepsilon}$$

$$\leq Cn^{s} s \mathbb{E} |L_{1}|^{1+\varepsilon} |L_{2} \dots L_{s} T_{1\dots s}|$$

$$\ll Cn^{s} \left(\mathbb{E} |L_{1}|^{(1+\varepsilon)q} \left(\mathbb{E} |L_{1}|^{q} \right)^{s-1} \right)^{1/q} \left(\mathbb{E} |T_{1\dots s}|^{\alpha} \right)^{1/\alpha}$$

$$\ll Cn^{s} \mathbb{E} |L_{1}|^{(1+\varepsilon)q} \left(\mathbb{E} |L_{1}|^{q} \right)^{s-1} + Cn^{s} \mathbb{E} |T_{1\dots s}|^{\alpha}$$

$$= Cn^{s} \mathbb{E} |L_{1}|^{(1+\varepsilon)q} \left(\mathbb{E} |L_{1}|^{q} \right)^{s-1} + C\gamma^{(\alpha)}.$$

We choose ε so that $(1+\varepsilon)q=3$. In view of $3/2<\alpha<2$, the number ε satisfies $0<\varepsilon<1$. Since $n\mathbb{E}\,|L_1|^3=\beta$, to conclude the estimation of $\|K_s-J_s\|$ it suffices to verify the inequality $n\mathbb{E}\,|L_1|^q\leq 1$. Recalling that $\beta\leq 1$ and applying Lemma 1, we have $n\mathbb{E}\,|L_1|^q\leq \beta^{q-2}\leq 1$, which completes the estimation of $\|K_s-J_s\|$. \square

Acknowledgements. The authors would like to thank the Editor, Associate Editor and referees for their comments and criticisms, which lead to this much improved version.

REFERENCES

- [1] Alberink, I. B. (2000). A Berry-Esseen bound for *U*-statistics in the non-i.i.d. case. *J. Theoret. Probab.* **13** 519-533.
- [2] Alberink, I. B. and Bentkus, V. (1999). Bounds for the concentration of asymptotically normal statistics. *Acta Applic. Math.* **58** 11-59.
- [3] Alberink, I. B. and Bentkus, V. (2001a). Lyapunov type bounds for *U*-statistics. *Theory Probab. Appl.* **46** 571-588.

- [4] Alberink, I. B. and Bentkus, V. (2001b). Berry-Esseen bounds for von Mises and U-statistics. Lithuanian Math. J. 41 1-16.
- [5] Bentkus, V., Götze, F., Paulauskas, V. and Račkauskas, A. (2000). The accuracy of Gaussian approximation in Banach spaces. Limit Theorems of Probability Theory, (Yu.V. Prokhorov, V.A. Statulevičius, eds.), 25-111, Springer Verlag, Berlin.
- [6] Bentkus, V., Götze, F. and van Zwet, W. R. (1997). An Edgeworth expansion for symmetric statistics. Ann. Statist. 25 851-896.
- [7] Bloznelis, M. and Gotze, F. (2001). Orthogonal decomposition of finite population statistics and its applications to distributional asymptotics. Ann. Statist. 29 353-365.
- [8] Borovskikh, Yu. V. (2000). On normal approximations to U-statistics. Probab. Theory and Appl. 45 406-423.
- [9] Borovskikh, Yu. V. and Korolyuk, V. S. (1985). Approximation of non-degenerate U-statistics. Theory Probab. Appl. 30 439-450.
- [10] Borovskikh, Yu. V. and Korolyuk, V. S. (1994). Theory of U-statistics. Kluwer Academic Publishers, Dordrecht.
- [11] Callaert, H. and Janssen, P. (1978). The Berry-Esseen theorem for U-statistics. Ann. Statist. 6 417-421.
- [12] Chen, L. H. Y. and Shao, Q. M. (2007). Normal approximation for nonlinear statistics using a concentration inequality approach. Bernoulli 13 581–599.
- [13] Feller, W. (1971). An Introduction to Probability Theory and Its Applications. Vol. II. 2nd Ed., Wiley, New York.
- [14] Friedrich, K. O. (1989). A Berry-Esseen bound for functions of independent random variables. Ann. Statist. 17 170-183.
- [15] Gine, E., Latala, R. and Zinn, J. (2000). Exponential and moment inequalities for U-statistics. Progress in Probability 47 13-38.
- [16] Hoeffding, W. (1948). A class of statistics with asymptotically normal distributions. Ann. Math. Stat. 19 293-325.
- [17] Jing, B.-Y.; Wang, Q.Y. (2003). Edgeworth expansion for U-statistics under minimal conditions. Ann. Statist. 31 1376-1391.
- [18] Karlin, S. and Rinott, Y. (1982). Applications of ANOVA type decompositions for comparisons of conditional variance statistics including jackknife estimates. Ann. Statist. 10 485-501.
- [19] Peccati, G. (2004). Hoeffding-ANOVA decompositions for symmetric statistics of exchangeable observations. Ann. Prob. 32 1796-1829.
- van Zwet, W. R. (1984). A Berry-Esseen bound for symmetric statistics. Z. Wahrsch. verw. Gebiete. 66 425-440.

Akademijos 4 VILNIUS, LITHUANIA E-MAIL: bentkus@ktl.mii.lt

Department of Mathematics INSTITUTE OF MATHEMATICS AND INFORMATICS HONG KONG UNIVERSITY OF SCIENCE AND TECHNOLOGY CLEAR WATER BAY, KOWLON, HONG KONG E-MAIL: majing@ust.hk

Department of Statistics and Applied Probability 6 Science Drive 2 National University of Singapore

Singapore 117546

E-MAIL: stazw@nus.edu.sg