

# CENTRAL LIMIT THEOREM FOR FOURIER TRANSFORMS OF STATIONARY PROCESSES\*

BY MAGDA PELIGRAD, WEI BIAOWU

*University of Cincinnati and University of Chicago*

We consider asymptotic behavior of Fourier transforms of stationary ergodic sequences with finite second moments. We establish a central limit theorem (CLT) for almost all frequencies and also an annealed CLT. The theorems hold for all regular sequences. Our results shed new light on the foundation of spectral analysis and on the asymptotic distribution of periodogram, and it provides a nice blend of harmonic analysis, theory of stationary processes and theory of martingales.

**1. Introduction.** In frequency or spectral domain analysis of time series, periodograms play a fundamental role. Since its introduction by Schuster in 1898, periodograms have been used in almost all scientific fields. Given a realization  $(X_j)_{j=1}^n$  of a stochastic process  $(X_j)_{j \in \mathbb{Z}}$ , the periodogram is defined as

$$I_n(\theta) = \frac{1}{2\pi n} \left| \sum_{j=1}^n X_j \exp(ji\theta) \right|^2, \quad \theta \in \mathbb{R},$$

where  $i = \sqrt{-1}$  is the imaginary unit. Periodogram is the building block in spectral domain analysis and a distributional theory is clearly needed in the related statistical inference. If  $(X_j)$  is a Gaussian process, then the Fourier transform

$$S_n(\theta) = \sum_{j=1}^n X_j \exp(ji\theta)$$

is complex Gaussian. Fisher (1929) proposed a test for hidden periodicities and obtained a distributional theory based on i.i.d. Gaussian random variables. If  $(X_j)$  is not Gaussian, the distribution of  $S_n(\theta)$  typically does not have a close form and one needs to resort to asymptotics. It is well-known

---

\* Magda Peligrad's research is supported in part by a Charles Phelps Taft Memorial Fund grant and NSA grant H98230-09-1-0005. Wei Biao Wu's research is supported in part by a Charles Phelps Taft Memorial Fund and the National Science Foundation DMS-0478704

*AMS 2000 subject classifications:* Primary 60F05; secondary 60F17

*Keywords and phrases:* Fourier transform, spectral analysis, martingale, central limit theorem, stationary process

since Wiener and Wintner (1941) (see also Lacey and Terwilleger, 2008) that for any stationary sequence  $(X_j)_{j \in \mathbb{Z}}$  in  $\mathcal{L}^1$  (namely  $E|X_0| < \infty$ ) there is a set  $\Omega'$  of probability 1 such that for all  $\theta$  and  $\omega \in \Omega'$ ,  $S_n(\theta)/n$  converges. Our problem is to investigate the speed of this convergence by providing a central limit theorem for the real and imaginary parts of  $S_n(\theta)/\sqrt{n}$ .

The above central limit problem was considered by many authors under various dependence conditions. We mention Rosenblatt (Theorem 5.3, p 131, 1985) who considered mixing processes; Brockwell and Davis (Theorem 10.3.2., p 347, 1991), Walker (1965) and Terrin and Hurvich (1994) discussed linear processes and Wu (2005) treated mixingales. Other contributions can be found in Olshen (1967), Rootzen (1976), Yajima (1989), Woodroffe (1992), Walker (2000), Lahiri (2003) and Liu and Lin (2009) among others.

To establish an asymptotic theory for  $S_n(\theta)/\sqrt{n}$ , we shall provide the framework of stationary processes that can be introduced in several equivalent ways. We assume that  $(\xi_n)_{n \in \mathbb{Z}}$  is a stationary ergodic Markov chain defined on a probability space  $(\Omega, \mathcal{F}, P)$  with values in a measurable space. The marginal distribution is denoted by  $\pi(A) = P(\xi_0 \in A)$ . Next let  $\mathcal{L}_0^2(\pi)$  be the set of functions such that  $\int h^2 d\pi < \infty$  and  $\int h d\pi = 0$ . Denote by  $\mathcal{F}_k$  the  $\sigma$ -field generated by  $\xi_j$  with  $j \leq k$ ,  $X_j = h(\xi_j)$ . For any integrable random variable  $X$  we denote  $E_k(X) = E(X|\mathcal{F}_k)$ . We assume  $h \in \mathcal{L}_0^2(\pi)$ , in other words we assume  $\|X_0\| := (E|X_0^2|)^{1/2} < \infty$  and  $E(X_0) = 0$ . Notice that any stationary sequence  $(Y_k)_{k \in \mathbb{Z}}$  can be viewed as a function of a Markov process  $\xi_k = (Y_j; j \leq k)$  with the function  $g(\xi_k) = Y_k$ .

The stationary stochastic processes may be also introduced in the following alternative way. Let  $T : \Omega \mapsto \Omega$  be a bijective bi-measurable transformation preserving the probability. Let  $\mathcal{F}_0$  be a  $\sigma$ -algebra of  $\mathcal{F}$  satisfying  $\mathcal{F}_1 \subseteq T^{-1}(\mathcal{F}_0)$ . We then define the nondecreasing filtration  $(\mathcal{F}_j)_{j \in \mathbb{Z}}$  by  $\mathcal{F}_j = T^{-j}(\mathcal{F}_0)$  (referred to as the stationary filtration). Let  $X_0$  be a random variable which is  $\mathcal{F}_0$ -measurable. We also define the stationary sequence  $(X_j)_{j \in \mathbb{Z}}$  by  $X_j = X_0 \circ T^j$ . In this paper we shall use both frameworks.

The rest of the paper is structured as follows. The main results are presented in Section 2 and proved in Section 4. Our proofs in Section 4 provide an interesting blend of harmonic analysis, martingale approximation and theory of stationary processes. Examples of regular processes and further extensions are given in Section 3.

**2. Main Results.** We shall assume that the following regularity condition holds:

$$(2.1) \quad E(X_0|\mathcal{F}_{-\infty}) = 0 \text{ } P\text{-almost surely}$$

and also that the sequence is stationary and ergodic. The regularity condition is quite mild and it is satisfied for many popular processes used in practice. Section 3 provides examples of stationary ergodic processes for which (2.1) holds.

We shall present first a central limit theorem for almost all frequencies. In Theorem 2.1, we let the parameter  $\theta$  be in the space  $[0, 2\pi]$ , endowed with Borelian sigma algebra and Lebesgue measure  $\lambda$ . We denote by " $\Rightarrow$ " the weak convergence, or convergence in distribution.

**THEOREM 2.1.** *Let  $(X_k)_{k \in \mathbb{Z}}$  be a stationary ergodic process such that (2.1) is satisfied. Then for almost all  $\theta \in (0, 2\pi)$ , the following convergence holds:*

$$(2.2) \quad \lim_{n \rightarrow \infty} \frac{E|S_n(\theta)|^2}{n} = g(\theta) \text{ (say),}$$

where  $g$  is integrable over  $\theta \in [0, 2\pi]$ , and

$$(2.3) \quad \frac{1}{\sqrt{n}}[\operatorname{Re}(S_n(\theta)), \operatorname{Im}(S_n(\theta))] \Rightarrow [N_1(\theta), N_2(\theta)] \text{ under } P,$$

where  $N_1(\theta)$  and  $N_2(\theta)$  are independent identically distributed normal random variables mean 0 and variance  $g(\theta)/2$ .

As implied by Lemma 4.2 in Section 4,  $g(\theta)/(2\pi)$  is actually the spectral density associated with the spectral distribution function induced by the covariances

$$(2.4) \quad c_j = \operatorname{cov}(X_0, X_j), \quad j \in \mathbb{Z}.$$

More specifically, by Herglotz's Theorem (Brockwell and Davis, 1991), there exists a non-decreasing function  $G$  (the spectral distribution function) on  $[0, 2\pi]$  such that, for all  $j \in \mathbb{Z}$ ,

$$(2.5) \quad c_j = \int_0^{2\pi} \exp(ij\theta) dG(\theta).$$

Hence, by Lemma 4.2,  $G$  is absolutely continuous and the spectral density  $G'(\theta)$  equals to  $g(\theta)/(2\pi)$  almost surely. By (2.5) or (4.3),  $\int_0^{2\pi} g(\theta) d\theta = 2\pi c_0$ . So a nice implication of our results is that, under the regularity condition, we obtain an interesting representation of the spectral densities; see Lemma 4.2 for details.

Following the proof of Theorem 2.1, by the Cramer-Wold device, for

$$V_n(\omega, \theta) = \frac{S_n}{\sqrt{n}}(\omega, \theta),$$

we have that, for almost all pairs  $(\theta', \theta'')$  (Lebesgue),  $V_n(\omega, \theta')$  and  $V_n(\omega, \theta'')$  are asymptotically independent. In this sense Theorem 2.1 justifies the folklore in the spectral domain analysis of time series: the Fourier transforms of stationary processes are asymptotically independent Gaussian. Namely, in the spectral or Fourier domain, the Fourier-transformed processes are asymptotically independent, while the original process can be very strongly dependent; see Example 3.3.

Theorem 2.1 substantially improves the result in Wu (2005) that proves (2.3) under the following stronger condition

$$(2.6) \quad \sum_{n=1}^{\infty} \frac{\|E(X_n|\mathcal{F}_0)\|^2}{n} < \infty.$$

We shall also establish the following characterization of the annealed CLT. Let  $Id_2$  denote the identity  $2 \times 2$  matrix.

**THEOREM 2.2.** *Under the same conditions as in Theorem 2.1 on the product space  $([0, 2\pi] \times \Omega, \mathcal{B} \times \mathcal{F}, \lambda \times P)$  we have*

$$(2.7) \quad \frac{1}{\sqrt{n}}[\operatorname{Re}(S_n(\theta)), \operatorname{Im}(S_n(\theta))] \Rightarrow [g(U)/2]^{1/2}N(0, Id_2) \text{ under } \lambda \times P.$$

Here  $U$  is a random variable independent of  $N(0, Id_2)$  and uniformly distributed on  $[0, 2\pi]$  and  $g(\cdot)$  is defined by (2.2).

Two types of stochastic processes can be considered concerning the partial sum  $S_n(\theta)$ . The process  $V_n(\omega, \theta)$  indexed by  $\theta$  is asymptotically Gaussian white noise. For another version, we consider

$$W_n(t, \omega, \theta) = \frac{S_{[nt]}(\omega, \theta)}{\sqrt{n}}, \quad 0 \leq t \leq 1,$$

where  $[x] = \max\{k \in \mathbb{Z} : k \leq x\}$  is the integer part of  $x$ . We shall prove the following invariance principle:

**PROPOSITION 2.1.** *Assume that  $(X_k)$  is stationary ergodic and satisfies (2.1). Then  $W_n(t, \omega, \theta)$  is tight in  $D(0, 1)$ , and*

$$[\operatorname{Re}(W_n(t, \omega, \theta)), \operatorname{Im}(W_n(t, \omega, \theta))] \Rightarrow [g(U)/2]^{1/2}[W'(t), W''(t)] \text{ under } \lambda \times P,$$

where  $(W'(t), W''(t))$  are two independent standard Brownian motions independent of  $U$ , and  $U$  is a random variable uniformly distributed on  $[0, 2\pi]$ .

We now give some remarks and discussions.

REMARK 2.1. (Non-adapted case) Our CLT also holds if  $X_0$  is not  $\mathcal{F}_0$ -measurable. Then clearly the regularity condition we shall impose is

$$X_0 \text{ is } \mathcal{F}_\infty\text{-measurable and } E(X_0|\mathcal{F}_\infty) = 0 \text{ almost surely.}$$

REMARK 2.2. (Adapted non-regular case) For general adapted sequences our CLT result still holds under centering. Let  $\tilde{S}_n(\theta) = S_n(\theta) - E(S_n(\theta)|\mathcal{F}_\infty)$  and  $\tilde{X}_k = X_k - E(X_k|\mathcal{F}_\infty)$ , where  $E(X_k|\mathcal{F}_\infty)$  denotes the following limit that holds almost surely and in  $\mathcal{L}^2$ :

$$\lim_{n \rightarrow \infty} E(X_k|\mathcal{F}_{-n}) = E(X_k|\mathcal{F}_\infty).$$

Then, for almost all  $\theta \in (0, 2\pi)$ , we have

$$\frac{1}{\sqrt{n}}[\operatorname{Re}(\tilde{S}_n(\theta)), \operatorname{Im}(\tilde{S}_n(\theta))] \Rightarrow N(0, \frac{g(\theta)}{2} Id_2) \text{ under } P,$$

by applying Theorem 2.1 to the stationary sequence  $\tilde{X}_k$ . Therefore the conclusion of Theorem 2.1 holds if we replace the assumption of regularity (2.1) by: for  $\lambda$ -almost all  $\theta$

$$\frac{1}{\sqrt{n}}E(S_n(\theta)|\mathcal{F}_\infty) \rightarrow 0 \text{ in probability.}$$

Now, since  $\|E(S_n(\theta)|\mathcal{F}_\infty)\|_2 \leq \|E(S_n(\theta)|\mathcal{F}_{-n})\|_2 \leq \|E(S_n(\theta)|\mathcal{F}_0)\|_2$ , Theorem 2.1 still holds if

$$\frac{1}{\sqrt{n}}\|E(S_n(\theta)|\mathcal{F}_{-n})\|_2 \rightarrow 0 \text{ as } n \rightarrow \infty$$

or under the condition

$$(2.8) \quad \frac{1}{\sqrt{n}}\|E(S_n(\theta)|\mathcal{F}_0)\|_2 \rightarrow 0 \text{ as } n \rightarrow \infty.$$

REMARK 2.3. (Conditional CLT) Since we use in the proof martingale approximation actually our CLT is a conditional CLT, that allows for a random change of measure. See Hall and Heyde (1980) and Dedecker and Merlevède (2002).

REMARK 2.4. (Periodogram) We notice that, as a consequence of Theorem 2.1, for sequences satisfying (2.1) the periodogram  $n^{-1}|S_n(\theta)|^2$  is asymptotically distributed as  $\frac{g(\theta)}{2}\chi^2(2)$  for almost all frequencies.

REMARK 2.5. (Resulting identities) By using the invariance principle in Proposition 2.1 we can get the convergence of many interesting functionals of  $S_n(\theta)$  and periodograms. As a consequence of Proposition 2.1 we can get for instance

$$\begin{aligned} \frac{1}{\|X_0\|^2 n \pi} \int_0^{2\pi} E \left[ \max_{1 \leq m \leq n} \sum_{k=1}^m X_k \cos(k\theta) \right]^2 d\theta &\rightarrow E \left| \sup_{0 \leq t \leq 1} W(t) \right|^2 \\ &= \int_0^\infty 2[1 - \Phi(\sqrt{y})] dy = 1 \end{aligned}$$

by noting that  $P(\sup_{0 \leq t \leq 1} W(t) \geq u) = 2P(W(1) \geq u)$  for  $u \geq 0$ . Here  $\Phi(\cdot)$  is the standard Gaussian distribution function.

**3. Examples.** Here we present several examples of processes for which the conclusions of Theorems 2.1 and 2.2 hold.

Clearly condition (2.1) is satisfied if the left tail sigma field  $\mathcal{F}_{-\infty}$  is trivial. These processes are called regular (see Chapter 2 Volume 1 in Bradley, 2007). Notice however that our condition (2.1) refers rather to the function  $X_0 = f(\xi_0)$  in relation to the tail field  $\mathcal{F}_\infty$ .

EXAMPLE 3.1. (Mixing sequences) We shall introduce the following mixing coefficients: For any two  $\sigma$ -algebras  $\mathcal{A}$  and  $\mathcal{B}$  define the strong mixing coefficient

$$\alpha(\mathcal{A}, \mathcal{B}) = \sup\{|P(A \cap B) - P(A)P(B)| : A \in \mathcal{A}, B \in \mathcal{B}\}$$

and the  $\rho$ -mixing coefficient, also known as maximal coefficient of correlation

$$\rho(\mathcal{A}, \mathcal{B}) = \sup\{\text{Cov}(X, Y) / \|X\|_2 \|Y\|_2 : X \in \mathcal{L}^2(\mathcal{A}), Y \in \mathcal{L}^2(\mathcal{B})\}.$$

For the stationary sequence of random variables  $(X_k)_{k \in \mathbb{Z}}$ ,  $\mathcal{F}^n$  denotes the  $\sigma$ -field generated by  $X_i$  with indices  $i \geq n$ , and  $\mathcal{F}_m$  denotes the  $\sigma$ -field generated by  $X_i$  with indices  $i \leq m$ . The sequences of coefficients  $\alpha(n)$  and  $\rho(n)$  are then defined by

$$\alpha(n) = \alpha(\mathcal{F}_0, \mathcal{F}^n) \quad \text{and} \quad \rho(n) = \rho(\mathcal{F}_0, \mathcal{F}^n),$$

respectively. For strongly mixing sequences, namely the strong mixing coefficients  $\alpha(n) \rightarrow 0$ , the tail sigma field is trivial; see Claim 2.17a in Bradley (2007). Examples of this type include Harris recurrent Markov chains. If  $\lim_{n \rightarrow \infty} \rho(n) < 1$ , then the tail sigma field is also trivial; see Proposition (5.6) in Bradley (2007).

EXAMPLE 3.2. (Functions of Gaussian processes.) Assume  $(Y_k)$  is a stationary Gaussian sequence and define  $X_n = f(Y_k, k \leq n)$ . Let  $f$  be such that  $E(X_0) = 0$  and  $E(X_0^2) < \infty$ . Since any Gaussian sequence can be represented as a function of i.i.d random variables, the process is then regular. Rosenblatt (1981) considered Fourier transforms of functionals of Gaussian sequences.

EXAMPLE 3.3. (Functions of i.i.d random variables) Let  $\varepsilon_k$  be i.i.d and consider  $X_n = f(\varepsilon_k, k \leq n)$ . These are regular processes and therefore Theorems 2.1 and 2.2 are applicable. Examples include linear processes, functions of linear processes and iterated random functions (Wu and Woodroffe, 2000) among others. For example let  $X_n = \sum_{j=0}^{\infty} a_j \varepsilon_{n-j}$ , where  $\varepsilon_j$  are i.i.d with mean 0 and variance 1 and  $a_j$  are real coefficients with  $\sum_{j=1}^{\infty} a_j^2 < \infty$ . In this case  $X_n$  is well-defined and, by Lemmas 4.1 and 4.2, the spectral density is  $g(\theta)/(2\pi)$ , where

$$g(\theta) = \left| \sum_{j=0}^{\infty} a_j \exp(ij\theta) \right|^2.$$

As a specific example, let  $a_j = j^{-1/2}/\log j$ ,  $j \geq 2$ , and  $a_0 = a_1 = 1$ . By elementary manipulations, the covariances  $c_j \sim (\log j)^{-1}$ , which decays very slowly as  $j \rightarrow \infty$ , hence suggesting strong dependence. For this example, condition (2.6) is violated. By the Tauberian theorem, as  $\theta \rightarrow 0$ ,  $g(\theta) \sim \pi/(|\theta| \log^2 |\theta|)$ , which has a pole at  $\theta = 0$ .

EXAMPLE 3.4. (Reversible Markov chains) As before, let  $\xi_j$  be a stationary ergodic Markov chain with values in a measurable space. We use the notations and constructions from the introduction. The marginal distribution and the transition metric are denoted by  $\pi(A) = P(\xi_0 \in A)$  and  $Q(\xi_0, A) = P(\xi_1 \in A | \xi_0)$ . In addition  $Q$  denotes the operator  $Qf(\xi) = \int f(z)Q(\xi, dz)$ . Let  $Q^*$  be the adjoint operator of the restriction of  $Q$  to  $\mathcal{L}^2(\pi)$  and assume  $Q = Q^*$ . Then, for any  $f \in \mathcal{L}_0^2(\pi)$  the central limit theorem of Theorem 2.1 holds. To see this we shall verify condition (2.8). By spectral calculus

$$\|E(S_n(\theta)|\mathcal{F}_0)\|^2 = \int_{-1}^1 \left| \sum_{k=1}^n (t \exp(i\theta))^k \right|^2 \rho_f(dt),$$

where  $\rho_f$  denotes the spectral measure of  $f$  with respect to  $Q$  (see for instance Borodin and Ibragimov (1994) for this identity). For  $\theta \neq 0$ ,  $\pi$  and  $-1 \leq t \leq 1$ , we have

$$\left| \sum_{k=1}^n (t \exp(i\theta))^k \right|^2 \leq 4|1 - \exp(i\theta)t|^{-2}$$

$$= 4(1 + t^2 - 2t \cos \theta)^{-1} \leq 4(1 - (\cos \theta)^2)^{-1}.$$

Therefore, for  $\lambda$ -almost all  $\theta$

$$\frac{1}{n} \|E(S_n(\theta)|\mathcal{F}_0)\|^2 \rightarrow 0.$$

**4. Proofs.** We shall establish first some preparatory lemmas. The almost sure convergence in Lemma 4.1 was shown by Wu (2005). The convergence in  $\mathcal{L}^2$  is new here. For  $k \in \mathbb{Z}$  we define the projection operator by

$$(4.1) \quad \mathcal{P}_k \cdot = E(\cdot|\mathcal{F}_k) - E(\cdot|\mathcal{F}_{k-1}).$$

LEMMA 4.1. *Let*

$$T_n(\theta) = \sum_{j=0}^n X_j \exp(ji\theta) = X_0 + S_n(\theta).$$

*Under (2.1), for  $\lambda$ -almost all  $\theta$  (Lebesgue), we have*

$$(4.2) \quad \mathcal{P}_0 T_n(\theta) \rightarrow \sum_{l=0}^{\infty} \mathcal{P}_0 X_l \exp(li\theta) =: D_0(\theta) \quad P\text{-almost surely and in } \mathcal{L}^2.$$

*Proof.* By (2.1),  $\sum_{k \in \mathbb{Z}} \|\mathcal{P}_k X_0\|_2^2 = \|X_0\|_2^2 < \infty$ , we have  $\sum_{k \in \mathbb{Z}} |\mathcal{P}_0 X_k|^2 < \infty$   $P$ -almost surely. Therefore by Carleson's (1966) theorem, for almost all  $\omega$ ,  $\sum_{1 \leq k \leq n} (\mathcal{P}_0 X_k) \exp(ik\theta)$  converges  $\lambda$ -almost surely, where  $\lambda$  is the Lebesgue measure on  $[0, 2\pi]$ . Denote the limit by  $D_0 = D_0(\theta)$ . We now consider the set

$$A = \{(\theta, \omega) \in [0, 2\pi] \times \Omega, \text{ where } \{\mathcal{P}_0 S_n(\theta)\}_n \text{ does not converge}\}$$

and notice that almost all sections for  $\omega$  fixed have Lebesgue measure 0. So by Fubini's theorem the set  $A$  has measure 0 in the product space and therefore, again by Fubini's theorem, almost all sections for  $\theta$  fixed have probability 0. It follows that for almost all  $\theta$ ,  $\mathcal{P}_0(S_n(\theta)) \rightarrow D_0$  almost surely under  $P$ . Next, by the maximal inequality in Hunt and Young (1974), there is a constant  $C$  such that

$$\int_0^{2\pi} \left[ \sup_n |\mathcal{P}_0(S_n(\theta))|^2 \right] \lambda(d\theta) \leq C \sum_k |\mathcal{P}_0 X_k|_2^2$$

and then we integrate

$$\int_0^{2\pi} E \left[ \sup_n |\mathcal{P}_0(S_n(\theta))|^2 \right] \lambda(d\theta) \leq C \|X_0\|_2^2 < \infty.$$

Therefore

$$E \left[ \sup_n |\mathcal{P}_0(S_n(\theta))|^2 \right] < \infty \text{ for almost all } \theta.$$

Since  $|\mathcal{P}_0 S_n(\theta)| < \sup_n |\mathcal{P}_0 S_n(\theta)|$ , and the last one is integrable for almost all  $\theta$ , by the Lebesgue dominated convergence we have that  $\mathcal{P}_0(S_n(\theta))$  converges in  $\mathcal{L}^2$ .  $\square$

LEMMA 4.2. *Let  $g(\theta) = E|D_0(\theta)|^2$ . For all  $j \in \mathbb{Z}$ , we have*

$$(4.3) \quad \int_0^{2\pi} g(\theta) \exp(ji\theta) d\theta = 2\pi c_j,$$

where  $c_j$  are defined by (2.4). So  $(c_j)$  are the Fourier coefficients of  $g$ . Additionally, for almost all  $\theta$ ,

$$(4.4) \quad \lim_{n \rightarrow \infty} \frac{E|S_n(\theta)|^2}{n} = g(\theta).$$

*Proof.* Without loss of generality we let  $j \geq 0$  and  $j < n$ . As before, let

$$\mathcal{P}_0 T_n(\theta) = \sum_{l=0}^n \mathcal{P}_0 X_l \exp(li\theta).$$

By elementary trigonometric identities, we have

$$(4.5) \quad \frac{1}{2\pi} \int_0^{2\pi} |\mathcal{P}_0 T_n(\theta)|^2 \exp(ji\theta) d\theta = \sum_{l=0}^{n-j} (\mathcal{P}_0 X_l)(\mathcal{P}_0 X_{l+j}).$$

Since  $X_j = \sum_{l \in \mathbb{Z}} \mathcal{P}_l X_j$ , by orthogonality of martingale differences and stationarity we have that

$$\begin{aligned} c_j &= \lim_{N \rightarrow \infty} E \left[ \left( \sum_{l=-N}^0 \mathcal{P}_l X_0 \right) \left( \sum_{l=-N}^0 \mathcal{P}_l X_j \right) \right] \\ &= \lim_{N \rightarrow \infty} \sum_{l=-N}^0 E[(\mathcal{P}_l X_0)(\mathcal{P}_l X_j)] \\ &= \lim_{n \rightarrow \infty} \sum_{l=0}^{n-j} E[(\mathcal{P}_0 X_l)(\mathcal{P}_0 X_{l+j})]. \end{aligned}$$

By (4.5) and the Lebesgue dominated convergence theorem, as in the proof of Lemma 4.1, (4.3) follows in view of Hunt's maximal inequality since  $\sup_n |\mathcal{P}_0 T_n(\theta)|$  is integrable.

Now we prove (4.4). By stationarity, we have

$$\begin{aligned}
\frac{1}{n}E|S_n(\theta)|^2 &= \frac{1}{n} \sum_{j=1}^n \sum_{l=1}^n E(X_j X_l) \exp(ij\theta) \exp(-il\theta) \\
&= \frac{1}{n} \sum_{j=1}^{n-1} \sum_{l=1}^n c_{j-l} \exp((j-l)i\theta) \\
(4.6) \qquad &= \sum_{j=-(n-1)}^{n-1} \left(1 - \frac{|j|}{n}\right) c_j \exp(ij\theta).
\end{aligned}$$

Namely  $E|S_n(\theta)|^2/n$  is the Cesaro average of the sum  $\sum_{j=-l}^l c_j \exp(ij\theta)$ . Note that  $g(\theta) = \|D_0(\theta)\|^2$  is integrable over  $[0, 2\pi]$ . Therefore by the Fejér-Lebesgue Theorem (cf Bary, 1964, p. 139 or Theorem 15.7 in Champeney, 1989), (4.4) holds for  $\lambda$ -almost all  $\theta \in [0, 2\pi]$  (Lebesgue).  $\square$

REMARK 4.1. In the proof of Lemma 4.2, (4.6) implies that the sequence  $(c_j \exp(ij\theta))$  is Cesaro summable. It turns out that, generally speaking,  $\sum_{j=0}^{\infty} c_j \exp(ij\theta)$  may not exist for almost all  $\theta$ . Consider the example in Kolmogorov (1923) (see also Theorem 3.1, p. 305, in Zygmund, 2002): there exists a sequence of non-negative trigonometric polynomials  $f_n$  with constant term  $1/2$ , a sequence of positive integers  $q_k \rightarrow \infty$ , and a positive sequence  $A_n \rightarrow \infty$ , such that the function

$$g(x) = \sum_{k=1}^{\infty} \frac{f_{n_k}(q_k x)}{A_{n_k}^{1/2}}$$

is integrable. However, for almost all  $\theta$ , the Fourier sum  $\sum_{l=1}^{\infty} c_l \exp(li\theta)$  diverges, where  $c_l = \int_0^{2\pi} g(\theta) \exp(li\theta) d\theta$  is the Fourier coefficient of  $g$ . Let  $G(x) = \int_0^x g(u) du$ . By Herglotz's Theorem (Brockwell and Davis, 1991), there exists a stationary process  $(X_j)$  such that its spectral distribution function is  $G$  and its covariance function is  $c_l$ .

LEMMA 4.3. *Assume (2.1). On the product space  $([0, 2\pi] \times \Omega, \mathcal{B} \times \mathcal{F}, \lambda \times P)$  we have that*

$$\left(\frac{\max_{1 \leq k \leq n} |S_k(\theta)|^2}{n}\right)_{n \geq 1} \text{ is uniformly integrable.}$$

*Proof.* Let  $m$  be a positive integer. We shall decompose the partial sums in a sum of  $m$  martingales and a remainder in the following way:

$$\frac{S_k(\theta)}{\sqrt{n}} = \frac{1}{\sqrt{n}} \sum_{l=1}^k \exp(il\theta) \sum_{j=0}^{m-1} \mathcal{P}_{l-j}(X_l) + \frac{1}{\sqrt{n}} \sum_{l=1}^k \exp(il\theta) E_{l-m}(X_l).$$

Notice that for any  $0 \leq j \leq m - 1$ ,

$$\sum_{l=1}^k \exp(il\theta) \mathcal{P}_{l-j}(X_l)$$

is a martingale adapted to the filtration  $(\mathcal{B} \times \mathcal{F}_k)$ . Moreover, since  $(X_k)_{k \in \mathbb{Z}}$  is a stationary sequence with variables square integrable it follows that  $(X_k^2)_{k \in \mathbb{Z}}$  is a uniformly integrable sequence. This fact implies that for  $j$  fixed the sequence  $(\mathcal{P}_{k-j}(X_k))_{k \in \mathbb{Z}}$  is also uniformly integrable. It follows that  $\sum_{l=1}^k \exp(il\theta) \mathcal{P}_{l-j}(X_l)$  is a martingale with uniformly integrable differences under the measure  $\lambda \times P$ . It is known that for a martingale with uniformly integrable differences we have

$$\frac{1}{n} \max_{1 \leq k \leq n} \left| \sum_{l=1}^k \exp(il\theta) \mathcal{P}_{l-j}(X_l) \right|^2$$

is uniformly integrable (see for instance Dedecker and Rio (2000), Proposition 1).

The result follows since by Hunt and Young (1974) maximal inequality

$$\int_0^{2\pi} \max_{1 \leq k \leq n} \left| \sum_{l=1}^k \exp(il\theta) E_{l-m}(X_l) \right|^2 d\theta \leq C \sum_{l=1}^n |E_{l-m}(X_l)|^2$$

and therefore, denoting by  $\mathbf{E}$  the expected value with respect to  $\lambda \times P$ , we have by regularity condition (2.1) that

$$\begin{aligned} \frac{1}{n} \mathbf{E} \left[ \max_{1 \leq k \leq n} \left| \sum_{l=1}^k \exp(il\theta) E_{l-m}(X_l) \right|^2 \right] &\leq \frac{C}{n} \sum_{l=1}^n E |E_{l-m}(X_l)|^2 \\ &= CE |E_{-m}(X_0)|^2 \rightarrow 0 \end{aligned}$$

as  $m \rightarrow \infty$  uniformly in  $n$ . So, the uniform integrability follows.  $\square$

4.1. *Proof of Theorem 2.1.* The first assertion of Theorem 2.1 is just Lemma 4.1. We now prove (2.3).

**Step 1. The construction of martingale.**

Define the projector operator by (4.1). Then we construct as in Lemma 4.1

$$\mathcal{P}_1(S_n(\theta)) = E(S_n(\theta)|\mathcal{F}_1) - E(S_n(\theta)|\mathcal{F}_0) = \sum_{k=1}^n \exp(ik\theta) \mathcal{P}_1(X_k)$$

and then by Lemma 4.1 for almost all  $\theta$

$$\mathcal{P}_1(S_n(\theta)) \rightarrow D_1(\theta) \text{ in } \mathcal{L}^2.$$

To verify it is a martingale we start from

$$E(\mathcal{P}_1(S_n(\theta))|\mathcal{F}_0) = 0 \text{ almost surely under } P$$

and by the contractive property of the conditional expectation

$$0 = E(\mathcal{P}_1(S_n(\theta))|\mathcal{F}_0) \rightarrow E(D_1(\theta)|\mathcal{F}_0) \text{ in } \mathcal{L}^2.$$

We then construct the sequence of stationary martingale differences  $(D_k(\theta))_{k \geq 1}$ , given by

$$\mathcal{P}_k(S_{n+k}(\theta) - S_k(\theta)) \rightarrow D_k(\theta) \text{ in } \mathcal{L}^2.$$

### Step 2. Martingale approximation.

Denote by

$$M_n(\theta) = \sum_{1 \leq k \leq n} \exp(ik\theta) D_k(\theta).$$

We show that, for almost all  $\theta$ ,

$$(4.7) \quad \frac{E|S_n(\theta) - M_n(\theta)|^2}{n} \rightarrow 0.$$

To this end, note that  $S_n(\theta) - E(S_n(\theta)|\mathcal{F}_0)$  and  $E(S_n(\theta)|\mathcal{F}_0)$  are orthogonal, we have

$$(4.8) \quad \|S_n(\theta)\|^2 = \|S_n(\theta) - E(S_n(\theta)|\mathcal{F}_0)\|^2 + \|E(S_n(\theta)|\mathcal{F}_0)\|^2.$$

For those  $\theta$  such that (4.2) holds, we have, by the orthogonality of martingale differences and the stationarity, that

$$(4.9) \quad \begin{aligned} \|S_n(\theta) - E(S_n(\theta)|\mathcal{F}_0) - M_n(\theta)\|^2 &= \sum_{k=1}^n \|\mathcal{P}_k(S_n(\theta) - M_n(\theta))\|^2 = \\ &= \sum_{k=1}^n \|\mathcal{P}_k S_n(\theta) - e^{ik\theta} D_k(\theta)\|^2 = \sum_{k=1}^n \|\mathcal{P}_0 T_{n-k}(\theta) - D_0(\theta)\|^2 = o(n) \end{aligned}$$

Hence, by (4.8) and (4.4), we have

$$\limsup_{n \rightarrow \infty} \frac{\|E(S_n(\theta)|\mathcal{F}_0)\|^2}{n} = \limsup_{n \rightarrow \infty} \frac{\|S_n(\theta)\|^2 - \|M_n(\theta)\|^2}{n} = 0$$

by noting that  $\|M_n(\theta)\|^2 = n\|D_0(\theta)\|^2$ . Hence we have (4.7) in view of (4.9).

**Step 3. The CLT for the approximating martingale.**

It remains just to prove central limit theorem for complex valued martingale:

$$\frac{1}{\sqrt{n}} \sum_{1 \leq k \leq n} \exp(ik\theta) D_k(\theta).$$

As a matter of fact we shall provide a central limit theorem for the real part and imaginary part and show that in the limit they are independent. The proof was carefully written down in Wu (2005). By the Cramer-Wold device we have to study the limiting distribution of the martingale

$$\frac{1}{\sqrt{n}} \sum_{1 \leq k \leq n} [s\operatorname{Re}(\exp(ik\theta)D_k(\theta)) + t\operatorname{Im}(\exp(ik\theta)D_k(\theta))].$$

By the Raikov type of argument, in order to prove the CLT we have only to show

$$\frac{1}{n} \sum_{1 \leq k \leq n} [s\operatorname{Re}(\exp(ik\theta)D_k(\theta)) + t\operatorname{Im}(\exp(ik\theta)D_k(\theta))]^2 \xrightarrow{p} \frac{(s^2 + t^2)\sigma^2(\theta)}{2}.$$

This follows from combining the following two facts. First by stationarity

$$\frac{1}{n} \sum_{1 \leq k \leq n} |\exp(ik\theta)D_k(\theta)|^2 = \frac{1}{n} \sum_{1 \leq k \leq n} |D_k(\theta)|^2 \rightarrow E|D_0(\theta)|^2$$

and then by Lemma 5 in Wu (2005) for almost all  $\theta$

$$\frac{1}{n} \sum_{1 \leq k \leq n} [\exp(ik\theta)D_k(\theta)]^2 \rightarrow 0.$$

The rest is simple algebra. □

4.2. *Proof of Theorem 2.2.* Consider the product space  $([0, 2\pi] \times \Omega, \mathcal{B} \times \mathcal{F}, \lambda \times P)$ . Let  $\mathbf{P} = \lambda \times P$  and  $\mathbf{E}$  the corresponding expected value. We already have shown in Theorem 2.1, that for  $\lambda$ -almost all  $\theta$ ,

$$E \exp \left[ \frac{i}{\sqrt{n}} (s\operatorname{Re}(S_n(\theta)) + t\operatorname{Im}(S_n(\theta))) \right] \rightarrow \exp \left[ -\frac{(s^2 + t^2)g(\theta)}{4} \right]$$

for  $s, t \in \mathbb{R}$ . Then we integrate with  $\theta$  and by the Lebesgue dominated convergence theorem we obtain

$$\mathbf{E} \exp \left[ \frac{i}{\sqrt{n}} (s\operatorname{Re}(S_n(\theta)) + t\operatorname{Im}(S_n(\theta))) \right] \rightarrow \frac{1}{2\pi} \int_0^{2\pi} \exp \left[ -\frac{(s^2 + t^2)g(\theta)}{4} \right] d\theta .$$

We then identify the limiting distribution as being a mixture of two independent random variables: a standard normal variable with a variable uniformly distributed on  $[0, 2\pi]$ .  $\square$

4.3. *Proof of Proposition 2.1.* It is easy to see that the finite dimensional distributions are convergent. So we just have to prove tightness. From Billingsley (1999), stationarity and standard considerations this follows by Lemma 4.3.  $\square$

**Acknowledgement.** The first author would like to thank Sergey Utev for his conjecture related to Theorem 2.1 that motivated this paper and Richard Bradley for useful discussions. The authors are grateful to the referee for carefully reading the manuscript and useful suggestions.

## References.

- [1] Bary, N. K. (1964). *A Treatise on Trigonometric Series*. New York, Macmillan.
- [2] Billingsley, P. (1999) *Convergence of probability measures*, 2nd ed. Wiley, New York.
- [3] Bradley, R.C. (2007). *Introduction to strong mixing conditions*. Vol. 1, 2, 3. Kendrick Press.
- [4] Borodin, A. N. and Ibragimov, I. A. (1994), Limit theorems for functionals of random walks. *Trudy Mat. Inst. Steklov.*, 195. Transl. into English: Proc. Steklov Inst. Math. (1995), v 195, no.2.
- [5] Brockwell, P. J. and Davis, R. A. (1991). *Time Series: Theory and Methods*. New York, Springer.
- [6] Carleson, L. (1966). On convergence and growth of partial sums of Fourier series. *Acta Math.* 116 135-157.
- [7] Champeney, D. C. (1989). *A handbook of Fourier theorems*, Cambridge University Press, UK.
- [8] Dedecker J. and Rio, E. (2000). On the functional central limit theorem for stationary processes. *Ann. Inst. H. Poincaré Probab. Statist.* 36 1-34.
- [9] Dedecker J. and Merlevede F (2002). Necessary and sufficient conditions for the conditional central limit theorem. *Annals of Probability* 30, 1044-1081.
- [10] Fisher, R. A. (1929) Tests of significance in harmonic analysis. *Proc. Roy. Soc., A*, 125 54–59.
- [11] Gordin, M. and Lifshitz, B. (1981). A remark about a Markov process with normal transition operator, *Third Vilnius Conf. Proba. Stat., Akad. Nauk Litovsk*, Vilnius, 1, 147–148 (in Russian).
- [12] Gray, H. L., Zhang, N.-F. and Woodward, W.A. (1989). On generalized fractional processes. *J. Time Ser. Anal.* 10 233-257.
- [13] Hall, P. and Heyde, C.C. (1980). *Martingale limit theory and its application*. Academic Press.
- [14] Heyde, C. C. (1974). On the central limit theorem for stationary processes. *Z. Wahrsch. verw. Gebiete.* 30, 315-320.
- [15] Hoffman, K. (1988). *Banach spaces of analytic functions*. Dover Publications, New York.
- [16] Hunt, R. I. and Young, W. S. (1974). A weighted norm inequality for Fourier series, *Bull. Amer. Math. Soc.* 80, 274–277.

- [17] Kolmogorov, A. (1923): Une série de Fourier-Lebesgue divergente presque partout, *Fund. Math.*, 4, 324–328
- [18] Lacey M. and Terwilleger E. (2008) A Wiener-Wintner theorem for the Hilbert transform, *Ark. Mat.* 46 (2008), no. 2, 315–336.
- [19] Lahiri, S. N. (2003). A necessary and sufficient condition for asymptotic independence of discrete Fourier transforms under short- and long-range dependence. *Ann. Statist.* 31 613–641. MR1983544
- [20] Liu W. and Z. Lin (2009) The maximum of the periodogram of stationary process. *Ann. Statist.*, To Appear.
- [21] Olshen, R. A. (1967). Asymptotic properties of the periodogram of a discrete stationary process. *J. Appl. Probab.* 4 508-528.
- [22] Rootzen, H. (1976). Gordin's theorem and the periodogram. *J. Appl. Probab.* 13 365-370.
- [23] Rosenblatt, M. (1981). Limit theorems for Fourier transforms of functionals of Gaussian sequences. *Z. Wahrsch. Verw. Gebiete* 55 123-132.
- [24] Rosenblatt, M. (1985). *Stationary sequences and random fields*. Birkhäuser, Boston.
- [25] Schuster, A. (1898) On the investigation of hidden periodicities with application to a supposed 26 day period of meteorological phenomena, *Terrestrial Magnetism and Atmospheric Electricity*, 3, 13–41.
- [26] Terrin, N. and Hurvich, C. M. (1994). An asymptotic Wiener-Ito representation for the low frequency ordinates of the periodogram of a long memory time series. *Stochastic Process. Appl.* 54 297-307.
- [27] Walker, A. M. (1965). Some asymptotic results for the periodogram of a stationary time series. *J. Austral. Math. Soc.* 5 107-128.
- [28] Walker, A. M. (2000). Some results concerning the asymptotic distribution of sample Fourier transforms and periodograms for a discrete-time stationary process with a continuous spectrum. *J. Time Ser. Anal.* 21 95-109.
- [29] Woodroffe, M. (1992). A central limit theorem for functions of a Markov chain with applications to shifts. *Stochastic Process. Appl.* 41 33-44.
- [30] Wu, W. B. and Woodroffe, M. (2000). A central limit theorem for iterated random functions. *J. Appl. Probab.* 37 748-755.
- [31] Wu, W. B. (2005). Fourier transforms of stationary processes. *Proc. Amer. Math. Soc.* 133, 285-293
- [32] Yajima, Y. (1989). A central limit theorem of Fourier transforms of strongly dependent stationary processes. *J. Time Ser. Anal.* 10 375-383.
- [33] Wiener N. and Wintner A. (1941), On the ergodic dynamics of almost periodic systems, *Amer. J. Math.* 63, 794–824.
- [34] Zygmund, A. (2002) *Trigonometric Series*. Cambridge University Press, UK.

DEPARTMENT OF MATHEMATICAL SCIENCES  
 UNIVERSITY OF CINCINNATI  
 PO BOX 210025  
 CINCINNATI, OHIO 45221  
 USA  
 E-MAIL: peligrm@math.uc.edu

DEPARTMENT OF STATISTICS  
 UNIVERSITY OF CHICAGO  
 5734 S. UNIVERSITY AVENUE  
 CHICAGO, ILLINOIS 60637  
 USA  
 E-MAIL: wbwu@galton.uchicago.edu