

LIMIT THEOREMS FOR ITERATION STABLE TESSELLATIONS

BY TOMASZ SCHREIBER* AND CHRISTOPH THÄLE

Nicolaus Copernicus University Toruń and University of Osnabrück

The intent of this paper is to describe the large scale asymptotic geometry of STIT tessellations in \mathbb{R}^d , which form a rather new, rich and flexible class of random tessellations considered in stochastic geometry. For this purpose, martingale tools are combined with second-order formulas proved earlier to establish limit theorems for STIT tessellations. More precisely, a Gaussian functional central limit theorem for the surface increment process induced by a STIT tessellation relative to an initial time moment is shown. As second main result, a central limit theorem for the total edge length/facet surface is obtained, with a normal limit distribution in the planar case and, most interestingly, with a non-normal limit showing up in all higher space dimensions.

1. Introduction and results. Random tessellations or mosaics of \mathbb{R}^d (with $d \geq 2$) are locally finite families of compact convex random polytopes, which have no common interior points and cover the whole space. They form a central object studied in stochastic geometry, spatial statistics and related fields, see [14, 21] and the references cited therein. However, there are only very few mathematically tractable models, the most prominent examples include hyperplane and Voronoi tessellations, where most often the Poisson case is considered. A new class, the so-called STIT tessellations, was introduced recently by Mecke, Nagel and Weiß in [10, 11, 12, 13] and has quickly attracted considerable interest. These tessellations clearly show the potential to become a new reference model for both, theoretical and practical purposes. Whereas most research on random tessellations in the last decades was about mean values and mean value relations (see [14] for the recent state of the art), modern stochastic geometry focuses on distributional aspects ([1, 6] for example) and limit theorems, see [4, 5] and the references therein.

In contrast to the tessellations studied so far, the STIT model has the

*26.06.1975-01.12.2010

AMS 2000 subject classifications: Primary 60D05, 60F17; secondary 60F05, 60J75

Keywords and phrases: Central limit theorem, functional limit theorem, iteration/nesting, Markov processes, martingale theory, random tessellations, stochastic stability, stochastic geometry

additional feature of arising as a result of a spatio-temporal *dynamic* construction. From this point of view, limit theorems for STIT tessellations are particularly interesting. As recently pointed out in [15], we expect that the large scale asymptotic of dynamic models for spatial random structures will become of great importance in stochastic geometry in the near future.

Let us recall the basic construction of tessellations that arise as a result of repeated cell division. To this end, we identify the space \mathcal{H} of hyperplanes in \mathbb{R}^d with the parameter space $\mathbb{R}_+ \times \mathcal{S}_{d-1}$ and the hyperplane $\{x \in \mathbb{R}^d : \langle x, u \rangle = r\}$ with the pair $(r, u) \in \mathbb{R}_+ \times \mathcal{S}_{d-1}$ and let Λ be a measure on \mathcal{H} , which admits under the described polar identification a representation

$$(1) \quad \Lambda = \ell_+ \otimes \mathcal{R},$$

where ℓ_+ is the Lebesgue measure on the positive real half-axis and where \mathcal{R} is a probability measure on the unit sphere \mathcal{S}_{d-1} . Throughout this paper we always require that the support of \mathcal{R} spans the whole space, i.e. that $\text{span}(\text{supp}(\mathcal{R})) = \mathbb{R}^d$, and we say in this case that Λ is non-degenerate. Further, let $t > 0$ be fixed and let $W \subset \mathbb{R}^d$ be a compact convex set with interior points in which our construction of a random tessellation $Y(t\Lambda, W)$ is carried out. In a first step, we assign to the window W a random lifetime. Upon expiry of its lifetime, the primordial cell W dies and splits into two sub-cells W^+ and W^- separated by a hyperplane hitting W , which is chosen according to suitable restriction and normalization of Λ . The resulting new cells W^+ and W^- are again assigned independently with random lifetimes and the entire construction continues recursively until the previously fixed deterministic time threshold t is reached. The described process of recursive cell divisions is called the *MNW-construction* through this paper (M-N-W stand for the inventors of the model) and the resulting random subdivision of W is denoted by $Y(t\Lambda, W)$, see Figure 1 for illustrations. Note that the cells of $Y(t\Lambda, W)$ are polyhedral except possibly those hitting the potentially curved boundary of W , so that upon boundary effects $Y(t\Lambda, W)$ is a random tessellation of W .

In order to ensure the Markov property of the above construction in the continuous-time parameter t , we assume from now on that the lifetimes arising in the MNW-construction (including that of the initial window W) are exponentially distributed. Moreover, we assume that the parameter of the exponentially distributed lifetime of an individual cell $[c]$ equals $\Lambda([c])$, where $[c]$ stands for the set of all parameter values of hyperplanes hitting c . In this special situation, the random tessellations $Y(t\Lambda, W)$ fulfill a stochastic stability property under the operation of iteration of tessellations and are for this reason called random *STIT tessellations*, see Section 2 below for

details.

Having studied the first- and second-order properties of STIT tessellations in [19, 20], we consider in this paper the central limit problem. This problem will be considered in two closely related settings, interestingly leading to results of very different qualitative nature. First, we shall focus our interest on the residual length/surface increment process, arising, respectively, as cumulative length or surface area of the cell-separating $(d - 1)$ -polyhedral facets born *after* a certain fixed initial time in the course of the MNW-construction. In this set-up we shall establish a central limit theorem with a Gaussian limiting variable. Next, we shall pass to the *total* length/surface, taking into account also the segments/facets born at the very initial *big bang* stages of the MNW-construction, as descriptively termed in [10]. It turns out that, whereas in dimension 2 the Gaussian convergence, is preserved, this is no more the case for dimensions 3 and higher, where non-Gaussian limits arise. This apparently surprising phenomenon is in fact due to the influence of the big bang phase in the MNW-construction itself, which is negligible in the planar case, but turns out to be crucial in higher dimensions.

We are now going to describe some of our limit theorems in more detail. For a compact convex set W as above, we put $W_R := RW$ for $R > 0$ and let Λ be some fixed hyperplane measure as in (1). Therefore and in order to simplify the notation we will write from now on $Y(t, W)$ instead of $Y(t\Lambda, W)$. Our first limit theorem deals with the total surface area $\text{Vol}_{d-1}(Y(t, W))$ induced by the MNW-construction within the time period $[s_0, 1]$, where $0 < s_0 < 1$ is some positive initial time moment.

THEOREM 1. *For each $0 < s_0 < 1$, the random variable*

$$\frac{1}{R^{d/2}} [(\text{Vol}_{d-1}(Y(1, W_R)) - \mathbb{E}\text{Vol}_{d-1}(Y(1, W_R))) \\ - (\text{Vol}_{d-1}(Y(s_0, W_R)) - \mathbb{E}\text{Vol}_{d-1}(Y(s_0, W_R)))]$$

converges, as $R \rightarrow \infty$, in law to $\mathcal{N}(0, V_W(\text{Vol}_{d-1}, \Lambda) \int_{s_0}^1 s^{1-d} ds)$, a normal distribution with mean 0 and variance $V_W(\text{Vol}_{d-1}, \Lambda) \int_{s_0}^1 s^{1-d} ds$, where $V_W(\text{Vol}_{d-1}, \Lambda)$ is explicitly given by (9) or alternatively (17) below.

This statement cannot be extended to $s_0 \downarrow 0$, as would be of interest as potentially leading to a Gaussian limit for the (centered and suitably normalized) total edge length/surface area $\text{Vol}_{d-1}(Y(1, W_R))$. The problem is that the variance integral $V_W(\text{Vol}_{d-1}, \Lambda) \int_{s_0}^t s^{1-d} ds$ diverges at 0. However, this difficulty can be overcome for $d = 2$, but not for $d > 2$. Indeed, in the planar case the asymptotic behavior of the total edge length turns out to be Gaussian:

THEOREM 2. For a STIT tessellation $Y(1, W_R)$ in the plane we have

$$\frac{1}{R\sqrt{\log R}}[\text{Vol}_1(Y(1, W_R)) - \mathbb{E}\text{Vol}_1(Y(1, W_R))] \Longrightarrow \mathcal{N}(0, V_W(\text{Vol}_1, \Lambda)),$$

where \Longrightarrow means convergence in law and where again $V_W(\text{Vol}_1, \Lambda)$ is explicitly known and given by (9) or (17) below.

In fact, Theorem 1 and Theorem 2 are direct consequences of our much stronger functional central limit theorems, Theorems 4 and 5 below.

For space dimensions $d > 2$ we claim that the Gaussian convergence cannot be preserved. Even though we are able to show this fact for all W and translation invariant Λ by establishing non-Gaussian tail decay, for simplicity and in order to keep the argument transparent we only give a proof for an more easily tractable particular case, in which all cells have the shape of cuboids (rectangular parallelepipeds). The study of more involved properties of the resulting random field is postponed to a future paper.

THEOREM 3. Fix $d > 2$, take $W = [0, 1]^d$ and consider the hyperplane measure

$$(2) \quad \Lambda := (2d)^{-1} \sum_{i=1}^d \int_{-\infty}^{+\infty} \delta_{re_i + e_i^\perp} dr,$$

where e_i , $i = 1, \dots, d$ are vectors of the standard orthonormal basis for \mathbb{R}^d and $\delta_{re_i + e_i^\perp}$ is the unit mass concentrated on the hyperplane orthogonal to e_i at distance r from the origin (here e_i^\perp stands for the orthogonal complement of e_i). In this setting,

$$\frac{1}{R^{d-1}}[\text{Vol}_{d-1}(1, W_R) - \mathbb{E}\text{Vol}_{d-1}(1, W_R)]$$

converges, as $R \rightarrow \infty$, to a non-Gaussian square-integrable random variable $\Xi(W)$ with explicitly known variance given by (23) below.

The structure of the paper is as follows: In the next section we recall some properties of STIT tessellations needed for the proofs of our limit theorems. We also recall there some of the facts from [19, 20] in order to keep the paper self-contained and present the exact statements of our functional central limit theorems. The proofs of our results are the content of Section 3.

We would like to remark that an extended version [18] of this text is available online and, moreover, that the results in the present paper form the basis of our works [16, 17].




FIG 1. Simulations of a planar and a spatial stationary and isotropic STIT tessellation (kindly provided by Joachim Ohser and Claudia Redenbach)

2. Background material and statement of the functional limit theorems. We start by rephrasing some of the properties of the STIT tessellations $Y(t\Lambda, W)$ as defined in the introduction, the proofs of which may be found in [13].

- $Y(t\Lambda, W)$ is consistent in that $Y(t\Lambda, W) \cap V \stackrel{D}{=} Y(t\Lambda, V)$ for convex $V \subset W$ (here and below $\stackrel{D}{=}$ stands for equality in distribution) and thus $Y(t\Lambda, W)$ can be extended to a random tessellation $Y(t\Lambda)$ in the whole space \mathbb{R}^d in such a way that $Y(t\Lambda) \cap W$ has the same distribution as $Y(t\Lambda, W)$.

This way, instead of interpreting $Y(t\Lambda, W)$ as outcome of the MNW-construction carried out in W , $Y(t\Lambda, W)$ can also be understood as restriction of the whole space random tessellation $Y(t\Lambda)$ (which is a proper tessellation in the usual sense of stochastic geometry as discussed in the introduction) to W .

- $Y(t\Lambda)$ is a stationary random tessellation, i.e. stochastically translation invariant. If, moreover, Λ is the unit-density isometry-invariant hyperplane measure Λ_{iso} , or equivalently if \mathcal{R} in (1) is the uniform distribution ν_{d-1} on \mathcal{S}_{d-1} , then $Y(t\Lambda_{\text{iso}})$ is even isotropic, i.e. stochastically invariant under rotations around the origin.
- $Y(t\Lambda)$ is stable under the operation of iteration, denoted by \boxplus . This

is to say

$$Y(t\Lambda) \stackrel{D}{=} m(Y((t/m)\Lambda) \boxplus \cdots \boxplus Y((t/m)\Lambda)), \quad m = 2, 3, \dots$$

For this reason, $Y(t\Lambda)$ is called a random STIT tessellation. This property was discussed in detail in [13, 19] and we refer to these papers and the references cited therein for further discussion, because our arguments do not explicitly use the stochastic stability, but its consequences.

- The surface density of $Y(t\Lambda)$, i.e. the mean surface area of cell boundaries of $Y(t\Lambda)$ per unit volume equals t . In particular, the mean surface area of facets arising in the MNW-construction during time $[0, t]$ within a compact convex $W \subset \mathbb{R}^d$ with interior points is given by $t \text{Vol}_d(W)$.
- STIT tessellations enjoy the following scaling property:

$$tY(t\Lambda) \stackrel{D}{=} Y(\Lambda),$$

i.e. the tessellation $Y(t\Lambda)$ of surface density t upon rescaling by factor t has the same distribution as $Y(\Lambda)$, the STIT tessellation with surface density 1.

- The intersection of a STIT tessellation in \mathbb{R}^d with a k -dimensional affine subspace L_k ($1 \leq k \leq d-1$) that is independent of the tessellation induces a STIT tessellation in L_k .
- STIT tessellations have Poisson typical cells, which is to say that the distribution of the interior of the typical cell $\text{TypicalCell}(Y(t\Lambda))$ of $Y(t\Lambda)$ coincides with that of a Poisson hyperplane tessellation $\text{PHT}(t\Lambda)$ having intensity measure $t\Lambda$, see the discussion after Theorem 1 in [19] or [12].

For the non-specialized reader let us remark that the typical cell of a tessellation is what we get when we choose equiprobably a cell of the tessellation at random out of a ‘large’ observation window. The exact definition makes use of Palm theory for which we refer to [14]. Moreover, a Poisson hyperplane tessellation $\text{PHT}(t\Lambda)$ is a random subdivision of \mathbb{R}^d induced by a Poisson point process on the space of hyperplanes \mathcal{H} having intensity measure $t\Lambda$.

The finite volume continuous-time incremental MNW-construction of random STIT tessellations, as discussed in the introduction, clearly has the Markov property in the continuous-time parameter, whence natural martingales arise, which will be of crucial importance for our further considerations. In fact, this observation was the starting point of [19], where a class of martingales associated to STIT tessellations was constructed. In order to streamline our discussion we do not repeat the full theory here, but

rephrase the martingale property of two stochastic processes, on which the proofs of our limit theorems are based on. To this end, let Y be some instant of $Y(t\Lambda, W)$ and let $\phi(\cdot)$ be a measurable facet functional of the form

$$(3) \quad \phi(f) = \text{Vol}_{d-1}(f)\zeta(\vec{\mathbf{n}}(f))$$

with $\vec{\mathbf{n}}(f)$ standing for the unit normal to a facet f and ζ for a bounded measurable function on \mathcal{S}_{d-1} . Moreover, denote the collection of cell-separating $(d-1)$ -dimensional facets, usually referred to as $(d-1)$ -dimensional maximal polytopes, arising in subsequent splits in the MNW-construction by $\text{MaxPolytopes}_{d-1}(Y)$. (Note that some of these polytopes can be chopped-off by the possibly curved boundary of the convex window W in which Y is constructed and are no polytopes in the usual sense. However, we somehow abuse notation and remark that this technical issue causes no difficulties in our theory, because of the special form of the facet functional ϕ .) Let us further define $\Sigma_\phi(Y)$ by

$$\Sigma_\phi(Y) := \sum_{f \in \text{MaxPolytopes}_{d-1}(Y)} \phi(f)$$

and $A_{\phi^2}(Y)$ to be

$$A_{\phi^2}(Y) := \int_{[W]} \sum_{f \in \text{Cells}(Y \cap H)} \phi^2(f) \Lambda(dH)$$

with $\text{Cells}(Y \cap H)$ standing for the set of $(d-1)$ -dimensional cells of the tessellation $Y \cap H$ induced by the intersection of Y with a hyperplane H (again, some of these cells may have a curved boundary, because of the intersection with the construction window). Let us further introduce the bar notation $\bar{\Sigma}_\phi(Y)$ for the centred version $\Sigma_\phi(Y) - \mathbb{E}\Sigma_\phi(Y)$ of $\Sigma_\phi(Y)$. Then, we have (see [19, 20]):

PROPOSITION 1. *The two stochastic processes*

$$(4) \quad \bar{\Sigma}_\phi(Y(t\Lambda, W)) \text{ and } \bar{\Sigma}_\phi^2(Y(t\Lambda, W)) - \int_0^t A_{\phi^2}(Y(s\Lambda, W)) ds$$

are both \mathfrak{S}_t -martingales, where \mathfrak{S}_t stands for the filtration generated by $(Y(s\Lambda, W))_{0 \leq s \leq t}$.

In particular, see [9, Paragraph I.8], by (4) the martingale $\bar{\Sigma}_\phi(Y(t\Lambda, W))$ has its predictable quadratic variation process $\langle \bar{\Sigma}_\phi(Y(\cdot, W)) \rangle$ absolutely continuous (in the sense of functions) and given by

$$(5) \quad \langle \bar{\Sigma}_\phi(Y(\cdot, W)) \rangle_t = \int_0^t A_{\phi^2}(Y(s\Lambda, W)) ds.$$

Besides these martingale tools, we will also make use of the following formula for the variance $\text{Var}(\Sigma_\phi(Y(t\Lambda, W)))$ of $\Sigma_\phi(Y(t\Lambda, W))$, $W \subset \mathbb{R}^d$ compact, convex and with interior points, established in full generality in [20] in order to calculate the variance of the limit random variable of our non-Gaussian limit theorem:

PROPOSITION 2. *For any non-degenerate translation-invariant Λ of the form (1) and ϕ as in (3), we have*

$$(6) \quad \text{Var}(\Sigma_\phi(Y(t\Lambda, W))) = \int_{[W]} \zeta^2(\vec{n}(H)) \int_{W \cap H} \int_{W \cap H} \frac{1 - \exp(-t\Lambda([xy]))}{\Lambda([xy])} dx dy \Lambda(dH),$$

where $[xy]$ stands for the set parameter values of hyperplanes hitting the line segment xy connecting x with y .

Let us further recall from [20] that the variance of the total edge length of a stationary and isotropic STIT tessellation $Y(t\Lambda_{\text{iso}}, W_R)$ in the plane behaves asymptotically like

$$(7) \quad \text{Var}(\text{Vol}_1(Y(t\Lambda_{\text{iso}}, W_R))) \sim \pi \text{Vol}_2(W) R^2 \log R, \quad \text{as } R \rightarrow \infty,$$

where W is again a compact convex set as above. Indeed, this can be seen from the general statement in Proposition 2 when combined with some integral-geometric tools. Note that the asymptotic variance expression for the total edge length is independent of t . However, for all space dimensions > 2 , the surface density t enters the asymptotic variance expression as shown in detail in [20].

We can now turn to the statement of our functional limit theorems, from which Theorems 1 and 2 are direct consequences of. As in introduction we fix a hyperplane measure Λ and suppress from now on the reference to Λ , for example by writing $Y(t, W_R)$ instead of $Y(t\Lambda, W_R)$.

THEOREM 4. *For each $0 < s_0 < 1$ the centred surface increment process*

$$\left(\mathcal{S}_{s_0, t}^{R, W} := \frac{1}{R^{d/2}} [\bar{\Sigma}_\phi(Y(t, W_R)) - \bar{\Sigma}_\phi(Y(s_0, W_R))] \right)_{t \in [s_0, 1]}$$

converges in law, as $R \rightarrow \infty$, on the space $\mathcal{D}[s_0, 1]$ of right continuous functions with left-hand limits (càdlàg) on $[s_0, 1]$, endowed with the Skorokhod J_1 -topology [9, Chapter 6.1], to a time-changed Wiener process

$$t \mapsto \mathcal{W}_{V_W(\phi, \Lambda)} \int_{s_0}^t s^{1-d} ds,$$

where $\mathcal{W}_{(\cdot)}$ is the standard Wiener process and $V_W(\phi, \Lambda)$ is given by (9) or alternatively (17) below. In particular,

$$\mathcal{S}_{s_0,1}^{R,W} = \frac{1}{R^{d/2}} [\bar{\Sigma}_\phi(Y(1, W_R)) - \bar{\Sigma}_\phi(Y(s_0, W_R))]$$

converges in law to $\mathcal{N}(0, V_W(\phi, \Lambda) \int_{s_0}^1 s^{1-d} ds)$, a normal distribution with mean 0 and variance $V_W(\phi, \Lambda) \int_{s_0}^1 s^{1-d} ds$.

We turn now the functional convergence of the total length process in the planar case. Write

$$\tau(s, R) := \exp([\log R - \log \log R](s - 1)) = R^{s-1}(\log R)^{1-s}$$

and define the *total length process*

$$\mathcal{L}_s^{R,W} := \frac{1}{R\sqrt{\log R}} \bar{\Sigma}_\phi(Y(\tau(s, R), W_R)), \quad s \in [0, 1].$$

THEOREM 5. *The total length process $(\mathcal{L}_s^{R,W})_{s \in [0,1]}$ converges in law, as $R \rightarrow \infty$, on the space $\mathcal{D}[0, 1]$ of càdlàg functions on $[0, 1]$, endowed with the Skorokhod J_1 -topology, to $(\sqrt{V_W(\phi, \Lambda)} \mathcal{W}_s)_{s \in [0,1]}$ where, again, $\mathcal{W}_{(\cdot)}$ stands for the standard Wiener process.*

REMARK 1. *We consider in this paper facet functionals of the form $\phi(\cdot) = \text{Vol}_{d-1}(\cdot) \zeta(\vec{\mathbf{n}}(\cdot))$, see (3). Taking $\zeta \equiv 1$, Theorems 4 - 5 reduce to the total surface area case discussed at the beginning. However, the additional flexibility induced by the introduction of $\zeta(\cdot)$ implies that our results allow to conclude limit theorems that are sensitive with respect to direction. Taking for example $\zeta(\cdot) = \mathbf{1}\{\vec{\mathbf{n}}(\cdot) \in U(n)\}$ to be the indicator function of a small neighbourhood $U(n)$ of a fixed direction $n \in \mathcal{S}_{d-1}$ satisfying $\mathcal{R}(U(n)) > 0$ (recall the decomposition (1)), yields central limit theorems also for the collection of tessellation facets having their normals in $U(n)$. This means that our results are not only valid for the whole STIT tessellation, but also for parts in arbitrary space directions.*

REMARK 2. *So far we have restricted our considerations to space dimensions $d \geq 2$. STIT tessellations and their limit theory on the line can also be considered. However, in [12] it was shown that a STIT tessellation on \mathbb{R} is nothing than a homogeneous Poisson point process, or more precisely the intervals between its points. These point processes and their limit theory are well known and for this reason we have focused on the cases $d \geq 2$.*

3. Proofs. After having rephrased some background material on STIT tessellations in the previous section, we are now prepared to present the proofs of our limit theorems. Let us briefly recall that we will deal with a fixed translation-invariant hyperplane measure Λ and that for this reason we shall write for example $Y(t)$ instead of $Y(t\Lambda)$ without confusion. Moreover, we fix some compact and convex set $W \subset \mathbb{R}^d$ having interior points and write $W_R = RW$ for W dilated by a factor $R > 0$. Moreover, recall that the facet functionals we are dealing with have the representation (3), that $\Sigma_\phi(Y(t))$ was defined in (4) and that the bar notation $\bar{\Sigma}_\phi(Y(t))$ stands for the centred version $\Sigma_\phi(Y(t)) - \mathbb{E}\Sigma_\phi(Y(t))$. We start now with the

PROOF OF THEOREM 4. Notice first that, because of $\Lambda([W_R]) = R\Lambda([W])$ we have,

$$\begin{aligned}
& \frac{1}{R^d} A_{\phi^2}(Y(1, W_R)) \\
&= \frac{1}{R} \int_{[W_R]} \frac{1}{R^{d-1}} \zeta^2(\bar{\mathbf{n}}(H)) \sum_{f \in \text{Cells}(Y(1, W_R) \cap H)} \text{Vol}_{d-1}^2(f) \Lambda(dH) \\
(8) \quad &= \int_{[W]} \frac{1}{R^{d-1}} \zeta^2(\bar{\mathbf{n}}(H)) \sum_{f \in \text{Cells}(Y(1, W_R) \cap RH)} \text{Vol}_{d-1}^2(f) \Lambda(dH).
\end{aligned}$$

We claim that, upon letting $R \rightarrow \infty$, this converges in probability to

$$\begin{aligned}
& V_W(\phi, \Lambda) := \\
& \int_{[W]} \zeta^2(\bar{\mathbf{n}}(H)) \text{Vol}_{d-1}(W \cap H) \frac{\mathbb{E} \text{Vol}_{d-1}^2(\text{TypicalCell}(Y(1) \cap H))}{\mathbb{E} \text{Vol}_{d-1}(\text{TypicalCell}(Y(1) \cap H))} \Lambda(dH) \\
(9) \quad &= \text{Vol}_d(W) \int_{\mathcal{S}_{d-1}} \zeta^2(u) \frac{\mathbb{E} \text{Vol}_{d-1}^2(\text{TypicalCell}(Y(1) \cap u^\perp))}{\mathbb{E} \text{Vol}_{d-1}(\text{TypicalCell}(Y(1) \cap u^\perp))} \mathcal{R}(du),
\end{aligned}$$

where \mathcal{R} is the directional distribution of the tessellation as given by (1). To see it, recall that $Y(1) \cap RH$ is a STIT tessellation in RH for each $R > 0$ and $H \in \mathcal{H}$. Thus, applying [14, (4.6) and Thm. 4.1.3] to this tessellation and the fact that $\mathbb{E} \text{Vol}_d(\text{TypicalCell}(Y(1) \cap u^\perp))$ is the same as the inverse cell density of the tessellation $Y(1) \cap u^\perp$, see (10.4) ibidem, we get

$$\begin{aligned}
& \lim_{R \rightarrow \infty} \frac{1}{R^{d-1}} \mathbb{E} \sum_{f \in \text{Cells}(Y(1, W_R) \cap RH)} \text{Vol}_{d-1}^2(f) \\
(10) \quad &= \text{Vol}_{d-1}(W \cap H) \frac{\mathbb{E} \text{Vol}_{d-1}^2(\text{TypicalCell}(Y(1) \cap H))}{\mathbb{E} \text{Vol}_{d-1}(\text{TypicalCell}(Y(1) \cap H))}.
\end{aligned}$$

Next, we observe that $Y(1, W_R) \cap RH \stackrel{D}{=} Y(1) \cap R \cdot_H (H \cap W)$, where \cdot_H is the scalar multiplication relative in H , that is to say $H \ni R \cdot_H x = p_H(0) + R(x - p_H(0))$, $x \in H$ with p_H standing for the orthogonal projection on H . Thus, using the recently developed strong mixing and tail triviality theory for STIT tessellations, especially [8, Thm. 2], noting that tail trivial stationary processes are ergodic [3, Prop. 14.9] and then applying the multidimensional ergodic theorem, see e.g. Corollary 14.A5 ibidem, to $\frac{1}{R^{d-1}} \sum_{f \in \text{Cells}(Y(1) \cap R \cdot_H (H \cap W))} \text{Vol}_{d-1}^2(f)$, we get from (10) that

$$\begin{aligned} & \lim_{R \rightarrow \infty} \frac{1}{R^{d-1}} \sum_{f \in \text{Cells}(Y(1, W_R) \cap RH)} \text{Vol}_{d-1}^2(f) \\ &= \text{Vol}_{d-1}(W \cap H) \frac{\mathbb{E} \text{Vol}_{d-1}^2(\text{TypicalCell}(Y(1) \cap H))}{\mathbb{E} \text{Vol}_{d-1}(\text{TypicalCell}(Y(1) \cap H))} \end{aligned}$$

in probability. Putting this together with (8) and integrating over $[W]$ yields

$$(11) \quad \lim_{R \rightarrow \infty} \frac{1}{R^d} A_{\phi^2}(Y(1, W_R)) = V_W(\phi, \Lambda) \quad \text{in probability,}$$

as required.

Note now that by the scaling properties of $Y(s, W_R)$ and ϕ^2 for $s > 0$ we have

$$(12) \quad \begin{aligned} \frac{1}{R^d} A_{\phi^2}(Y(s, W_R)) &\stackrel{D}{=} \frac{1}{R^d s^{2d-1}} A_{\phi^2}(Y(1, W_{sR})) \\ &\stackrel{D}{=} \frac{1}{s^{d-1}} \frac{1}{(Rs)^d} A_{\phi^2}(Y(1, W_{sR})). \end{aligned}$$

Thus, combining (11) with the scaling relation (12) we get

$$(13) \quad \lim_{R \rightarrow \infty} \frac{1}{R^d} A_{\phi^2}(Y(s, W_R)) = \frac{1}{s^{d-1}} V_W(\phi, \Lambda)$$

in probability uniformly in $s \in [s_0, 1]$.

This crucial statement puts us now in context of the general martingale limit theory. Indeed, using Proposition 1 we see that

$$\mathcal{S}_{s_0, s}^{R, W} = \frac{1}{R^{d/2}} [\bar{\Sigma}_{\phi}(Y(1, W_R)) - \bar{\Sigma}_{\phi}(Y(s_0, W_R))]$$

is a martingale with absolutely continuous predictable quadratic variation process

$$(14) \quad \langle \mathcal{S}_{s_0, \cdot}^{R, W} \rangle_t = \int_{s_0}^t \frac{1}{R^d} A_{\phi^2}(Y(s, W_R)) ds$$

by (4) and (11), see again Paragraph I.8 in [9]. In these terms, (13) yields for each t

$$(15) \quad \lim_{R \rightarrow \infty} \langle \mathcal{S}_{s_0, \cdot}^{R,W} \rangle_t = \int_{s_0}^t \frac{1}{s^{d-1}} V_W(\phi, \Lambda) ds \quad \text{in probability.}$$

We now want to apply the martingale functional central limit theorem. Whereas this is well known for continuous martingales, we need a version for martingales in the Skorokhod space $\mathcal{D}[s_0, 1]$. In this paper, we will make use of the version formulated as Theorem 2.1 in the survey article [22]. In order to apply this theorem, several conditions have to be checked. Condition (ii.6) in [22, Thm. 2.1] is just (15), whereas condition (ii.4) there is trivially verified, because the predictable quadratic variation $\langle \mathcal{S}_{s_0, \cdot}^{R,W} \rangle$ has no jumps by (14). It remains to check the condition (ii.5) ibidem, which is that the second moment of the maximum jump $\mathcal{J}(\mathcal{S}_{s_0, \cdot}^{R,W}; 1)$ of the process $(\mathcal{S}_{s_0, s}^{R,W})_{s \in [s_0, 1]}$ goes to 0 as $R \rightarrow \infty$. More precisely,

$$\mathcal{J}(\mathcal{S}_{s_0, \cdot}^{R,W}; 1) = \sup_{s_0 \leq t \leq 1} |\mathcal{S}_{s_0, t}^{R,W} - \mathcal{S}_{s_0, t-}^{R,W}|, \quad \mathcal{S}_{s_0, t-}^{R,W} = \lim_{s \uparrow t} \mathcal{S}_{s_0, s}^{R,W},$$

and we have to check that $\mathbb{E} \mathcal{J}^2(\mathcal{S}_{s_0, \cdot}^{R,W}; 1) \rightarrow 0$, as $R \rightarrow \infty$. To this end, note first that, with probability one, $\mathcal{J}(\mathcal{S}_{s_0, \cdot}^{R,W}; 1)$ is bounded from above by a constant multiple of $R^{-d/2}$ times the $(d-1)$ -th power of the diameter of the largest cell of $Y(s_0, W_R)$. Since the typical cell distribution of $Y(s_0)$ is the same as that of a Poisson hyperplane tessellation with intensity measure $s_0 \Lambda$ (see Theorem 1 in [19] or Section 2 above), we conclude by standard properties of Poisson hyperplane tessellations that the expected number $e(Y(s_0, W_R), D)$ of cells in $Y(s_0, W_R)$ with diameter exceeding D is of the order $O(R^d \exp(-D))$. Indeed, writing $\text{diam}(c)$ for the diameter of a cell c and $\mathbf{1}\{\cdot\}$ for the usual indicator function, we can rewrite $e(Y(s_0, W_R), D)$ as

$$e(Y(s_0, W_R), D) = \mathbb{E} \sum_{c \in \text{Cells}(Y(s_0, W_R))} \mathbf{1}\{\text{diam}(c) > D\},$$

which by Theorem 4.1.3 in [14] is of the same order as the mean number $N(Y(s_0, W_R)) = \mathbb{E} \sum_{c \in \text{Cells}(Y(s_0, W_R))} 1$ of cells in $Y(s_0, W_R)$ times the probability that the typical cell diameter of $Y(s_0)$ exceeds D (the additional condition in Theorem 4.3.1 in [14] is easily verified by using Steiner's formula together with the fact that the typical cell of a Poisson hyperplane tessellation has finite mean intrinsic volumes, see Theorem 10.3.3 ibidem). Thus, $e(Y(s_0, W_R), D)$ satisfies

$$e(Y(s_0, W_R), D) = O(N(Y(s_0, W_R)) \mathbb{P}(\text{diam}(\text{TypicalCell}(Y(s_0))) > D)).$$

Stationarity of the tessellation implies that the first factor is of volume order R^d . We claim that the second term is bounded from above by $c_1 e^{-c_2 s_0 D}$, where c_1 and c_2 are constants, which depend on the hyperplane measure Λ . To this end we notice first that the typical cell of $Y(s_0)$ is stochastically smaller than the almost surely uniquely determined cell Z_0 of $Y(s_0)$ containing the origin, cf. Corollary 10.4.1 in [14]. Moreover, equation (20) in [6] with $\Sigma = \text{diam}$, $\varepsilon = 0$ and $\alpha = 1$ there (the other parameters are then $k = \tau = 1$) implies that there are constants $c_1, c_2 > 0$ depending on the hyperplane measure Λ such that $\mathbb{P}(\text{diam}(Z_0) > D) \leq c_1 e^{-c_2 s_0 D}$, which implies

$$\mathbb{P}(\text{diam}(\text{TypicalCell}(Y(s_0))) > D) \leq \mathbb{P}(\text{diam}(Z_0) > D) \leq c_1 e^{-c_2 s_0 D}.$$

Putting these two issues together leads to the desired order $O(R^d \exp(-D))$ for the expected number of cells in $Y(s_0, W_R)$ with diameter exceeding D . Recalling that $\mathcal{J}(\mathcal{S}_{s_0, \cdot}^{R, W}; 1)$ is bounded from above by a constant multiple of $R^{-d/2}$ times the $(d-1)$ -th power of the diameter of the largest cell of $Y(s_0, W_R)$ and putting $u = D^{d-1} R^{-d/2}$ we find

$$(16) \quad \mathbb{P}(\mathcal{J}(\mathcal{S}_{s_0, \cdot}^{R, W}; 1) > u) = O(R^d \exp(-R^{d/(2d-2)} u^{1/(d-1)})).$$

Clearly, (16) is sufficient to guarantee that

$$\lim_{R \rightarrow \infty} \mathbb{E} \mathcal{J}^2(\mathcal{S}_{s_0, \cdot}^{R, W}; 1) = 0,$$

which gives the required condition (ii.5) of Theorem 2.1 in [22]. This result yields now the functional convergence as stated in our Theorem 4. \square

Before turning to the proof of Theorem 5 we provide an alternative formula for the factor $V_W(\phi, \Lambda)$. Readers not specialized in convex or stochastic geometry could also skip this alternative representation and directly jump to the next paragraph, because Proposition 3 will not be used in the sequel. However, having such a more explicit variance expression is useful for other purposes and has already been used in our work [17]. We denote, as in [14] or [19], by Π the associated zonoid of a Poisson hyperplane tessellation with intensity measure Λ , by Π° its polar body and by \mathcal{R} the directional distribution of the STIT tessellation from (1), see [14] for the precise definitions of Π and Π° .

PROPOSITION 3. *We have*

$$(17) \quad V_W(\phi, \Lambda) = \text{Vol}_d(W) \frac{(d-1)!}{2^{d-1}} \int_{\mathcal{S}_{d-1}} \zeta^2(u) \text{Vol}_{d-1}((\Pi|u^\perp)^\circ) \mathcal{R}(du),$$

where $\Pi|u^\perp$ stands for the orthogonal projection of Π onto the hyperplane u^\perp , and where the polar body $(\Pi|u^\perp)^\circ$ is considered relative to u^\perp . In the isotropic case, i.e. when $\mathcal{R} = \nu_{d-1}$, the uniform distribution on the unit sphere \mathcal{S}_{d-1} , this reduces to

$$V_W(\phi, \Lambda_{\text{iso}}) = \text{Vol}_d(W) 2^{d-1} \pi^{d-\frac{3}{2}} \frac{\Gamma\left(\frac{d+1}{2}\right)^{d-1}}{\Gamma\left(\frac{d}{2}\right)^{d-2}} \int_{\mathcal{S}_{d-1}} \zeta^2(u) \nu_{d-1}(du).$$

In particular for $\zeta \equiv 1$, $W = B_1^d$ the unit ball and $d = 2$ and $d = 3$ we conclude the exact values

$$V_{B_1^2}(\text{Vol}_1, \Lambda_{\text{iso}}) = \pi^2 \quad \text{and} \quad V_{B_1^3}(\text{Vol}_2, \Lambda_{\text{iso}}) = \frac{32}{3} \pi^2.$$

PROOF OF PROPOSITION 3. At first, [2, Cor. 3.7] provides a general formula for the second moment of the volume of the typical Poisson cell of a stationary Poisson hyperplane tessellation $\text{PHT}(\Lambda)$ in \mathbb{R}^d having intensity measure Λ . In terms of the zonoid Π it reads

$$\mathbb{E} \text{Vol}_d^2(\text{TypicalCell}(\text{PHT}(\Lambda))) = \frac{d!}{2^d} \frac{\text{Vol}_d(\Pi^\circ)}{\text{Vol}_d(\Pi)},$$

where we have used [14, Eq. (4.63)]. Moreover, the first volume moment of the typical cell of a Poisson hyperplane tessellation $\text{PHT}(\Lambda)$ equals $1/\text{Vol}_d(\Pi)$ according to [14, Thm. 10.3.3 and (10.4)]. Using now Eq. (4.61) ibidem and the fact that STIT tessellations have Poisson typical cell distributions and replacing d by $d-1$ in the last two formulas we obtain (17) immediately from (9). The precise value in the stationary and isotropic case can be calculated from the fact that in this case, Π is a ball with a known radius, see [14]. \square

PROOF OF THEOREM 5. Recall that $\tau(s, R)$ is defined by

$$\tau(s, R) = \exp([\log R - \log \log R](s-1)) = R^{s-1}(\log R)^{1-s}$$

and note that this implies

$$(18) \quad \tau(0, R) = \frac{\log R}{R}, \quad \tau(1, R) = 1, \quad \frac{\partial}{\partial s} \tau(s, R) = \tau(s, R)[\log R - \log \log R].$$

Thus, defining the auxiliary process

$$M_s^{R,W} = M_s := \frac{1}{R\sqrt{\log R - \log \log R}} [\bar{\Sigma}_\phi(Y(\tau(s, R), W_R)) - \bar{\Sigma}_\phi(Y(\tau(0, R), W_R))]$$

and using (4) with $W_R := RW$ and under variable substitution $s := \tau(u, R)$ and $t := s$ with left hand side variables corresponding to the notation of (4) and those on the right hand side to that used here, we see that, by (18),

$$(M_s)_{s \in [0,1]} \text{ and } \left(M_s^2 - \int_0^s \frac{\tau(u, R)}{R^2} A_{\phi^2}(Y(\tau(u, R), W_R)) du \right)_{s \in [0,1]}$$

are $\mathfrak{S}_{\tau(s, R)}$ -martingales. In particular, see once more [9, Paragraph I.8], the predictable quadratic variation process $\langle M \rangle_s$ is given by

$$(19) \quad \langle M \rangle_s = \int_0^s \frac{\tau(u, R)}{R^2} A_{\phi^2}(Y(\tau(u, R), W_R)) du, \quad s \in [0, 1].$$

Repeating the argument leading to (13) we see that

$$(20) \quad \lim_{R \rightarrow \infty} \frac{\tau(s, R)}{R^2} A_{\phi^2}(Y(\tau(s, R), W_R)) = V_W(\phi, \Lambda)$$

in probability, uniformly in $s \in [0, 1]$. Note that the uniformity in s comes, as in the case of (13), from the relation (12) implying that, in distribution, all instances of the left hand side for different values of s are just re-scalings of the same object $\tilde{R}^{-2} A_{\phi^2}(Y(1, W_{\tilde{R}}))$ for $\tilde{R} = R/\tau(s, R)$ and thus, in terms of the considered convergence in probability to a deterministic limit, we are just dealing with a single asymptotic statement. Consequently, by (20) and in full analogy to (15),

$$(21) \quad \lim_{R \rightarrow \infty} \langle M \rangle_s = \int_0^s V_W(\phi, \Lambda) du = s V_W(\phi, \Lambda) \quad \text{in probability.}$$

Thus, we are again in a position to apply the martingale functional central limit theorem [22, Thm 2.1] yielding the functional convergence in law, as $R \rightarrow \infty$, in $\mathcal{D}[0, 1]$ of $(M_s)_{s \in [0,1]}$ to the random process $(\sqrt{V_W(\phi, \Lambda)} \mathcal{W}_s)_{s \in [0,1]}$. Indeed, condition (ii.6) there is just (21), condition (ii.4) is trivial in view of (19), whereas the condition (ii.5) is verified by noting that, with probability one, $\mathcal{J}(M; 1) = \frac{1}{R\sqrt{\log R}} O(R \text{diam}(W)) = O(1/\sqrt{\log R})$, so that in particular $\lim_{R \rightarrow \infty} \mathbb{E} \mathcal{J}^2(M; 1) = 0$, as required.

Denoting now by $C^{R, W}$ the correction term $\frac{1}{R\sqrt{\log R}} \bar{\Sigma}_{\phi}(Y(\tau(0, R), W_R))$, such that

$$\mathcal{L}_s^{R, W} = C^{R, W} + \sqrt{\frac{\log R - \log \log R}{\log R}} M_s,$$

noting that $\log R - \log \log R \sim \log R$ and that, by the scaling property of STIT tessellations and by (7),

$$\begin{aligned} \text{Var}(C^{W, R}) &= O([R^{-2}(\log R)^{-1}][R^2/(\log R)^2][(\log R)^2(\log \log R)]) \\ &= O(\log \log R / \log R), \end{aligned}$$

we see that the processes M_s and $\mathcal{L}_s^{R,W}$ are asymptotically equivalent in $\mathcal{D}[0, 1]$, as $R \rightarrow \infty$. This completes the proof of Theorem 5. \square

REMARK 3. *In the context of proof of Theorem 5 it should be remarked that the ‘negligible correction term’ $C^{R,W}$ has its variance of order*

$$O(\log \log R / \log R)$$

and is thus indeed tending to 0, but extremely slowly. Consequently, although the Gaussian CLT holds for $\mathcal{L}_1^{R,W}$, it is quite natural to expect that the convergence rates are extremely slow, conjecturedly logarithmic. This is due to the fact that dimension 2 is the largest dimension (critical dimension) where the Gaussian limits are still present. In dimensions 3 and higher there is no Gaussian CLT and the ‘correction term’ analogous to $C^{R,W}$ will turn out order-determining rather than negligible, as shown by Theorem 3.

We turn now to the higher-dimensional situations. Even if in the formulation of Theorem 3 we have used the surface functional, we will show the statement in a more general context, where $\text{Vol}_{d-1}(Y(1, W_R))$ is replaced by a general cumulative facet functional $\Sigma_\phi(Y(1, W_R))$ satisfying (3).

We claim that the argument from the proof of Theorem 5 cannot be repeated for $d > 2$. Intuitively, this is due to the fact that for $d > 2$ the variance order of $\bar{\Sigma}_\phi(Y(1, W_R))$ is $O(R^{2(d-1)})$, see below, whereas the variance order of the increment $\bar{\Sigma}_\phi(Y(1, W_R)) - \bar{\Sigma}_\phi(Y(s_0, W_R))$, with some time instant $0 < s_0 < 1$, is $O(R^d)$ as seen from Theorem 4. Hence, for $d > 2$ we conclude that even the very first facets born in the MNW-cell-division process already bring a non-negligible contribution to the overall variance. Thus, we cannot split the whole STIT construction into the *warm-up phase* ($t \in [0, R^{-1} \log R]$ for $d = 2$) with negligible variance contribution and the *proper phase* unfolding already in a typical STIT environment. In fact, the claim is that the CLT does not hold for STIT surface functionals in dimension greater than 2!

PROOF OF THEOREM 3. Recall that we do not show this fact in full generality for all non-degenerate hyperplane measures Λ and all windows W , but restrict ourself to a particular case, where Λ is given by (2) and where $W = [0, 1]^d$. To see the non-Gaussianity, observe first that, by the scaling property of STIT tessellations,

$$(22) \quad R^{-(d-1)} \bar{\Sigma}_\phi(Y(1, W_R)) \stackrel{D}{=} \bar{\Sigma}_\phi(Y(R, W)),$$

which implies that the variance $\text{Var}(\Sigma_\phi(Y(1, W_R)))$ is of order $O(R^{2(d-1)})$. Indeed, this follows directly from the special form (3) of the facet functional

ϕ and the scaling relation $Y(1, W_R) \stackrel{D}{=} RY(R, W)$. Further, recall that by (4) the process $R \mapsto \bar{\Sigma}_\phi(Y(R, W))$ is a square-integrable martingale with absolutely continuous predictable quadratic variation process given in (5) and, moreover, by Proposition 2 we have

$$\begin{aligned} \text{Var}(\Sigma_\phi(Y(R, W))) &= \\ &= \int_{[W]} \zeta^2(\bar{\mathbf{n}}(H)) \int_{W \cap H} \int_{W \cap H} \frac{1 - \exp(-R\Lambda([xy]))}{\Lambda([xy])} dx dy \Lambda(dH), \end{aligned}$$

which is bounded uniformly in R . Consequently, by the martingale convergence theorem (cf. Corollary 7.22 in [7]) there exists a centred square-integrable random variable $\Xi(W)$ such that

$$\Xi(W) = \lim_{R \rightarrow \infty} \bar{\Sigma}_\phi(Y(R, W))$$

a.s. and in L^2 and, moreover,

$$\begin{aligned} \text{Var}(\Xi(W)) &= \lim_{R \rightarrow \infty} \text{Var}(\Sigma_\phi(Y(R, W))) \\ (23) \quad &= \int_{[W]} \zeta^2(\bar{\mathbf{n}}(H)) \int_{W \cap H} \int_{W \cap H} \frac{1}{\Lambda([xy])} dx dy \Lambda(dH). \end{aligned}$$

Using now (22) we readily conclude that

$$R^{-(d-1)} \bar{\Sigma}_\phi(Y(1, W)) \Longrightarrow \Xi(W)$$

as $R \rightarrow \infty$, where \Longrightarrow stands for convergence in distribution.

We show now that the variable $\Xi(W)$ cannot be Gaussian. To see it, consider the event \mathcal{E}_N , $N > 0$, that only hyperplanes orthogonal to e_1 have been born during the time $[0, 1]$ in the MNW-construction and that their number exceeds N . Observe that, in view of the special form (2) of the hyperplane measure Λ , $\mathbb{P}(\mathcal{E}_N) = \exp(-d) \sum_{k=N+1}^{\infty} \frac{1}{k!}$ and thus

$$(24) \quad \mathbb{P}(\mathcal{E}_N) = \exp(-\Theta(N \log N)),$$

where by $\Theta(\cdot)$ we mean a function bounded both from below and above by multiples of the argument. Further, given the fixed collection of all hyperplanes H_1, \dots, H_k , $k > N$, born at times between 0 and 1, on the event \mathcal{E}_N , we see that the conditional law of $\Xi(W)$ coincides with that of $k - d$ plus the sum of independent copies ξ_1, \dots, ξ_{k+1} of $\Xi(W_1), \dots, \Xi(W_{k+1})$ respectively, where W_j , $j = 1, \dots, k + 1$, are the parallelepipeds into which $W = [0, 1]^d$ is partitioned by H_1, \dots, H_k . More formally, we have the relation

$$\mathbb{P}(\Xi(W) > u | \mathcal{E}_{N,k}) = \mathbb{P}(k - d + (\xi_1 + \dots + \xi_{k+1}) > u), \quad u \in \mathbb{R},$$

where $\mathcal{E}_{N,k}$ is the event that exactly the hyperplanes H_1, \dots, H_k ($k > N$) orthogonal to e_1 are born within time $[0, 1]$. Note that the extra k above is the sum of the $(d-1)$ -volumes of $W \cap H_i$, whereas $-d = -\mathbb{E}\Sigma_\phi(Y(1, W))$ is the centering term.

Since $\text{Var}(\xi_1 + \dots + \xi_{k+1}) = \sum_{j=1}^{k+1} \text{Var}(\Xi(W_j))$, which is bounded from above by $\text{Var}(\Xi(W))$ in view of (23), by Chebyshev's inequality we get

$$\begin{aligned} \mathbb{P}(\xi_1 + \dots + \xi_{k+1} \geq -2\sqrt{\text{Var}(\Xi(W))}) &\geq 1 - \frac{\text{Var}(\xi_1 + \dots + \xi_{k+1})}{4 \text{Var}(\Xi(W))} \\ &\geq 1 - \frac{1}{4} = \frac{3}{4}. \end{aligned}$$

Thus, in view of (24) we end up with

$$\mathbb{P}(\Xi(W) > N) \geq \frac{3}{4} \mathbb{P}\left(\mathcal{E}_{N+2\sqrt{\text{Var}(\Xi(W))+d}}\right) = \exp(-\Theta(N \log N)).$$

Since Gaussian variables exhibit tail decay of the order $\exp(-\Theta(N^2))$, the random variable $\Xi(W)$ cannot be Gaussian. \square

Acknowledgment. The second author is indebted to Werner Nagel and Matthias Reitzner for interesting discussions, suggestions and remarks. Moreover, the suggestions and remarks of an anonymous referee were helpful in improving the presentation and the style of the paper.

REFERENCES

- [1] BAUMSTARK, V. AND LAST, G. (2009). Gamma distributions for stationary Poisson flat processes, *Adv. Appl. Probab.* **41** 911–939.
- [2] FAVIS, W. AND WEISS, V. (1998). Mean values of weighted cells of stationary Poisson hyperplane tessellations in \mathbb{R}^d , *Math. Nachr.* **193** 37–48.
- [3] GEORGII, H.-O. (1988). *Gibbs measures and phase transitions*, de Gruyter, Berlin.
- [4] HEINRICH, L. (2009). Central limit theorems for motion-invariant Poisson hyperplanes in expanding convex windows, *Rendiconti del Circolo Matematico di Palermo Series II, Suppl.* **81** 187–212.
- [5] HEINRICH, L. AND MUCHE, L. (2008). Second-order properties of the point process of nodes in a stationary Voronoi tessellation, *Math. Nachr.* **281** 350 – 375.
- [6] HUG, D. AND SCHNEIDER, R. (2007). Asymptotic shapes of large cells in random tessellations, *Geom. Funct. Anal.* **17** 156–191.
- [7] KALLENBERG, O. (2002). *Foundations of Modern Probability*, Springer, New York.

- [8] LACHIÈZE-REY, R. (2011). Mixing properties for STIT tessellations, *Adv. Appl. Probab.* **43** 40–48.
- [9] LIPSTER, R.S. AND SHIRYAYEV, A.N. (1989). *Theory of Martingales*, Kluwer Academic Publishers, Dordrecht.
- [10] MECKE, J.; NAGEL, W. AND WEISS, V. (2008). A global construction of homogeneous random planar tessellations that are stable under iteration, *Stochastics* **80** 51–67.
- [11] MECKE, J.; NAGEL, W. AND WEISS, V. (2008). The iteration of random tessellations and a construction of a homogeneous process of cell division, *Adv. Appl. Probab.* **40**, 49–59.
- [12] NAGEL, W. AND WEISS, V. (2003). Limits of sequences of stationary planar tessellations, *Adv. Appl. Probab.* **35** 123–138.
- [13] NAGEL, W. AND WEISS, V. (2005). Crack STIT tessellations: characterization of stationary random tessellations stable with respect to iteration, *Adv. Appl. Probab.* **37** 859–883.
- [14] SCHNEIDER, R. AND WEIL, W. (2008). *Stochastic and Integral Geometry*, Springer, Berlin.
- [15] SCHREIBER, T. (2010). Limit theorems in stochastic geometry, In *New Perspectives in Stochastic Geometry*, edited by W.S. Kendall and I. Molchanov, Oxford University Press, Oxford.
- [16] SCHREIBER, T. AND THÄLE, C. (2010). Second-order properties and central limit theory for the vertex process of iteration infinitely divisible and iteration stable random tessellations in the plane, *Adv. Appl. Probab.* **42** 913–935.
- [17] SCHREIBER, T. AND THÄLE, C. (2011). Intrinsic volumes of the maximal polytope process in higher dimensional STIT tessellations, *Stoch. Proc. Appl.* **121** 989–1012.
- [18] SCHREIBER, T. AND THÄLE, C. (2010). Typical geometry, second-order properties and central limit theory for iteration stable tessellations, *arXiv: 1001.0990 [math.PR]*.
- [19] SCHREIBER, T. AND THÄLE, C. (2011). Geometry of iteration stable tessellations: Connection with Poisson hyperplanes, *arXiv: 1103.3958 [math.PR]*.
- [20] SCHREIBER, T. AND THÄLE, C. (2011). Second-order theory for iteration stable tessellations, *arXiv: 1103.3959 [math.PR]*.
- [21] STOYAN, D.; KENDALL, W. AND MECKE, J. (1995). *Stochastic Geometry and its Applications*, 2nd ed., Wiley, Chichester.
- [22] WHITT, W. (2007). Proofs of the martingale FCLT, *Probability Surveys* **4** 268–302.

TOMASZ SCHREIBER
FACULTY OF MATHEMATICS AND COMPUTER SCIENCE
NICOLAUS COPERNICUS UNIVERSITY
TORUŃ, POLAND

UNIVERSITÄT OSNABRÜCK
INSTITUT FÜR MATHEMATIK
ALBRECHTSTRASSE 28A
D-49076 OSNABRÜCK
GERMANY
E-MAIL: cthale@uos.de