

The first order correction to harmonic measure for random walks of rotationally invariant step distribution

Longmin Wang, Kainan Xiang and Lang Zou*
 School of Mathematical Sciences, LPMC, Nankai University
 Tianjin City, 300071, P. R. China
 Emails: wanglm@nankai.edu.cn (Wang)
 kainanxiang@nankai.edu.cn (Xiang)
 lang.zou.proba@gmail.com (Zou)

Abstract

Let $D \subset \mathbb{R}^d$ ($d \geq 2$) be an open simply-connected bounded domain with smooth boundary ∂D and $\mathbf{0} = (0, \dots, 0) \in D$. Fix any rotationally invariant probability μ on closed unit ball $\{z \in \mathbb{R}^d : |z| \leq 1\}$ with $\mu(\{\mathbf{0}\}) < 1$. Let $\{S_n^\mu\}_{n=0}^\infty$ be the random walk with step-distribution μ starting at $\mathbf{0}$. Denote by $\omega_\delta(\mathbf{0}, dz; D)$ the discrete harmonic measure for $\{\delta S_n^\mu\}_{n=0}^\infty$ ($\delta > 0$) exiting from D , which is viewed as a probability on ∂D by projecting suitably the first exiting point to ∂D . Denote by $\omega(\mathbf{0}, dz; D)$ the harmonic measure for the d -dimensional standard Brownian motion exiting from D . Then in the weak convergence topology,

$$\lim_{\delta \rightarrow 0} \frac{1}{\delta} [\omega_\delta(\mathbf{0}, dz; D) - \omega(\mathbf{0}, dz; D)] = c_\mu \rho_D(z) |dz|,$$

where $\rho_D(\cdot)$ is a smooth function depending on D but not on μ , c_μ is a constant depending only on μ , and $|dz|$ is the Lebesgue measure with respect to ∂D . Additionally, $\rho_D(z)$ is determined by the following equation: For any smooth function g on ∂D ,

$$\int_{\partial D} g(z) \rho_D(z) |dz| = \int_{\partial D} \frac{\partial f}{\partial \mathbf{n}_z}(z) H_D(\mathbf{0}, z) |dz|,$$

where f is the harmonic function in D with boundary values given by g , $H_D(\mathbf{0}, z)$ is the Poisson kernel and derivative $\frac{\partial f}{\partial \mathbf{n}_z}$ is with respect to the inward unit normal \mathbf{n}_z at $z \in \partial D$.

1 Introduction

We study in this paper the universality for the first order correction between discrete and continuous harmonic measures in \mathbb{R}^d with $d \geq 2$, which was initiated by Kennedy [11] and Jiang & Kennedy [10] in \mathbb{R}^2 .

To begin, denote $\mathbf{0} = (0, \dots, 0) \in \mathbb{R}^d$, $x = (x_1, \dots, x_d)$ for any $x \in \mathbb{R}^d$. Let $\{X_i\}_{i=0}^\infty$ be an i.i.d sequence of random variables in \mathbb{R}^d with common distribution μ such that

$$\begin{cases} \mathbb{E}[X_i] = \mathbf{0}, & \mathbb{E}[X_i^{(n)} X_i^{(m)}] = 0, \quad n \neq m, \\ \mathbb{E}\left[\left(X_i^{(1)}\right)^2\right] = \dots = \mathbb{E}\left[\left(X_i^{(d)}\right)^2\right] = \kappa \in (0, \infty), \end{cases} \quad (1.1)$$

where $X_i = (X_i^{(1)}, \dots, X_i^{(d)})$. For any $s_0 \in \mathbb{R}^d$, we define the random walk $\{S_n^\mu\}_{n \geq 0}$ with step distribution μ starting at s_0 by

$$S_0^\mu = s_0, \quad S_n^\mu = s_0 + X_1 + \dots + X_n, \quad n \geq 1.$$

The rescaled process $\left\{ \delta \left(S_{[\delta^{-2}t]}^\mu - s_0 \right) \right\}_{t \geq 0}$ converges in law to $\{B(\kappa t)\}_{t \geq 0}$ as $\delta \rightarrow 0$, where $[x]$ is the integer part of $x \in \mathbb{R}$ and $\{B(t)\}_{t \geq 0}$ is the d -dimensional standard Brownian motion in \mathbb{R}^d starting at $\mathbf{0}$.

* Correspondence to Lang Zou. Email: lang.zou.proba@gmail.com.

The project is supported partially by CNNSF (No. 11671216).

MSC2010 subject classifications. 60J10, 05C81, 60D05, 60G17.

Key words and phrases. Harmonic measure; Discrete harmonic measure; Random walk; First order correction.

Let $D \subset \mathbb{R}^d$ be an open simply-connected bounded domain with smooth boundary ∂D and $\mathbf{0} \in D$. It is known that D is regular ([7] p.27 and [12] pp.245-250), in the sense that,

$$\mathbb{P}_z\{\tau_D = 0\} = 1, \quad \forall z \in \partial D;$$

where $\tau_D = \inf\{t \geq 0 : B(t) \notin D\}$ and \mathbb{P}_z is the law of $\{B(t)\}_{t \geq 0}$ starting from z . Let $\omega(x, dz; D)$ be the continuous harmonic measure for $\{B(t)\}_{t \geq 0}$ exiting from D when starting at $x \in D$, that is,

$$\omega(x, dz; D) = \mathbb{P}_x(B(\tau_D) \in dz).$$

It is known that for fixed $x \in D$, $\omega(x, dz; D)$ is absolutely continuous with respect to $|dz|$, the Lebesgue measure on ∂D . More precisely,

$$\omega(x, dz; D) = H_D(x, z) |dz|,$$

where $H_D(x, z)$ is the Poisson kernel in D and may be defined as the derivative of the Green function $G_D(x, z)$ in the direction \mathbf{n}_z , the inward unit normal at $z \in \partial D$, i.e.,

$$H_D(x, z) = \frac{\partial G_D(x, z)}{\partial \mathbf{n}_z}.$$

Since ∂D is smooth, we have that $H_D(x, z)$ is smooth in $z \in \partial D$ (c.f. [9] and [7]). For more details on harmonic measures please refer to [1, 5, 12, 15].

Now we back to the discrete setting. Without loss of generality, in the rest of this paper we will always assume $S_0^\mu = s_0 = \mathbf{0}$, unless otherwise specified. Let

$$T_D = T_D(\delta) = \inf\{n \geq 0 : \delta S_n^\mu \notin D\}.$$

Define the discrete harmonic measure $\hat{\omega}_\delta(\mathbf{0}, dz; D)$ for $\{\delta S_n^\mu\}_{n \geq 0}$ exiting from D by

$$\hat{\omega}_\delta(\mathbf{0}, \Gamma; D) = \mathbb{P}\left(\overline{\delta S_{T_D}^\mu} \in \Gamma\right), \quad \forall \text{ measurable } \Gamma \subseteq \partial D, \quad (1.2)$$

where $\overline{\delta S_{T_D}^\mu}$ is the point on ∂D with the smallest distance to $\delta S_{T_D}^\mu$. Note that the choice for $\overline{\delta S_{T_D}^\mu}$ is unique when δ is sufficiently small. Indeed, the following map

$$\varphi_a : z \in \partial D \rightarrow z - a\mathbf{n}_z \in \partial D(a) := \{y - a\mathbf{n}_y : y \in \partial D\}$$

is smooth for any $a \in [0, 1]$, and $\varphi_a(z)$ is smooth in $(z, a) \in \partial D \times [0, 1]$; and the Jacobian determinant $J_a(z)$ of φ_a is smooth in $z \in \partial D$ for any $a \in [0, 1]$ and $J_a(z)$ is smooth in $(z, a) \in \partial D \times [0, 1]$. Thus from $J_0(z) \equiv 1$, for small enough $a > 0$, $J_a(z) > 0$, $z \in \partial D$, namely $\varphi_a(\cdot)$ is injective; which implies that for any $x \notin \partial D$ with $\text{dist}(x, \partial D) \leq a$, there is a unique $y \in \partial D$ satisfying $\text{dist}(x, \partial D) = |x - y|$. Here $\text{dist}(x, \partial D) := \inf_{z \in \partial D} |x - z|$ is the distance between x and ∂D .

Note that $\hat{\omega}_\delta(\mathbf{0}, \cdot; D)$ converges weakly to $\omega(\mathbf{0}, \cdot; D)$ as $\delta \downarrow 0$. Our primary motivation is to study the following conjecture of universality problem which was pioneered by Kennedy in [11].

$$\lim_{\delta \rightarrow 0} \frac{1}{\delta} [\hat{\omega}_\delta(\mathbf{0}, dz; D) - \omega(\mathbf{0}, dz; D)] = c_\mu \rho_D(z) |dz| \quad (\text{in the weak convergence topology}),$$

where $c_\mu > 0$ is a constant depending on μ but not on D , and $\rho_D(z)$ is a measurable function on ∂D independent of μ .

The above universality may not hold for the random walk $\{\delta S_n^\mu\}_{n \geq 0}$ whose μ is not rotationally invariant. The reason is that the standard d -dimensional Brownian motion starting at $\mathbf{0}$ is rotationally invariant. Thus it is natural to use the following $\omega_\delta(\mathbf{0}, \cdot; D)$ to replace $\hat{\omega}_\delta(\mathbf{0}, \cdot; D)$:

$$\omega_\delta(\mathbf{0}, dz; D) = \int_{O(d)} \hat{\omega}_{\delta, \alpha}(\mathbf{0}, dz; D) d\tilde{m}(\alpha);$$

where \tilde{m} is the normalized Haar measure on $O(d)$, the group of $d \times d$ orthogonal matrices, and $\hat{\omega}_{\delta, \alpha}(\mathbf{0}, dz; D)$ is the image of $\hat{\omega}_\delta(\mathbf{0}, dz; D)$ under rotation $\alpha \in O(d)$. For a detailed discussion on the above replacement in \mathbb{R}^2 , refer to Kennedy [11].

Now we restate the universality problem as follows.

Conjecture 1.1 Assume that $D \subset \mathbb{R}^d$ is an open simply-connected bounded domain with $\mathbf{0} \in D$ and ∂D is smooth. Let g be any smooth function on ∂D . Then

$$\lim_{\delta \rightarrow 0} \frac{1}{\delta} \left(\int_{\partial D} g(z) \omega_\delta(\mathbf{0}, dz; D) - \int_{\partial D} g(z) \omega(\mathbf{0}, dz; D) \right) = c_\mu \int_{\partial D} g(z) \rho_D(z) |dz|, \quad (1.3)$$

where $c_\mu > 0$ is a constant depending only on μ , and $\rho_D(z)$ is a measurable function on ∂D independent of μ .

Moreover, $\rho_D(z)$ is determined by the following equation:

$$\int_{\partial D} g(z) \rho_D(z) |dz| = \int_{\partial D} \frac{\partial f}{\partial \mathbf{n}_z}(z) H_D(\mathbf{0}, z) |dz|, \quad (1.4)$$

where f is the harmonic function in D with boundary values given by g , and derivative $\frac{\partial f}{\partial \mathbf{n}_z}$ is with respect to the inward unit normal \mathbf{n}_z at $z \in \partial D$.

For any $r > 0$ and $z_0 \in \mathbb{R}^d$, write

$$\mathbb{B}^d(z_0, r) = \{|z - z_0| < r : z \in \mathbb{R}^d\}, \quad \overline{\mathbb{B}^d(z_0, r)} = \{|z - z_0| \leq r : z \in \mathbb{R}^d\}.$$

Let $\partial \mathbb{B}^d(z_0, r)$ be the boundary of $\mathbb{B}^d(z_0, r)$. For convenience, put

$$\mathbb{B}_r^d := \mathbb{B}^d(\mathbf{0}, r), \quad \mathbb{B}^d := \mathbb{B}^d(\mathbf{0}, 1).$$

When μ is the uniform probability measure on \mathbb{B}^2 , Jiang and Kennedy [10] proved Conjecture 1.1 in \mathbb{R}^2 . We attribute the above conjecture to [11] and [10], though in [11] this conjecture was stated for three kinds of random walk models (simple random walk, the nearest neighbor random walk not allowed to backtrack, the smart kinetic walk) on square, triangular and hexagonal planar lattices; and we are responsible for any possible mistake.

In the rest of the paper, we assume that μ is a rotationally invariant probability measure on $\overline{\mathbb{B}^d}$ such that $\mu(\{\mathbf{0}\}) < 1$. Under this setting, there is a probability measure ν on $[0, 1]$ with $\nu(\{0\}) < 1$ such that

$$\int f d\mu = \int_0^1 \int_{\mathbb{S}^{d-1}} f(r\alpha) d\nu(r) dm(\alpha) \quad (1.5)$$

for all positive measurable function f , where m is the uniform distribution on the unit sphere \mathbb{S}^{d-1} . Hence, the rotational invariant property of μ implies that

$$\omega_\delta(\mathbf{0}, \cdot; D) = \hat{\omega}_\delta(\mathbf{0}, \cdot; D) = \hat{\omega}_{\delta, \alpha}(\mathbf{0}, \cdot; D).$$

Let

$$\mathbb{H}^d = \{(x_1, x_2, \dots, x_d) \in \mathbb{R}^d : x_d > 0\}, \quad \partial \mathbb{H}^d = \{(x_1, x_2, \dots, x_d) \in \mathbb{R}^d : x_d = 0\},$$

and $T_{\mathbb{H}^d} := \inf \{n \geq 0, S_n^\mu \notin \mathbb{H}^d\}$. Define $h^\mu(\ell)$ on $[0, 1]$ by

$$h^\mu(\ell) = \int_{[\ell, 1]} \frac{\Gamma(d/2)}{\sqrt{\pi} \Gamma(\frac{d-1}{2})} \left[\frac{(r^2 - \ell^2)^{(d-1)/2}}{(d-1)r^{d-2}} + {}_2F_1 \left(\frac{1}{2}, \frac{3-d}{2}; \frac{3}{2}; \frac{\ell^2}{r^2} \right) \frac{\ell^2}{r} \right] d\nu(r) - \frac{\ell}{2} \nu([\ell, 1]), \quad (1.6)$$

where ${}_2F_1(a, b; c; z)$ is the hypergeometric function given by

$${}_2F_1(a, b; c; z) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{z^n}{n!},$$

and $(x)_n = x(x+1) \cdots (x+n-1)$ is the Pochhammer symbol. For hypergeometric functions, see [6]. Let

$$c_\mu = \frac{2}{\kappa} \int_0^1 \left(\ell + \mathbb{E}^\ell \left[V \left(S_{T_{\mathbb{H}^d}}^\mu \right) \right] \right) h^\mu(\ell) d\ell, \quad (1.7)$$

where \mathbb{E}^ℓ is the expectation with respect to the distribution of $\{S_n^\mu\}_{n \geq 0}$ starting at $s_0 = (0, \dots, 0, \ell) \in \mathbb{H}^d$, and

$$V \left(S_{T_{\mathbb{H}^d}}^\mu \right) = \left| S_{T_{\mathbb{H}^d}}^\mu - \overline{S_{T_{\mathbb{H}^d}}^\mu} \right|.$$

We state our main result as follows.

Theorem 1.2 *Assume that μ is a rotationally invariant probability on $\overline{\mathbb{B}^d}$ and $\mu(\{\mathbf{0}\}) < 1$. Then Conjecture 1.1 holds for constant c_μ specified in (1.7) and smooth function $\rho_D(z)$ specified by*

$$\rho_D(z) = \lim_{\delta \rightarrow 0} \frac{1}{\delta} \left\{ \int_{\partial D} H_D(\mathbf{0}, \zeta) H_D(z + \delta \mathbf{n}_z, \zeta) |\mathrm{d}\zeta| - H_D(\mathbf{0}, z) \right\}, \quad z \in \partial D. \quad (1.8)$$

The limit of (1.8) does exist, see the proof of Lemma 2.12.

Remark 1.3 (i) *The most difficult situation for Conjecture 1.1 is the case that μ is not a rotationally invariant probability measure. Refer to [11] and [4] for several very interesting discrete random walks.*

(ii) *As pointed out by Kennedy [11], there is another very natural way to define the ‘exit’ point in ∂D when the random walk exits D : By linearly interpolating between the steps of the random walk so that it becomes a piece-wise linear curve in \mathbb{R}^d , we can consider the first point where this curve intersects ∂D as the exit point. In this setting, Theorem 1.2 also holds.*

The paper is organized as follows. In Section 2, we recall firstly some preliminary facts on harmonic measures and random walks; then after giving a series of lemmas on discrete and continuous harmonic measures, we prove Theorem 1.2. In Section 3, we take several examples for Theorem 1.2.

2 Proof of Theorem 1.2

In Subsection 2.1, we recall some preliminary knowledge. Then in Subsection 2.2, we prove a series of lemmas which will be used to prove Theorem 1.2. Finally in Subsection 2.3, we finish the proof of Theorem 1.2.

Recall in \mathbb{R}^2 , by using the conformal mapping from D to \mathbb{B}^2 , Kennedy computed $\rho_D(z)$ in [11] Section 3 and Jiang & Kennedy deduced a formula for $\rho_D(z)$ in [10] Proposition 3. In this paper, we have found a more general approach without resorting the conformal mapping to compute explicitly the function $\rho_D(z)$ for d -dimension ($d \geq 2$) domain D . This together with Jiang and Kennedy [10] lead that we can extend main result of [10] to higher dimensional \mathbb{R}^d and any non-degenerate rotationally invariant step-distribution on $\overline{\mathbb{B}^d}$. Our proof is similar to that of [10], but we need some new insights in higher dimensional \mathbb{R}^d which scatter in proofs of lemmas in Subsection 2.2.

2.1 Preliminaries

For small $\delta > 0$, let

$$D_2 = \{z \in D : \mathrm{dist}(z, \partial D) < \delta\}, \quad D_3 = \{z \in \mathbb{R}^d \setminus D : \mathrm{dist}(z, \partial D) < \delta\}, \quad (2.1)$$

$$D_1 = D \setminus D_2, \quad D_+ = D \cup D_3. \quad (2.2)$$

For any continuous function g on ∂D , consider the following Dirichlet problem:

$$\begin{cases} \Delta f(z) = 0, & z \in D, \\ f(z) = g(z), & z \in \partial D. \end{cases} \quad (2.3)$$

Recall that D is regular. The unique solution to (2.3) can be written as

$$f(z) = \int_{\partial D} g(\xi) \omega(z, \mathrm{d}\xi; D). \quad (2.4)$$

Furthermore, if g is smooth on ∂D , then $f \in C^\infty(\overline{D})$ ([7] p.111 Theorem 6.19) and therefore f has a C^∞ -extension to a larger domain \overline{D}_+ , which is still denoted by f by abusing notations.

Clearly the generator Δ_δ for the random walk $\{\delta S_n^\mu\}_{n \geq 0}$ is given by

$$\Delta_\delta f(z) = \int_{\mathbb{B}^d} [f(z + \delta\xi) - f(z)] d\mu(\xi), \quad (2.5)$$

for any bounded measurable function f on \mathbb{R}^d . Consider the following discrete Dirichlet problem:

$$\begin{cases} \Delta_\delta f_\delta(z) = 0, & z \in D, \\ f_\delta(z) = g(\tilde{z}), & z \in D_3; \end{cases} \quad (2.6)$$

where D_3 is given by (2.1), and $\tilde{z} \in \partial D$ satisfies $|\tilde{z} - z| = \min\{|\zeta - z| : \zeta \in \partial D\}$. It is easy to see that the function f_δ defined by

$$f_\delta(z) = \int_{\partial D} g(\xi) \omega_\delta(z, d\xi; D) \quad (2.7)$$

is the unique solution to (2.6) (the uniqueness follows from the maximum principle).

Let $\mathbb{P}^{x, \delta}$ be the law of $\{\delta S_n^\mu\}_{n \geq 0}$ with $\delta S_0^\mu = x$, and $\mathbb{E}^{x, \delta}$ the corresponding expectation. Define the following probability on \mathbb{B}^d : For any measurable subset A of \mathbb{B}^d ,

$$\widehat{\mu}(A) = \frac{1}{1 - \mu(\{\mathbf{0}\})} \mu(A \setminus \{\mathbf{0}\}).$$

Note that the random walks $\{\delta S_n^\mu\}_{n \geq 0}$ and $\{\delta S_n^{\widehat{\mu}}\}_{n \geq 0}$ have the same discrete harmonic measure. Thus in the rest of this paper we assume that

$$\mu(\{\mathbf{0}\}) = 0.$$

Notice that μ is rotationally invariant on $\overline{\mathbb{B}^d}$ ($d \geq 2$) with $\mu(\{\mathbf{0}\}) = 0$. By [17], the k -fold convolution

$$\underbrace{\mu * \mu * \cdots * \mu}_k$$

with $k \geq 2$ is absolutely continuous with respect to the d -dimensional Lebesgue measure. Define the transition probability density for the random walk $\{\delta S_n^\mu\}_{n \geq 0}$:

$$\begin{aligned} p_\delta(0, x, y) &= \delta(x, y), \\ p_\delta(n, x, y) &= \lim_{\epsilon \rightarrow 0} \frac{\mathbb{P}^{x, \delta}(|\delta S_n^\mu - y| \leq \epsilon)}{2\pi^{d/2}\epsilon^d/(d\Gamma(d/2))}, \quad n \geq 2, \end{aligned} \quad (2.8)$$

where $\delta(x, y)$ is the Dirac delta function giving unit mass to x . In general, for $n = 1$, the limit above does not exist and so $p_\delta(1, x, y)$ may be a distribution rather than a function.

Likewise, define the transition probability density for $\{\delta S_n^\mu\}_{n \geq 0}$ killed on exiting D as follows: For any $x \in D$ and $y \in \mathbb{R}^d$,

$$\begin{aligned} p_{D, \delta}(0, x, y) &= \delta(x, y), \\ p_{D, \delta}(n, x, y) &= \lim_{\epsilon \rightarrow 0} \frac{\mathbb{P}^{x, \delta}(|\delta S_n^\mu - y| \leq \epsilon, n < T_D)}{2\pi^{d/2}\epsilon^d/(d\Gamma(d/2))}, \quad n \geq 2. \end{aligned}$$

Here $p_{D, \delta}(n, x, y)$ does exist by (2.8) for $n \geq 2$.

2.2 Some lemmas

By the Markov property for $\{\delta S_n^\mu\}_{n \geq 0}$, we may rewrite $p_{D, \delta}(n, x, y)$ as

$$p_{D, \delta}(n, x, y) = \int_{\mathbb{B}^d} p_{D, \delta}(n-1, x, y + \delta\xi) d\mu(\xi), \quad x, y \in D, \quad n \geq 3. \quad (2.9)$$

The killed discrete Green function is defined by

$$G_\delta(x, y) = \sum_{n=0}^{\infty} p_{D, \delta}(n, x, y), \quad x \in D, \quad y \in D \setminus \overline{\mathbb{B}^d(x, 2\delta)},$$

where $p_{D, \delta}(1, x, y)$ is viewed as 0 due to $|y - x| > 2\delta$. In fact in the above definition, $p_{D, \delta}(n, x, y)$ with $n \leq 2$ make no contribution to $G_\delta(x, y)$. Recall D_3 from (2.1).

Lemma 2.1 For any fixed $x \in D$,

$$\begin{cases} \Delta_\delta G_\delta(x, y) = 0, & y \in D \setminus \overline{\mathbb{B}^d(x, 2\delta)}, \\ G_\delta(x, y) = 0, & y \in D_3. \end{cases} \quad (2.10)$$

Proof. The boundary condition is obvious. By (2.9) and the Fubini theorem, for any $y \in D \setminus \overline{\mathbb{B}^d(x, 2\delta)}$,

$$\begin{aligned} G_\delta(x, y) &= \sum_{n=0}^2 p_{D,\delta}(n, x, y) + \sum_{n=3}^{\infty} \int_{\mathbb{B}^d} p_{D,\delta}(n-1, x, y + \delta\xi) \, d\mu(\xi) \\ &= \int_{\mathbb{B}^d} \sum_{n=3}^{\infty} p_{D,\delta}(n-1, x, y + \delta\xi) \, d\mu(\xi) = \int_{\mathbb{B}^d} G_\delta(x, y + \delta\xi) \, d\mu(\xi), \end{aligned}$$

where we used the facts that $p_{D,\delta}(0, x, y + \delta\xi) = p_{D,\delta}(1, x, y + \delta\xi) = 0$ for $y \in D \setminus \overline{\mathbb{B}^d(x, 2\delta)}$ and $\xi \in \overline{\mathbb{B}^d}$. Hence,

$$\Delta_\delta G_\delta(x, y) = \int_{\mathbb{B}^d} G_\delta(x, y + \delta\xi) \, d\mu(\xi) - G_\delta(x, y) = 0, \quad y \in D \setminus \overline{\mathbb{B}^d(x, 2\delta)}.$$

□

Similarly to [10, Lemma 3], we can prove

Lemma 2.2 In the settings of Theorem 1.2, for smooth function g on ∂D ,

$$f_\delta(\mathbf{0}) - f(\mathbf{0}) = \int_{D_2} G_\delta(\mathbf{0}, z) \Delta_\delta f(z) \, dz. \quad (2.11)$$

Define potential kernel $a_\delta(x)$ for the random walk $\{\delta S_n^\mu\}_{n \geq 0}$ by

$$a_\delta(x) = \sum_{n=3}^{\infty} p_\delta(n, \mathbf{0}, x), \quad x \in \mathbb{R}^d \quad (2.12)$$

for $d \geq 3$ and by

$$a_\delta(x) = \sum_{n=3}^{\infty} [p_\delta(n, \mathbf{0}, \mathbf{0}) - p_\delta(n, \mathbf{0}, x)], \quad x \in \mathbb{R}^2 \quad (2.13)$$

for $d = 2$.

For convenience, when $\delta = 1$, we define

$$a(x) := a_1(x), \quad p(n, x, y) := p_1(n, x, y).$$

Lemma 2.3 Assume $d \geq 3$. Then $a(x)$ is well-defined, and

$$a(x) = \frac{\Gamma(\frac{d-2}{2})}{2\kappa\pi^{d/2}} \frac{1}{|x|^{d-2}} + O(|x|^{-d}), \quad |x| \rightarrow \infty, \quad (2.14)$$

where the constant in big O term only depends on μ .

Proof. Let $X = (X^{(1)}, X^{(2)}, \dots, X^{(d)})$ be a random variable with distribution μ , and $i = \sqrt{-1}$. Let

$$\phi(\theta) = \mathbb{E}(e^{i\theta \cdot X}) = \mathbb{E}\left(\exp\left\{i \sum_{j=1}^d \theta_j X^{(j)}\right\}\right)$$

be the characteristic function of X , where $\theta = (\theta_1, \dots, \theta_d) \in \mathbb{R}^d$. By the rotational invariance of μ , for any $n \geq 1$,

$$\mathbb{E}[(\theta \cdot X)^n] = \mathbb{E}\left[\left\{(0, \dots, 0, |\theta|) \cdot (X^{(1)}, \dots, X^{(d)})\right\}^n\right] = |\theta|^n \mathbb{E}\left[\left(X^{(d)}\right)^n\right];$$

and $\phi(\theta)$ is rotationally invariant in θ , namely a function in $|\theta|$:

$$\phi(\theta) = \chi(|\theta|). \quad (2.15)$$

Note $|\theta \cdot X| \leq |\theta|$. Then as $\theta \rightarrow \mathbf{0}$,

$$\mathbb{E} \left[\sum_{n=3}^{\infty} \frac{(-1)^n}{(2n)!} (\theta \cdot X)^{2n} \right] = \sum_{n=3}^{\infty} \frac{(-1)^n}{(2n)!} \mathbb{E} [(\theta \cdot X)^{2n}] = \sum_{n=3}^{\infty} \frac{(-1)^n}{(2n)!} |\theta|^{2n} \mathbb{E} \left[(X^{(d)})^{2n} \right] = O(|\theta|^6).$$

Since the Taylor series at 0 of $\cos(x)$ is $\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$, we have that

$$\begin{aligned} \phi(\theta) &= \mathbb{E} (\cos(\theta \cdot X) + i \sin(\theta \cdot X)) = \mathbb{E} (\cos(\theta \cdot X)) \\ &= \mathbb{E} \left(1 - \frac{(\theta \cdot X)^2}{2} + \frac{(\theta \cdot X)^4}{24} + \sum_{n=3}^{\infty} \frac{(-1)^n}{(2n)!} (\theta \cdot X)^{2n} \right) \\ &= 1 - \frac{\kappa|\theta|^2}{2} + \frac{1}{24} \mathbb{E} \left[(X^{(d)})^4 \right] |\theta|^4 + O(|\theta|^6), \quad \theta \rightarrow \mathbf{0}. \end{aligned} \quad (2.16)$$

From (2.27) in [8], the function $\chi(\cdot)$ has the following form:

$$\chi(q) = \Gamma(d/2) \int_0^1 \left(\frac{2}{rq} \right)^{d/2-1} J_{d/2-1}(rq) \, d\nu(r), \quad q \geq 0. \quad (2.17)$$

Here $J_v(z)$ is the Bessel function of the first kind of order v (see [16]), which is defined by the series

$$J_v(z) = \left(\frac{1}{2}z \right)^v \sum_{k=0}^{\infty} \frac{(-\frac{1}{4}z^2)^k}{k! \Gamma(v+k+1)}.$$

Clearly, for any $n \geq 3$, $p(n, \mathbf{0}, x)$ is rotationally invariant in x . Let

$$p(n, \mathbf{0}, x) = \psi_n(|x|).$$

Then by (2.34) in [8],

$$\psi_n(\rho) = \frac{2}{(4\pi)^{d/2}} \int_0^{\infty} \left(\frac{2}{q\rho} \right)^{d/2-1} J_{d/2-1}(q\rho) q^{d-1} \chi^n(q) \, dq, \quad (2.18)$$

where $\chi(q)$ is given by (2.17). By (2.15) and (2.16), and $\ln(1+\epsilon) = \epsilon + O(\epsilon^2)$ ($\epsilon \rightarrow 0$), we get that

$$\chi^n(q) = e^{n \ln(\chi(q))} \sim e^{-\frac{\kappa}{2} n q^2}, \quad \text{as } q \rightarrow 0. \quad (2.19)$$

Recall the following definite integral in [16] p.394:

$$\int_0^{\infty} q^{v+1} J_v(\beta q) e^{-\alpha^2 q^2} \, dq = \frac{\beta^v}{(2\alpha^2)^{v+1}} e^{-\beta^2/(4\alpha^2)}. \quad (2.20)$$

Thus the large n behaviour of the integral in (2.18) is governed by the small q behaviour of $\chi(q)$, and by (2.19)-(2.20),

$$p(n, \mathbf{0}, x) = \psi_n(|x|) \sim \frac{1}{(2\pi\kappa n)^{d/2}} e^{-\frac{|x|^2}{2n\kappa}}. \quad (2.21)$$

For any $n \geq 1$ and $x \in \mathbb{R}^d$, let

$$\bar{p}(n, \mathbf{0}, x) = \frac{1}{(2\pi\kappa n)^{d/2}} e^{-\frac{|x|^2}{2n\kappa}}. \quad (2.22)$$

By [14, Lemma 4.3.2], we have that for any $b > 1$, as $r \rightarrow \infty$,

$$\sum_{n=3}^{\infty} n^{-b} e^{-r/n} = \frac{\Gamma(b-1)}{r^{b-1}} + O\left(\frac{1}{r^{b+1}}\right). \quad (2.23)$$

Plugging $b = d/2$, $r = \frac{|x|^2}{2\kappa}$ into (2.23), we obtain that as $|x| \rightarrow \infty$,

$$\sum_{n=3}^{\infty} \bar{p}(n, \mathbf{0}, x) = \frac{\Gamma(\frac{d-2}{2})}{2\kappa\pi^{d/2}} \frac{1}{|x|^{d-2}} + O\left(\frac{1}{|x|^{d+2}}\right). \quad (2.24)$$

Let $E(n, x) = |p(n, \mathbf{0}, x) - \bar{p}(n, \mathbf{0}, x)|$. Similarly to the proof of [13, Lemma 1.5.2], we can verify that

$$\sum_{n=3}^{\infty} E(n, x) = O\left(\frac{1}{|x|^d}\right), \quad (2.25)$$

where the constant in $O(\cdot)$ does not depend on x but on μ .

Now from (2.24)-(2.25), we see the lemma holds immediately. \square

Lemma 2.4 *When $d = 2$, $a(x)$ is well-defined, and there exists a constant C_0 depending on μ such that as $|x| \rightarrow \infty$,*

$$a(x) = \frac{1}{\kappa\pi} \ln|x| + C_0 + O(|x|^{-2}), \quad (2.26)$$

where the constant in big O term only depends on μ .

Proof. We rewrite

$$a(x) = \sum_{3 \leq n \leq |x|^2/\kappa} [p(n, \mathbf{0}, \mathbf{0}) - p(n, \mathbf{0}, x)] + \sum_{n > |x|^2/\kappa} [p(n, \mathbf{0}, \mathbf{0}) - p(n, \mathbf{0}, x)]. \quad (2.27)$$

For any $n \geq 1$ and $x \in \mathbb{R}^2$, let

$$\bar{p}(n, \mathbf{0}, x) = \frac{1}{2\pi\kappa n} e^{-\frac{|x|^2}{2n\kappa}}.$$

It is easy to see that (2.16)-(2.21) are also true for $d = 2$.

Let $E(n, x) = |p(n, \mathbf{0}, x) - \bar{p}(n, \mathbf{0}, x)|$. Similarly to the proof of Theorem 1.2.1 in [13], we can check that

$$E(n, x) = O\left(\frac{1}{n^2}\right), \quad n \rightarrow \infty,$$

where the constant in $O(\cdot)$ does not depend on x but on μ .

Hence

$$\begin{aligned} p(n, \mathbf{0}, \mathbf{0}) &= \frac{1}{2\pi\kappa} \frac{1}{n} + O\left(\frac{1}{n^2}\right), \quad n \geq 3; \\ \sum_{3 \leq n \leq |x|^2/\kappa} p(n, \mathbf{0}, \mathbf{0}) &= \sum_{3 \leq n \leq |x|^2/\kappa} \frac{1}{2\pi\kappa} \frac{1}{n} + \sum_{3 \leq n \leq |x|^2/\kappa} O\left(\frac{1}{n^2}\right). \end{aligned}$$

Recall that

$$\sum_{3 \leq n \leq |x|^2/\kappa} \frac{1}{n} = 2 \ln|x| + \gamma - \ln \kappa - \frac{3}{2} + O(|x|^{-2}),$$

where γ is Euler's constant. So for some constant C_1 depending on μ ,

$$\sum_{3 \leq n \leq |x|^2/\kappa} p(n, \mathbf{0}, \mathbf{0}) = \frac{1}{\kappa\pi} \ln|x| + C_1 + O(|x|^{-2}).$$

By the martingale maximal inequality, there exist $\beta > 0, c > 0$ depending on μ such that for all n and all $s > 0$,

$$\mathbb{P} \left\{ \max_{0 \leq j \leq n} |S_j^\mu| \geq s\sqrt{n} \right\} \leq ce^{-\beta s^2}.$$

This implies that $\sum_{n < |x|} p(n, \mathbf{0}, x)$ decays faster than any power of $|x|$ as $|x| \rightarrow \infty$, particularly

$$\sum_{n < |x|} p(n, \mathbf{0}, x) = o(|x|^{-2}).$$

Similarly to the proof of Theorem 4.4.4 in [14], we can show that for some constant c depending on μ ,

$$\sum_{|x| \leq n \leq |x|^2/\kappa} |p(n, \mathbf{0}, x) - \bar{p}(n, \mathbf{0}, x)| \leq c|x|^{-2}.$$

Therefore, as $|x| \rightarrow \infty$,

$$\begin{aligned} \sum_{|x| \leq n \leq |x|^2/\kappa} p(n, \mathbf{0}, x) &= \sum_{|x| \leq n \leq |x|^2/\kappa} \bar{p}(n, \mathbf{0}, x) + O(|x|^{-2}) \\ &= \sum_{|x| \leq n \leq |x|^2/\kappa} \frac{1}{2\pi n \kappa} e^{-\frac{|x|^2}{2n\kappa}} + O(|x|^{-2}) \\ &= \frac{1}{2\pi \kappa} \int_{|x|}^{\frac{|x|^2}{\kappa}} \frac{1}{t} e^{-\frac{|x|^2}{2\kappa t}} dt + O(|x|^{-2}) \\ &= \frac{1}{2\pi \kappa} \int_1^\infty \frac{1}{y} e^{-y/2} dy + O(|x|^{-2}). \end{aligned}$$

So far we have proved that

$$\sum_{n \leq |x|^2/\kappa} [p(n, \mathbf{0}, \mathbf{0}) - p(n, \mathbf{0}, x)] = \frac{1}{\kappa\pi} \ln |x| + C' + O(|x|^{-2}), \quad |x| \rightarrow \infty, \quad (2.28)$$

holds for some constant C' depending on μ .

Likewise, consider the case when $n > |x|^2/\kappa$, similarly to the proof of Theorem 4.4.4 in [14], we can verify that

$$\sum_{n > |x|^2/\kappa} [p(n, \mathbf{0}, \mathbf{0}) - p(n, \mathbf{0}, x)] = \frac{1}{2\pi \kappa} \int_0^1 \frac{1}{y} (1 - e^{-y/2}) dy + O(|x|^{-2}), \quad |x| \rightarrow \infty. \quad (2.29)$$

By (2.28) and (2.29), we finish proving the lemma. \square

Let $G_D(\mathbf{0}, \cdot)$ be the Green function for the killed Brownian motion in \bar{D} starting at $\mathbf{0}$. Then $G_D(\mathbf{0}, z)$ is the unique harmonic function on $D \setminus \{\mathbf{0}\}$ such that $G_D(\mathbf{0}, z) \rightarrow 0$ as $z \rightarrow \partial D$. When $d \geq 3$, let $\varphi(z)$ be the harmonic function in $z \in D$ satisfying that

$$\begin{cases} \Delta \varphi(z) = 0, & z \in D, \\ \varphi(z) = -\frac{1}{4} \pi^{-d/2} \Gamma\left(\frac{d-2}{2}\right) \frac{1}{|z|^{d-2}}, & z \in \partial D. \end{cases} \quad (2.30)$$

It is well known that for $d \geq 3$,

$$G_D(\mathbf{0}, z) = \frac{1}{4} \pi^{-d/2} \Gamma\left(\frac{d-2}{2}\right) \frac{1}{|z|^{d-2}} + \varphi(z), \quad z \in \bar{D} \setminus \{\mathbf{0}\};$$

and $\frac{1}{4} \pi^{-d/2} \Gamma\left(\frac{d-2}{2}\right) \frac{1}{|z|^{d-2}}$ is harmonic in $z \in \mathbb{R}^d \setminus \{\mathbf{0}\}$, and thus $\varphi(z)$ can be extended to a harmonic function in the domain \bar{D}_+ for small enough $\delta > 0$.

Lemma 2.5 *There exists a constant $C > 0$ depending on μ but not on δ such that*

$$\left| \delta^2 G_\delta(\mathbf{0}, z) - \frac{2}{\kappa} G_D(\mathbf{0}, z) \right| \leq C\delta \quad (2.31)$$

holds uniformly in $z \in D$ with $|z| > \delta^{1/d}$.

Proof. When $d = 2$, proof of the lemma is similar to that of Lemma 5 in [10].

Assume $d \geq 3$, and δ is small enough such that $\mathbb{B}_{\delta^{1/d}}^d \subseteq D$. Let

$$H_\delta(z) = \delta^2 \sum_{n=0}^2 p_\delta(n, \mathbf{0}, z) + \delta^{2-d} a(z/\delta), \quad z \in \mathbb{R}^d \setminus \overline{\mathbb{B}_{2\delta}^d}.$$

Then by Lemma 2.3, we have that

$$H_\delta(z) = \frac{\Gamma(\frac{d-2}{2})}{2\kappa\pi^{d/2}} \frac{1}{|z|^{d-2}} + O(\delta^2/|z|^d), \quad z \in D \setminus \mathbb{B}_{\delta^{1/d}}^d.$$

Define the following function on \mathbb{R}^d :

$$e_\delta(z) = \begin{cases} H_\delta(z) - \delta^2 G_\delta(\mathbf{0}, z), & z \in \mathbb{R}^d \setminus \overline{\mathbb{B}_{2\delta}^d}, \\ \delta^{2-d} a(z/\delta) - \delta^2 \sum_{n=3}^{\infty} p_{D,\delta}(n, \mathbf{0}, z), & z \in \overline{\mathbb{B}_{2\delta}^d}. \end{cases}$$

Then by (2.10), we obtain that

$$\begin{cases} \Delta_\delta e_\delta(z) = 0, & z \in D, \\ e_\delta(z) = \frac{\Gamma(\frac{d-2}{2})}{2\kappa\pi^{d/2}} \frac{1}{|z|^{d-2}} + O(\delta^2/|z|^d), & z \in D_3. \end{cases} \quad (2.32)$$

Recall that $\varphi(z)$ can be extended to a harmonic function in the domain $\overline{D_+}$ for small enough $\delta > 0$. Therefore, for sufficiently small $\delta > 0$,

$$\begin{cases} \Delta_\delta \varphi(z) = 0, & z \in D, \\ \varphi(z) = -\frac{1}{4}\pi^{-d/2} \Gamma(\frac{d-2}{2}) \frac{1}{|z|^{d-2}} + O(\delta), & z \in D_3. \end{cases} \quad (2.33)$$

Combining with (2.32)-(2.33), we get that

$$\begin{cases} \Delta_\delta [e_\delta(z) + \frac{2}{\kappa}\varphi(z)] = 0, & z \in D, \\ e_\delta(z) + \frac{2}{\kappa}\varphi(z) = O(\delta), & z \in D_3. \end{cases}$$

Then the maximum principle for Δ_δ implies that

$$e_\delta(z) + \frac{2}{\kappa}\varphi(z) = O(\delta), \quad z \in D.$$

Therefore, when δ is small enough, for any $z \in D$ with $|z| > \delta^{1/d}$,

$$\begin{aligned} \delta^2 G_\delta(\mathbf{0}, z) &= H_\delta(z) - e_\delta(z) = \frac{2}{\kappa} \frac{\Gamma(\frac{d-2}{2})}{4\pi^{d/2}} \frac{1}{|z|^{d-2}} + \frac{2}{\kappa}\varphi(z) + O(\delta) \\ &= \frac{2}{\kappa} G_D(\mathbf{0}, z) + O(\delta). \end{aligned}$$

□

Proofs of the following Lemmas 2.6-2.7 are similar to those of Lemmas 6 and 8 in [10] respectively.

Case 2. $d \geq 3$. For any $\xi \in \overline{\mathbb{B}^d}$, write $\rho := \rho(r, \varphi_1, \varphi_2, \dots, \varphi_{d-1})$ for ξ in the spherical polar coordinates. Notice (2.36). Then

$$\begin{aligned}
& \Delta_\delta f(x + \ell \mathbf{n}_x) \\
&= \int_{\overline{\mathbb{B}^d}} [f(x + \ell \mathbf{n}_x + \delta \xi) - f(x + \ell \mathbf{n}_x)] \, d\xi \\
&= \int_0^1 \int_0^{2\pi} \int_0^\pi \cdots \int_0^\pi [f(x + \ell \mathbf{n}_x + \delta \rho) - f(x + \ell \mathbf{n}_x)] J_d \, d\mu(r, \varphi_1, \dots, \varphi_{d-1}) \\
&= \int_{[0, \ell/\delta]} \int_0^{2\pi} \int_0^\pi \cdots \int_0^\pi [f(x + \ell \mathbf{n}_x + \delta \rho) - f(x + \ell \mathbf{n}_x)] J_d \, d\mu(r, \varphi_1, \dots, \varphi_{d-1}) \\
&\quad + \int_{[\ell/\delta, 1]} \int_0^{2\pi} \int_0^\pi \cdots \int_0^\pi [f(x + \ell \mathbf{n}_x + \delta \rho) - f(x + \ell \mathbf{n}_x)] J_d \, d\mu(r, \varphi_1, \dots, \varphi_{d-1}) \\
&= \int_{[\ell/\delta, 1]} \int_0^{2\pi} \int_0^\pi \cdots \int_0^\pi [f(x + \ell \mathbf{n}_x + \delta \rho) - f(x + \ell \mathbf{n}_x)] J_d \, d\mu(r, \varphi_1, \dots, \varphi_{d-1}) \\
&= \int_{[\ell/\delta, 1]} \int_0^{2\pi} \int_0^\pi \cdots \int_0^{\pi/2 + \arcsin(\frac{\ell}{\delta r})} [f(x + \ell \mathbf{n}_x + \delta \rho) - f(x + \ell \mathbf{n}_x)] J_d \, d\mu(r, \varphi_1, \dots, \varphi_{d-1}) \\
&\quad + \int_{[\ell/\delta, 1]} \int_0^{2\pi} \int_0^\pi \cdots \int_0^{\arccos(\frac{\ell}{\delta r})} [f(x + \ell \mathbf{n}_x - \delta \rho) - f(x + \ell \mathbf{n}_x)] J_d \, d\mu(r, \varphi_1, \dots, \varphi_{d-1}) \\
&= I_1(x, \ell) + I_2(x, \ell). \tag{2.37}
\end{aligned}$$

The smoothness and boundedness of ∂D implies that there are positive constants C_1, C_2, \dots, C_d such that for any $x \in \partial D$,

$$\partial D \cap \overline{\mathbb{B}^d(x, \delta)} \subset [-C_1\delta, C_1\delta] \cdot \mathbf{t}_x^1 + [-C_2\delta, C_2\delta] \cdot \mathbf{t}_x^2 + \cdots + [-C_d\delta^2, C_d\delta^2] \cdot \mathbf{n}_x,$$

where $\{\mathbf{t}_x^1, \mathbf{t}_x^2, \dots, \mathbf{t}_x^{d-1}\}_{x \in \partial D}$ is a moving tangent orthonormal frame of ∂D .

Since f has a C^2 -extension, which is still denoted by f , to \overline{D}_+ , by the Taylor expansion of f at $x \in \partial D$ with respect to coordinate directions \mathbf{n}_x and $\mathbf{t}_x^1, \dots, \mathbf{t}_x^{d-1}$, we get that

$$I_1(x, \ell) = \frac{\partial f(x)}{\partial \mathbf{n}_x} \int_{[\ell/\delta, 1]} \int_0^{2\pi} \int_0^\pi \cdots \int_0^{\pi/2 + \arcsin(\frac{\ell}{\delta r})} \delta r \cos(\varphi_1) J_d \, d\mu(r, \varphi_1, \dots, \varphi_{d-1}) + O(\delta^2).$$

Recall

$$\int_0^{2\pi} \int_0^\pi \cdots \int_0^\pi (\sin \varphi_2)^{d-3} \cdots \sin \varphi_{d-2} \, d\varphi_2 \cdots d\varphi_{d-2} d\varphi_{d-1} = \frac{2\pi^{(d-1)/2}}{\Gamma(\frac{d-1}{2})}.$$

Hence by (2.36),

$$\begin{aligned}
& I_1(x, \ell) \\
&= \frac{\partial f(x)}{\partial \mathbf{n}_x} \int_{[\ell/\delta, 1]} \int_0^{\pi/2 + \arcsin(\frac{\ell}{\delta r})} \frac{2\pi^{(d-1)/2}}{\Gamma(\frac{d-1}{2})} \frac{\Gamma(d/2)}{2\pi^{d/2} r^{d-1}} \delta r^d \cos(\varphi_1) \sin^{d-2}(\varphi_1) \, d\varphi_1 d\nu(r) + O(\delta^2) \\
&= \frac{\partial f(x)}{\partial \mathbf{n}_x} \int_{[\ell/\delta, 1]} \frac{\Gamma(d/2)}{\sqrt{\pi} \Gamma(\frac{d-1}{2})} \frac{(\delta^2 r^2 - \ell^2)^{(d-1)/2}}{(d-1)(\delta r)^{d-2}} \, d\nu(r) + O(\delta^2).
\end{aligned}$$

Likewise, we have that

$$\begin{aligned}
I_2(x, \ell) &= \frac{\partial f(x)}{\partial \mathbf{n}_x} \int_{[\ell/\delta, 1]} \int_0^\pi \cdots \int_0^{\arccos(\frac{\ell}{\delta r})} -\ell J_d \frac{\Gamma(d/2)}{2\pi^{d/2} (\delta r)^{d-1}} \, d\varphi_1 \cdots d\varphi_{d-2} d\nu(r) + O(\delta^2) \\
&= \frac{\partial f(x)}{\partial \mathbf{n}_x} \int_{[\ell/\delta, 1]} \int_0^{\arccos(\frac{\ell}{\delta r})} -\ell \frac{\Gamma(d/2)}{\sqrt{\pi} \Gamma(\frac{d-1}{2})} \sin^{d-2}(\varphi_1) \, d\varphi_1 d\nu(r) + O(\delta^2).
\end{aligned}$$

From [2],

$$\int \sin^{d-2}(\varphi_1) d\varphi_1 = -\cos(\varphi_1) {}_2F_1\left(\frac{1}{2}, \frac{3-d}{2}; \frac{3}{2}; \cos^2(\varphi_1)\right) + C, \quad 0 \leq \varphi_1 \leq \pi,$$

$${}_2F_1\left(\frac{1}{2}, \frac{3-d}{2}; \frac{3}{2}; 1\right) = \frac{\sqrt{\pi}\Gamma(\frac{d-1}{2})}{2\Gamma(d/2)}.$$

Therefore, the lemma follows by a simple computation. \square

Lemma 2.9 Fix $a \in (0, \infty)$ and $\theta \in [\frac{1}{8}\pi, \pi]$. Let

$$\Gamma = \Gamma(a, \theta) = \{x = (x_1, \dots, x_d) \in \partial\mathbb{B}_a^d : x_d \geq a \cos \theta\}.$$

Then there exists a constant C , depending on μ , but being independent of a and θ , such that when δ is small enough,

$$\omega_\delta(u, \Gamma; \mathbb{B}_a^d) \geq C > 0 \quad \text{for all } u \in \mathbb{B}_{a/2}^d.$$

Proof. The proof is similar to that of Lemma 2.10 in [3]. Write $x = (x_1, \dots, x_d)$ in the d -dimensional spherical polar coordinates transform given by (2.34). For any $\rho \in (0, 1/4)$, let

$$\Pi_1(\rho) = \{x \in \mathbb{R}^d : r = |x| \in (1 - \rho, 1 + \rho), \varphi_1 \in [0, \pi/8]\},$$

$$\Pi_2(\rho) = \{x \in \mathbb{R}^d : r = |x| \in (1 - \rho, 1 + \rho), \varphi_1 \in (\pi/8, \pi]\};$$

and $h_\rho(\cdot)$ be the continuous harmonic measure of $\{x \in \mathbb{R}^d : |x| = 1 + \rho, \varphi_1 \in [0, \frac{1}{9}\pi]\}$, and

$$c_1 := \frac{1}{2} \liminf_{\rho \rightarrow 0} \min_{|x| \leq 1/2} \{h_\rho(x)\} > 0.$$

Choose ρ small enough such that

$$\min_{|x| \leq \frac{1}{2} + \rho} \{h_\rho(x)\} \geq 3c_1/2 \quad \text{and} \quad \max_{x \in \Pi_2(\rho)} \{h_\rho(x)\} \leq c_1/2.$$

Then choose constant $c_2 > 0$ sufficiently small such that $c_2(1 + \rho)^2 \leq \frac{1}{2}c_1$. Let

$$\phi_0(x) := h_\rho(x) - c_1 + c_2|x|^2, \quad x \in \overline{\mathbb{B}_{1+\rho}^d}.$$

Then smooth function $\phi_0 : \mathbb{B}_{1+\rho}^d \rightarrow \mathbb{R}$ satisfies that

- (i) $\phi_0(x) \leq 1$ for all $x \in \Pi_1(\rho)$;
- (ii) $\phi_0(x) \leq 0$ for any $x \in \Pi_2(\rho)$;
- (iii) ϕ_0 is subharmonic, and $\Delta\phi_0 \geq 2dc_2$;
- (iv) $\phi_0(x) \geq c_1/2$ for all $x \in \mathbb{B}_{1/2+\rho}^d$.

Let

$$\phi(x) = \phi_0\left(\frac{x}{a}\right), \quad x \in \overline{\mathbb{B}_a^d}.$$

Then $\phi \leq 1$ on Γ and $\phi \leq 0$ on $\partial\mathbb{B}_a^d \setminus \Gamma$.

Notice $\limsup_{\delta \rightarrow 0} \sup_{x \in \overline{\mathbb{B}_a^d}} |(\frac{\kappa}{2}\Delta\phi - \Delta_\delta\phi)(x)| = 0$. By (iii), for small enough δ , ϕ is Δ_δ -subharmonic in \mathbb{B}_a^d .

By the maximum principle, for some constant $C > 0$ which depends on μ but is independent of a and θ , when δ is sufficiently small,

$$\omega_\delta(x, \Gamma; \mathbb{B}_a^d) \geq \phi(x) \geq C > 0, \quad x \in \mathbb{B}_{a/2}^d.$$

\square

Lemma 2.10 *There exist two constants $\beta > 0$ and $C > 0$ depending on μ such that for any simply-connected bounded regular domain \mathcal{D} , $u \in \mathcal{D}$ and $V \subseteq \partial\mathcal{D}$, and sufficiently small $\delta > 0$,*

$$\mathbb{P}^{u,\delta} \left(\overline{\delta S_{T_{\mathcal{D}}}^{\mu}} \in V \right) \leq C \left[\frac{\text{dist}(u, \partial\mathcal{D})}{\text{dist}_{\mathcal{D}}(u; V)} \right]^{\beta},$$

where $\overline{\delta S_{T_{\mathcal{D}}}^{\mu}}$ is the orthogonal projection of $\delta S_{T_{\mathcal{D}}}^{\mu}$ onto $\partial\mathcal{D}$, and

$$\text{dist}_{\mathcal{D}}(u; V) = \inf \{ R > 0 : u \text{ and } V \text{ are connected in } \mathcal{D} \cap \mathbb{B}^d(u, R) \}.$$

Proof. The proof is similar to that of Proposition 2.11 in [3]. Let $a = \text{dist}(u, \partial\mathcal{D})$ and $r = \text{dist}_{\mathcal{D}}(u; V)$. Recall that $\omega_{\delta}(u, V; \mathcal{D})$ is equal to the probability that the first exit point of $\{\delta S_n^{\mu}\}_{n \geq 0}$ ($\delta S_0^{\mu} = u$) from \mathcal{D} whose orthogonal projection onto $\partial\mathcal{D}$ is in V . By Lemma 2.9, without loss of generality, we assume $a < r/2$. It is easy to show that for each $a \leq r' \leq \frac{r}{2}$ the probability to cross the annulus $\mathbb{B}^d(u, 2r') \setminus \mathbb{B}^d(u, r')$ inside \mathcal{D} without touching its boundary is bounded above by some constant $p > 0$ depending on μ that does not depend on r' and \mathcal{D} . Hence,

$$\omega_{\delta}(u, V; \mathcal{D}) \leq p^{\log_2(r/a)-1} = p^{-1} \cdot (a/r)^{-\log_2 p},$$

so the lemma holds true with the exponent $\beta = -\log_2 p$. \square

Note Lemma 2.5. The proof of the following Lemma 2.11 is similar to that of [10] Proposition 2.

Lemma 2.11 *Given any $\epsilon \in (0, 1/2)$. Then as $\delta \rightarrow 0$, for any $\ell \in [0, \delta]$ and $x \in \partial D$,*

$$\delta^2 G_{\delta}(\mathbf{0}, z) - \frac{2}{\kappa} G_D(\mathbf{0}, z) - \frac{2}{\kappa} H_D(\mathbf{0}, x) \mathbb{E}^{\ell} \left[V \left(S_{T_{\mathbb{H}^d}}^{\mu} \right) \right] = O(\delta^{1+\epsilon\beta}) + O(\delta^{2-2\epsilon}),$$

where $z = x + \ell \mathbf{n}_x \in D_2$ and the big O terms depend on D and μ .

Lemma 2.12 *Let f be the harmonic function in D with boundary values given by the smooth function g . Then (1.4) holds for C^{∞} -function $\rho_D(z)$ specified in (1.8).*

Proof. Let $h(\cdot)$ be the harmonic function on D with boundary values given by the smooth function $H_D(\mathbf{0}, \cdot)$. Then $h \in C^{\infty}(D)$. Since $\Delta f(z) = 0$, $z \in D$, by the divergence theorem,

$$\begin{aligned} \int_{\partial D} H_D(\mathbf{0}, z) \frac{\partial f}{\partial \mathbf{n}_z}(z) |dz| &= - \int_D \nabla h(z) \cdot \nabla f(z) + h(z) \cdot \Delta f(z) \, dz \\ &= - \int_D \nabla h(z) \cdot \nabla f(z) \, dz. \end{aligned}$$

Moreover, note $\Delta h(z) = 0$, $z \in D$, by the divergence theorem again,

$$\begin{aligned} \int_{\partial D} H_D(\mathbf{0}, z) \frac{\partial f}{\partial \mathbf{n}_z}(z) |dz| &= - \int_D \Delta h(z) \cdot f(z) + \nabla h(z) \cdot \nabla f(z) \, dz \\ &= \int_{\partial D} g(z) \frac{\partial h}{\partial \mathbf{n}_z}(z) |dz|. \end{aligned}$$

The mean value and regularity property of harmonic function $h(\cdot)$ implies that for any $z \in \partial D$,

$$\begin{aligned} \frac{\partial h}{\partial \mathbf{n}_z}(z) &= \lim_{\delta \rightarrow 0} \frac{1}{\delta} [h(z + \delta \mathbf{n}_z) - h(z)] \\ &= \lim_{\delta \rightarrow 0} \frac{1}{\delta} \left\{ \int_{\partial D} H_D(\mathbf{0}, \zeta) H_D(z + \delta \mathbf{n}_z, \zeta) |d\zeta| - H_D(\mathbf{0}, z) \right\}. \end{aligned}$$

For any $z \in \partial D$, let

$$\rho_D(z) = \frac{\partial h}{\partial \mathbf{n}_z}(z).$$

Thus (1.4) holds for C^{∞} -function $\rho_D(z)$ specified in (1.8). \square

2.3 Proof of Theorem 1.2

Recall (2.4) and (2.7). By Lemmas 2.2 and 2.7, we get that

$$\begin{aligned} f_\delta(\mathbf{0}) - f(\mathbf{0}) &= \int_{D_2} G_\delta(\mathbf{0}, z) \Delta_\delta f(z) \, dz \\ &= (1 + O(\delta)) \int_{\partial D} \int_0^\delta G_\delta(\mathbf{0}, x + \ell \mathbf{n}_x) \Delta_\delta f(x + \ell \mathbf{n}_x) \, d\ell |dx|. \end{aligned}$$

By Lemma 2.8,

$$\begin{aligned} \Phi(\delta) &:= \int_{\partial D} \int_0^\delta G_\delta(\mathbf{0}, x + \ell \mathbf{n}_x) \Delta_\delta f(x + \ell \mathbf{n}_x) \, d\ell |dx| \\ &= \int_{\partial D} \frac{\partial f(x)}{\partial \mathbf{n}_x} \int_0^\delta G_\delta(\mathbf{0}, x + \ell \mathbf{n}_x) [h^\mu(\ell, \delta) + O(\delta^2)] \, d\ell |dx|. \end{aligned}$$

Combining with Lemma 2.11, we have that

$$\begin{aligned} \Phi(\delta) &= \int_{\partial D} \frac{\partial f(x)}{\partial \mathbf{n}_x} \int_0^\delta \frac{2}{\kappa \delta^2} \left[G_D(\mathbf{0}, x + \ell \mathbf{n}_x) + H_D(\mathbf{0}, x) \mathbb{E}^\ell \left[V \left(S_{T_{\mathbb{H}^d}}^\mu \right) \right] + o(\delta) \right] \\ &\quad \times [h^\mu(\ell, \delta) + O(\delta^2)] \, d\ell |dx|. \end{aligned}$$

By Lemma 2.6,

$$\begin{aligned} \Phi(\delta) &= \int_{\partial D} \frac{\partial f(x)}{\partial \mathbf{n}_x} \int_0^\delta \frac{2}{\kappa \delta^2} \left[\ell H_D(\mathbf{0}, x) + H_D(\mathbf{0}, x) \mathbb{E}^\ell \left[V \left(S_{T_{\mathbb{H}^d}}^\mu \right) \right] + o(\delta) \right] \\ &\quad \times [h^\mu(\ell, \delta) + O(\delta^2)] \, d\ell |dx|. \end{aligned}$$

A straightforward calculation gives that

$$\Phi(\delta) = c_\mu \delta \int_{\partial D} \frac{\partial f(x)}{\partial \mathbf{n}_x} H_D(\mathbf{0}, x) \, |dx| + o(\delta),$$

where c_μ is given by (1.7). Therefore

$$f_\delta(\mathbf{0}) - f(\mathbf{0}) = \delta c_\mu \int_{\partial D} \frac{\partial f(x)}{\partial \mathbf{n}_x} H_D(\mathbf{0}, x) \, |dx| + O(\delta^2),$$

combining with Lemma 2.12, then (1.3), (1.4) and (1.8) hold. So far we have completed proving Theorem 1.2. \square

3 Several examples for Theorem 1.2

In this section, we consider a class of random walks $\{\delta S_n^\mu\}_{n \geq 0}$ for Theorem 1.2 such that μ has the form specified by (1.5) with

$$\nu(r) := \nu([0, r]) = 1 - \left[1 - r^{1/\alpha} \right]^{1/\beta}, \quad r \in [0, 1], \quad (3.1)$$

where $\alpha \geq 0$ and $\beta > 0$. A tedious and straightforward calculation shows that κ in (1.1) is given by

$$\kappa = \kappa(d, \alpha, \beta) = \frac{\Gamma(d/2) \Gamma(2 + 2\alpha) \Gamma(1/\beta)}{2\beta(1 + 2\alpha) \Gamma(d/2 + 1) \Gamma(1/\beta + 2\alpha + 1)}. \quad (3.2)$$

Let $d = 2, \beta = 1, \alpha = \frac{1}{2}$, then $\nu(r) = r^2, r \in [0, 1], \kappa = \frac{1}{4}$, and the random walk $\{\delta S_n^\mu\}_{n \geq 0}$ is exactly the case of continuous-state random walk in [10].

The following corollary is a consequence of some tedious but straightforward calculus.

Corollary 3.1 In (3.1), let d, β, α be as follows.

(i) When $d = 2, \beta = 1$ and $\alpha = 0$, i.e. $\nu(r) = I_{\{r=1\}}$, $\kappa = \frac{1}{2}$, then step-distribution of random walk $\{S_n^\mu\}_{n \geq 0}$ is the uniform distribution on the unit circle, and

$$c_\mu = \frac{4}{9\pi} + \frac{4}{\pi} \int_0^1 \left[\sqrt{1-\ell^2} - \ell \arccos(\ell) \right] \mathbb{E}^\ell \left[V \left(S_{T_{\mathbb{H}^2}}^\mu \right) \right] d\ell.$$

(ii) When $d = 3, \beta = 1$ and $\alpha = 0$, i.e. $\nu(r) = I_{\{r=1\}}$, $\kappa = \frac{1}{3}$, then step-distribution of random walk $\{S_n^\mu\}_{n \geq 0}$ is the uniform distribution on the unit spherical surface in \mathbb{R}^3 , and

$$c_\mu = \frac{1}{8} + \int_0^1 \left(\frac{3}{2} - 3\ell + \frac{3\ell^2}{2} \right) \mathbb{E}^\ell \left[V \left(S_{T_{\mathbb{H}^3}}^\mu \right) \right] d\ell.$$

(iii) When $d = 3, \beta = 1$ and $\alpha = 1/3$, i.e. $\nu(r) = r^3$, $r \in [0, 1]$, $\kappa = \frac{1}{5}$, then step-distribution of random walk $\{S_n^\mu\}_{n \geq 0}$ is the uniform distribution on the closed 3-dimensional unit ball, and

$$c_\mu = \frac{5}{48} + \frac{5}{8} \int_0^1 (3 + \ell)(1 - \ell)^3 \mathbb{E}^\ell \left[V \left(S_{T_{\mathbb{H}^3}}^\mu \right) \right] d\ell.$$

(iv) When $d = 4, \beta = 1$ and $\alpha = 0$, i.e. $\nu(r) = I_{\{r=1\}}$, $\kappa = \frac{1}{4}$, then step-distribution of random walk $\{S_n^\mu\}_{n \geq 0}$ is the uniform distribution on the unit spherical surface in \mathbb{R}^4 , and

$$c_\mu = \frac{16}{45\pi} + \int_0^1 \left(\frac{16}{3\pi} (1 - \ell^2)^{3/2} - \frac{8\ell}{\pi} \arccos(\ell) + \frac{8\ell^2}{\pi} \sqrt{1 - \ell^2} \right) \mathbb{E}^\ell \left[V \left(S_{T_{\mathbb{H}^4}}^\mu \right) \right] d\ell.$$

(v) When $d = 4, \beta = 1$ and $\alpha = 1/4$, i.e. $\nu(r) = r^4$, $r \in [0, 1]$, $\kappa = \frac{1}{6}$, then step-distribution of random walk $\{S_n^\mu\}_{n \geq 0}$ is the uniform distribution on the closed 4-dimensional unit ball, and

$$c_\mu = \frac{32}{105\pi} + \int_0^1 \left(\frac{32}{5\pi} (1 - \ell^2)^{5/2} - \frac{4}{\pi} \ell^2 (2\ell^2 - 5) \sqrt{1 - \ell^2} - \frac{12}{\pi} \ell \arccos(\ell) \right) \mathbb{E}^\ell \left[V \left(S_{T_{\mathbb{H}^4}}^\mu \right) \right] d\ell.$$

Acknowledgements. We would like to express our sincere appreciation to the reviewers for their insightful comments and we are grateful to Dr. Jianping Jiang for many helpful conversations.

References

- [1] S. Axler, P. Bourdon and R. Wade. (2001). *Harmonic function theory*. Springer Science and Business Media, New York.
- [2] Y. A. Brychkov. (2008). *Handbook of special functions. Derivatives, integrals, series and other formulas*. CRC Press, Boca Raton, FL.
- [3] D. Chelkak and S. Smirnov. (2011). Discrete complex analysis on isoradial graphs. *Adv. Math.* **228(3)**, 1590-1630.
- [4] Y. Dai. (2017). The exit distribution for smart kinetic walk with symmetric transition probability. *J. Stat. Phys.* **166(6)**, 1455-1463.
- [5] J. Garnett and D. Marshall. (2005). *Harmonic measure*. Cambridge Univ. Press, Cambridge.
- [6] G. Gasper and M. Rahman. (2004). *Basic hypergeometric series* (2nd edition). Encyclopedia of Mathematics and its Applications **96**. Cambridge Univ. Press, Cambridge.
- [7] D. Gilbarg and N. S. Trudinger. (2001). *Elliptic partial differential equations of second order*. Reprint of the 1998 edition. Classics in Mathematics. Springer-Verlag, Berlin.
- [8] B. D. Hughes. (1995). *Random walks and random environments* (Volume 1). Oxford Univ. Press, Oxford.

- [9] D. Jerison. (1990). Regularity of the Poisson kernel and free boundary problems. *Colloq. Math.* **60(2)**, 547-568.
- [10] J. Jiang and T. Kennedy. (2017). The difference between a discrete and continuous harmonic measure. *J. Theoret. Probab.* **30(4)**, 1424-1444.
- [11] T. Kennedy. (2016). The first order correction to the exit distribution for some random walks. *J. Stat. Phys.* **164(1)**, 174-189.
- [12] I. Karatzas and S. E. Shreve. (1991). *Brownian motion and stochastic calculus* (2nd edition). Graduate Texts in Mathematics, **113**. Springer-Verlag, New York.
- [13] G. F. Lawler. (1991). *Intersections of random walks*. Birkhäuser, Boston.
- [14] G. F. Lawler and V. Limic. (2010). *Random walk: a modern introduction*. Cambridge Univ. Press, Cambridge.
- [15] P. Möters and Y. Peres. (2010). *Brownian motion*. Cambridge Univ. Press, Cambridge.
- [16] G. N. Watson. (1944). *A treatise on the theory of Bessel functions* (2nd edition). Cambridge Univ. Press, Cambridge.
- [17] J. Zabczyk. (1970). Sur la théorie semi-classique du potentiel pour les processus à accroissements indépendants. *Studia Math.* **35**, 227-247.